APPENDIX

Pediatric Crohn's Disease Activity Index (PCDAI) (4)			
History (Recall, 1 week)			
Abdominal pain			Score
0 = None	5 = Mild: brief, does not interfere with activities	10 = Moderate/severe: daily, longer lasting, affects activities, nocturnal	_____
Patient functioning, general well-being			Score
0 = No limitation of activities, well	5 = Occasional difficulty in maintaining age appropriate activities, below par	10 = Frequent limitation of activity, very poor	_____
Stools (per day)			Score
0 = 0–1 liquid stools, no blood	5 = Up to 2 semi-formed with small blood, or 2–5 liquid	10 = Gross bleeding, or ≥ 6 liquid, or nocturnal diarrhea	_____
Laboratory			
Hematocrit			Score
< 10 years:	11–14 years (male):		
0 = > 33% 2.5 = 28–32% 5 = < 28%	0 = ≥35% 2.5 = 30–34%	5 = < 30%	
11–19 years (female):	15–19 years (male):		
0 = ≥34% 2.5 = 29–33% 5 = < 29%	0 = ≥37% 2.5 = 32–36%	5 = < 32	_____
Erythrocyte sedimentation rate			Score
0 = < 20 mm/h	2.5 = 20 – 50 mm/h	5 = > 50 mm/h	_____
Albumin			Score
0 = ≥3.5 g/dl	5 = 3.1–3.4 g/dl	10 = ≤ 3.0 g/dl	_____
Examination			
Weight			Score
0 = Weight gain or voluntary weight stable/loss	5 = Involuntary weight stable, weight loss 1–9%	10 = Weight loss ≥ 10%	_____
Height			Score
If completed at diagnosis:			
0 = < 1 channel decrease	5 = ≥ 1 < 2 channel decrease	10 = > 2 channel decrease	
If completed after diagnosis:			
0 = Height velocity ≥ – 1 s.d.	5 = Height velocity < – 1 s.d., > – 2 s.d.	10 = Height velocity ≤ – 2 s.d.	_____
Abdominal examination			Score
0 = No tenderness, no mass	5 = Tenderness, or mass without tenderness	10 = Tenderness, involuntary guarding, definite mass	_____
Perirectal disease			Score
0 = None, asymptomatic tags	5 = 1–2 indolent fistula, scant drainage, no tenderness	10 = Active fistula, drainage, tenderness, or abscess	_____
Extra-intestinal manifestations			Score
(Fever ≥38.5 °C for 3 days over past week, definite arthritis, uveitis, Erythema nodosum, Pyoderma gangrenosum)			
0 = None	5 = One	10 = ≥ Two	_____
		Total score:	