

**Practice Guidelines for Perioperative Blood Management: An Updated Report**  
*American Society of Anesthesiologists*

**Bibliography in Alphabetical Order**

1. Abdelmalak BB, Cata JP, Bonilla A, You J, Kopyeva T, Vogel JD, Campbell S, Sessler DI: Intraoperative tissue oxygenation and postoperative outcomes after major non-cardiac surgery: an observational study. *Br J Anaesth* 2013; 110:241-249
2. Ahn SW, Shim JK, Youn YN, Song JW, Yang SY, Chung SC, Kwak YL: Effect of tranexamic acid on transfusion requirement in dual antiplatelet-treated anemic patients undergoing off-pump coronary artery bypass graft surgery. *Circ J* 2012; 76:96-101
3. Ak K, Isbir CS, Tetik S, Atalan N, Tekeli A, Aljodi M, Civelek A, Arsan S: Thromboelastography-based transfusion algorithm reduces blood product use after elective CABG: a prospective randomized study. *J Card Surg* 2009; 24:404-410
4. Aksoy MC, Tokgozoglu AM: Erythropoietin for autologous blood donation in total hip arthroplasty patients. *Arch Orthop Trauma Surg* 2001; 121:162-165
5. Alanay A, Acaroglu E, Ozdemir O, Ercelen O, Bulutcu E, Surat A: Effects of deamino-8-D-arginin vasopressin on blood loss and coagulation factors in scoliosis surgery. A double-blind randomized clinical trial. *Spine* 1999; 24:877-882
6. Almeida MF: Preoperative normovolemic hemodilution in aesthetic plastic surgery. *Aesthetic Plast Surg* 1999; 23:445-449
7. Amanor-Boadu SD, Malomo A, Komolafe EO, Adeolu AA, Abdullahi A, Shokunbi MT: Acute isovolaemic haemodilution in two Jehovah's Witnesses presenting for major intracranial surgery. *Afr J Med Sci* 2002; 31:79-81
8. Andersen ND, Bhattacharya SD, Williams JB, Fosbol EL, Lockhart EL, Patel MB, Gaca JG, Welsby IJ, Hughes GC: Intraoperative use of low-dose recombinant activated factor VII during thoracic aortic operations. *Ann Thorac Surg* 2012; 93:1921-1928
9. Andreasen JB, Hvas AM, Christiansen K, Ravn HB: Can RoTEM® analysis be applied for haemostatic monitoring in paediatric congenital heart surgery? *Cardiol Young* 2011; 21:684-691
10. Andreasen JJ, Dethlefsen C, Modrau IS, Baech J, Schonheyder HC, Moeller JK, Johnsen SP: Storage time of allogeneic red blood cells is associated with risk of severe postoperative infection after coronary artery bypass grafting. *Eur J Cardiothorac Surg* 2011; 39:329-334
11. Andreasen JJ, Nielsen C: Prophylactic tranexamic acid in elective, primary coronary artery bypass surgery using cardiopulmonary bypass. *Eur J Cardiothorac Surg* 2004; 26:311-317
12. Andrews CM, Lane DW, Bradley JG: Iron pre-load for major joint replacement. *Transfus Med* 1997; 7:281-286
13. Avall A, Hyllner M, Bengtson JP, Carlsson L, Bengtsson A: Greater increase in cytokine concentration after salvage with filtered whole blood than with washed red cells, but no difference in postoperative hemoglobin recovery. *Transfusion* 1999; 39:271-276
14. Avall A, Hyllner M, Bengtson JP, Carlsson L, Bengtsson A: Postoperative inflammatory response after autologous and allogeneic blood transfusion. *Anesthesiology* 1997; 87:511-516

15. Avidan MS, Alcock EL, Da Fonseca J, Ponte J, Desai JB, Despotis GJ, Hunt BJ: Comparison of structured use of routine laboratory tests or near-patient assessment with clinical judgement in the management of bleeding after cardiac surgery. *Br J Anaesth* 2004; 92:178-186
16. Axford TC, Dearani JA, Ragno G, MacGregor H, Patel MA, Valeri CR, Khuri SF: Safety and therapeutic effectiveness of reinfused shed blood after open heart surgery. *Ann Thorac Surg* 1994; 57:615-622
17. Ayers DC, Murray DG, Duerr DM: Blood salvage after total hip arthroplasty. *J Bone Joint Surg Am* 1995; 77:1347-1351
18. Bacher A, Mayer N, Rajek AM, Haider W: Acute normovolaemic haemodilution does not aggravate gastric mucosal acidosis during cardiac surgery. *Intensive Care Med* 1998; 24:313-321
19. Badner NH, Nicolaou G, Clarke CF, Forbes TL: Use of spinal near-infrared spectroscopy for monitoring spinal cord perfusion during endovascular thoracic aortic repairs. *J Cardiothorac Vasc Anesth* 2011; 25:316-319
20. Banbury MK, Brizzio ME, Rajeswaran J, Lytle BW, Blackstone EH: Transfusion increases the risk of postoperative infection after cardiac surgery. *J Am Coll Surg* 2006; 202:131-138
21. Barnette RE, Wendling WW, Schweiger JW, Brister NW, Schartel SA, Chen D, Shuman CA, McClurken JB, Jeevanandam V: Intravenous vitamin K1 prior to orthotopic heart transplantation: effects in vivo and in vitro. *Acta Anaesthesiol Scand* 1997; 41:78-83
22. Bartels C, Bechtel JV, Winkler C, Horsch S: Intraoperative autotransfusion in aortic surgery: comparison of whole blood autotransfusion versus cell separation. *J Vasc Surg* 1996; 24:102-108
23. Bartels C, Claeys L, Ktenidis K, Nigbur H, Horsch S: Intraoperative whole blood autotransfusion during venous thrombectomy. *J Cardiovasc Surg* 1994; 35:109-114
24. Beholz S, Liu J, Thoelke R, Spiess C, Konertz W: Use of desmopressin and erythropoietin in an anaemic Jehovah's Witness patient with severely impaired coagulation capacity undergoing stentless aortic valve replacement. *Perfusion* 2001; 16:485-489
25. Benavides S, Nicol K, Koranyi K, Nahata MC: Yersinia septic shock following an autologous transfusion in a pediatric patient. *Transfus Apher Sci* 2003; 28:19-23
26. Benli IT, Akalin S, Duman E, Citak M, Kis M: The results of intraoperative autotransfusion in orthopaedic surgery. *Bull Hosp Jt Dis* 1999; 58:184-187
27. Bennett J, Haynes S, Torella F, Grainger H, McCollum C: Acute normovolemic hemodilution in moderate blood loss surgery: a randomized controlled trial. *Transfusion* 2006; 46:1097-1103
28. Bennett SR: Perioperative autologous blood transfusion in elective total hip prosthesis operations. *Ann R Coll Surg Engl* 1994; 76:95-98
29. Bennett-Guerrero E, Sorohan JG, Canada AT, Ayuso L, Newman MF, Reves JG, Mythen MG:  $\epsilon$ -aminocaproic acid plasma levels during cardiopulmonary bypass. *Anesth Analg* 1997; 85:248-251
30. Benoist S, Panis Y, Pannegeon V, Alves A, Valleur P: Predictive factors for perioperative blood transfusions in rectal resection for cancer: A multivariate analysis of a group of 212 patients. *Surgery* 2001; 129:433-439

31. Benoni G, Fredin H: Fibrinolytic inhibition with tranexamic acid reduces blood loss and blood transfusion after knee arthroplasty: a prospective, randomised, double-blind study of 86 patients. *J Bone Joint Surg Br* 1996; 78:434-440
32. Benoni G, Lethagen S, Nilsson P, Fredin H: Tranexamic acid, given at the end of the operation, does not reduce postoperative blood loss in hip arthroplasty. *Acta Orthop Scand* 2000; 71:250-254
33. Bhardwaj M, Bunsell R: Beriplex P/N: an alternative to fresh frozen plasma in severe haemorrhage. *Anaesthesia* 2007; 62:832-834
34. Bhatt K, Cherian S, Agarwal R, Jose S, Cherian KM: Perioperative management of sickle cell disease in paediatric cardiac surgery. *Anaesth Intensive Care* 2007; 35:792-795
35. Biesma DH, Marx JJ, Kraaijenhagen RJ, Franke W, Messinger D, van de Wiel A: Lower homologous blood requirement in autologous blood donors after treatment with recombinant human erythropoietin. *Lancet* 1994; 344:367-370
36. Bilgin YM, van de Watering LM, Eijnsman L, Versteegh MI, Brand R, van Oers MH, Brand A: Double-blind, randomized controlled trial on the effect of leukoreduced erythrocyt transfusions in cardiac valve surgery. *Circulation* 2004; 109:2755-2760
37. Billote DB, Glisson SN, Green D, Wixson RL: A prospective, randomized study of preoperative autologous donation for hip replacement surgery. *J Bone Joint Surg Am* 2002; 84-A:1299-304
38. Billote DB, Glisson SN, Green D, Wixson RL: Efficacy of preoperative autologous blood donation: analysis of blood loss and transfusion practice in total hip replacement. *J Clin Anesth* 2000; 12:537-542
39. Bjessmo S, Ivert T: Blood loss after coronary artery bypass surgery: relations to patient variables and antithrombotic treatment. *Scand Cardiovasc J* 2000; 34:438-445
40. Blais RE, Hadjipavlou AG, Shulman G: Efficacy of autotransfusion in spine surgery: comparison of autotransfusion alone and with hemodilution and apheresis. *Spine* 1996; 21:2795-2800
41. Blumberg N, Heal JM, Gettings KF, Phipps RP, Masel D, Refaai MA, Kirkley SA, Fialkow LB: An association between decreased cardiopulmonary complications (TRALI and TACO) and implementation of universal leukoreduction of blood transfusions. *Transfusion* 2010; 50:2738-2744
42. Bocchieri KA, Scheinerman SJ, Graver LM: Exchange transfusion before cardiopulmonary bypass in sickle cell disease. *Ann Thorac Surg* 2010; 90:323-324
43. Body SC, Birmingham J, Parks R, Ley C, Maddi R, Shernan SK, Siegel LC, Stover EP, D'Ambra MN, Levin J, Mangano DT, Spiess BD: Safety and efficacy of shed mediastinal blood transfusion after cardiac surgery: a multicenter observational study. Multicenter Study of Perioperative Ischemia Research Group. *J Cardiothorac Vasc Anesth* 1999; 13:410-416
44. Boezeman RP, Kelder JC, Waanders FG, de Vries JP: Continuous surveillance of lower limb perfusion during aortic surgery with near-infrared spectroscopy: a pilot study. *Vasc Endovascular Surg* 2011; 45:407-411
45. Bokesch PM, Szabo G, Wojdyga R, Grocott HP, Smith PK, Mazer CD, Veticaden S, Wheeler A, Levy JH: A phase 2 prospective, randomized, double-blind trial comparing the effects of tranexamic acid with ecallantide on blood loss from high-risk cardiac surgery with cardiopulmonary bypass (CONSERV-2 Trial). *J Thorac Cardiovasc Surg* 2012; 143:1022-1029

46. Bokesch PM, Szabo G, Wojdyga R, Grocott HP, Smith PK, Mazer CD, Veticaden S, Wheeler A, Levy JH: A phase 2 prospective, randomized, double-blind trial comparing the effects of tranexamic acid with ecallantide on blood loss from high-risk cardiac surgery with cardiopulmonary bypass (CONSERV-2 Trial). *J Thorac Cardiovasc Surg* 2012; 143:1022-1029
47. Boldt J, Weber A, Mailer K, Papsdorf M, Schuster P: Acute normovolaemic haemodilution vs controlled hypotension for reducing the use of allogeneic blood in patients undergoing radical prostatectomy. *Br J Anaesth* 1999; 82:170-174
48. Bottner F, Pavone V, Johnson T, Heitkemper S, Sculco TP: Blood management after bilateral total knee arthroplasty. *Clin Orthop* 2003; 410:254-261
49. Bouchard D, Marcheix B, Al-Shamary S, Vanden Eynden F, Demers P, Robitaille D, Pellerin M, Perrault LP, Carrier M: Preoperative autologous blood donation reduces the need for allogeneic blood products: a prospective randomized study. *Can J Surg* 2008; 51:422-427
50. Boyer-Neumann C, Dreyfus M, Wolf M, Veyradier A, Meyer D: Multi-therapeutic approach to manage delivery in an alloimmunized patient with type 3 von Willebrand disease. *J Thromb Haemost* 2003; 1:190-192
51. Boylan JF, Klinck JR, Sandler AN, Arellano R, Greig PD, Nierenberg H, Roger SL, Glynn MF: Tranexamic acid reduces blood loss, transfusion requirements, and coagulation factor use in primary orthotopic liver transplantation. *Anesthesiology* 1996; 85:1043-1048
52. Bracey AW, Radovancevic R, Riggs SA, Houston S, Cozart H, Vaughn WK, Radovancevic B, McAllister HA, Cooley DA: Lowering the hemoglobin threshold for transfusion in coronary artery bypass procedures: effect on patient outcome. *Transfusion* 1999; 39:1070-1077
53. Braga M, Gianotti L, Vignali A, Gentilini O, Servida P, Bordignon C, Di Carlo V: Evaluation of recombinant human erythropoietin to facilitate autologous blood donation before surgery in anaemic patients with cancer of the gastrointestinal tract. *Br J Surg* 1995; 82:1637-1640
54. Brener SJ, Bhatt DL, Moliterno DJ, Schneider JP, Ellis SG, Topol EJ: Revisiting optimal anticoagulation with unfractionated heparin during coronary stent implantation. *Am J Cardiol* 2003; 92:1468-1471
55. Bridgens JP, Evans CR, Dobson PM, Hamer AJ: Intraoperative red blood-cell salvage in revision hip surgery. A case-matched study. *J Bone Joint Surg Am* 2007; 89:270-275
56. Brown RS, Thwaites BK, Mongan PD: Tranexamic acid is effective in decreasing postoperative bleeding and transfusions in primary coronary artery bypass operations: a double-blind, randomized, placebo-controlled trial. *Anesth Analg* 1997; 85:963-970
57. Bruce D, Nokes TJ: Prothrombin complex concentrate (Beriplex P/N) in severe bleeding: experience in a large tertiary hospital. *Crit Care* 2008; 12:R105
58. Bui JD, Despotic GD, Trulock EP, Patterson GA, Goodnough LT: Fatal thrombosis after administration of activated prothrombin complex concentrates in a patient supported by extracorporeal membrane oxygenation who had received activated recombinant factor VII. *J Thorac Cardiovasc Surg* 2002; 124:852-854
59. Burman JF, Westlake AS, Davidson SJ, Rutherford LC, Rayner AS, Wright AM, Morgan CJ, Pepper JR: Study of five cell salvage machines in coronary artery surgery. *Transfus Med* 2002; 12:173-179

60. Bush RL, Pevec WC, Holcroft JW: A prospective, randomized trial limiting perioperative red blood cell transfusions in vascular patients. *Am J Surg* 1997; 174:143-148
61. Caglar GS, Tasci Y, Kayikcioglu F, Haberal A: Intravenous tranexamic acid use in myomectomy: a prospective randomized double-blind placebo controlled study. *Eur J Obstet Gynecol Reprod Biol* 2008; 137:227-231
62. Camarasa MA, Ollé G, Serra-Prat M, Martín A, Sánchez M, Ricós P, Pérez A, Opisso L: Efficacy of aminocaproic, tranexamic acids in the control of bleeding during total knee replacement: a randomized clinical trial. *Br J Anaesth* 2006; 96:576-582
63. Canver CC, Kroncke GM, Nichols RD, Murray EL, Mentzer RM Jr: The effects of intraoperative autologous whole blood sequestration on the need for transfusion of allogenic blood and blood products in coronary bypass operations. *J Cardiovasc Surg (Torino)* 1995; 36:423-438
64. Capraro L, Kuitunen A, Salmenpera M, Kekomaki R: On-site coagulation monitoring does not affect hemostatic outcome after cardiac surgery. *Acta Anaesthesiologica Scand* 2001; 45:200-206
65. Carson JL, Brooks MM, Abbott JD, Chaitman B, Kelsey SF, Triulzi DJ, Srinivas V, Menegus MA, Marroquin OC, Rao SV, Noveck H, Passano E, Hardison RM, Smitherman T, Vagaonescu T, Wimmer NJ, Williams DO: Liberal versus restrictive transfusion thresholds for patients with symptomatic coronary artery disease. *Am Heart J* 2013; 165:964-971
66. Carson JL, Terrin ML, Barton FB, Aaron R, Greenburg AG, Heck DA, Magaziner J, Merlino FE, Bunce G, McClelland B, Duff A, Noveck H: A pilot randomized trial comparing symptomatic vs. hemoglobin-level-driven red blood cell transfusions following hip fracture. *Transfusion* 1998; 38:522-529
67. Carson JL, Terrin ML, Noveck H, Sanders DW, Chaitman BR, Rhoads GG, Nemo G, Dragert K, Beaupre L, Hildebrand K, Macauley W, Lewis C, Cook DR, Dobbin G, Zakriya KJ, Apple FS, Horney RA, Magaziner J: Liberal or restrictive transfusion in high-risk patients after hip surgery. *N Engl J Med* 2011; 365:2453-2462
68. Casati A, Fanelli G, Pietropaoli P: Continuous monitoring of cerebral oxygen saturation in elderly patients undergoing general abdominal surgery minimizes brain exposure to potential hypoxia. *Anesth Analg* 2005; 101:740-747
69. Casati A, Fanelli G, Pietropaoli P: Monitoring cerebral oxygen saturation in elderly patients undergoing general abdominal surgery: a prospective cohort study. *Eur J Anaesthesiol* 2007; 24:59-65
70. Casati V, Bellotti F, Gerli C, Franco A, Oppizzi M, Cossolini M, Calori G, Benussi S, Alfieri O, Torri G: Tranexamic acid administration after cardiac surgery: A prospective, randomized, double-blind, placebo-controlled study. *Anesthesiology* 2001; 94:8-14
71. Casati V, D'Angelo A, Barbato L, Turolla D, Villa F, Grasso MA, Porta A, Guerra F: Perioperative management of four anaemic female Jehovah's Witnesses undergoing urgent complex cardiac surgery. *Br J Anaesth* 2007; 99:349-352
72. Casati V, Della Valle P, Benussi S, Franco A, Gerli C, Baili P, Alfieri O, D'Angelo A: Effects of tranexamic acid on postoperative bleeding and related hematochemical variables in coronary surgery: comparison between on-pump and off-pump techniques. *J Thorac Cardiovasc Surg* 2004; 128:83-91

73. Casati V, Gerli C, Franco A, Torri G, D'Angelo A, Benussi S, Alfieri O: Tranexamic acid in off-pump coronary surgery: A preliminary, randomized, double-blind, placebo-controlled study. *Ann Thorac Surg* 2001; 72:470-475
74. Casati V, Sandrelli L, Speziali G, Calori G, Grasso MA, Spagnolo S: Hemostatic effects of tranexamic acid in elective thoracic aortic surgery: a prospective randomized, double-blinded, placebo-controlled study. *J Thorac Cardiovasc Surg* 2002; 123:1084-1091
75. Casati V, Speziali G, D'Alessandro C, Cianchi C, Antonietta Grasso M, Spagnolo S, Sandrelli L: Intraoperative low-volume acute normovolemic hemodilution in adult open-heart surgery. *Anesthesiology* 2002; 97:367-373
76. Cash KL, Brown T, Sausais L, Uehlinger J, Reed LJ: Severe delayed hemolytic transfusion reaction secondary to anti-At(a). *Transfusion* 1999; 39:834-837
77. Cata JP, Klein EA, Hoeltge GA, Dalton JE, Mascha E, O'Hara J, Russell A, Kurz A, Ben-Elihayhu S, Sessler DI: Blood storage duration and biochemical recurrence of cancer after radical prostatectomy. *Mayo Clin Proc* 2011; 86:120-127
78. Chan ACW, Blumgart LH, Wuest DL, Melendez JA, Fong Y: Use of preoperative autologous blood donation in liver resections for colorectal metastases. *Am J Surg* 1998; 175:461-465
79. Chanda A, Smith DR, Nanda A: Autotransfusion by cell saver technique in surgery of lumbar and thoracic spinal fusion with instrumentation. *J Neurosurg* 2002; 96:298-303
80. Chapman WC, Clavien PA, Fung J, Khanna A, Bonham A: Effective control of hepatic bleeding with a novel collagen-based composite combined with autologous plasma: results of a randomized controlled trial. *Arch Surg* 2000; 135:1200-1204
81. Chelemer SB, Prato BS, Cox PM Jr, O'Connor GT, Morton JR: Association of bacterial infection and red blood cell transfusion after coronary artery bypass surgery. *Ann Thorac Surg* 2002; 73:138-142
82. Chen G, Zhang FJ, Gong M, Yan M: Effect of perioperative autologous versus allogeneic blood transfusion on the immune system in gastric cancer patients. *J Zhejiang Univ Sci B* 2007; 8:560-565
83. Chen J, Singhapricha T, Memarzadeh M, Ziman A, Yuan S, Hu KQ, Steadman RH, Busuttill RW, Xia VW: Storage age of transfused red blood cells during liver transplantation and its intraoperative and postoperative effects. *World J Surg* 2012; 36:2436-2442
84. Cheng HW, Chang HH, Chen YJ, Chang WK, Chan KH, Chen PT: Clinical value of application of cerebral oximetry in total replacement of the aortic arch and concomitant vessels. *Acta Anaesthesiol Taiwan* 2008; 46:178-183
85. Chikada M, Furuse A, Kotsuka Y, Yagyu K: Open-heart surgery in Jehovah's Witness patients. *Cardiovasc Surg* 1996; 4:311-314
86. Choi WS, Irwin MG, Samman N: The effect of tranexamic acid on blood loss during orthognathic surgery: a randomized controlled trial. *J Oral Maxillofac Surg* 2009; 67:125-133
87. Chowdhury M, Shore-Lesserson L, Mais AM, Leyvi G: Thromboelastography with platelet mapping predicts postoperative chest tube drainage in patients undergoing coronary artery bypass grafting. *J Cardiothorac Vasc Anesth* 2014; 28:217-223
88. Christodoulakis M, Tsiftsis DD: Preoperative epoetin alfa in colorectal surgery: a randomized, controlled study. *Ann Surg Oncol* 2005; 12:718-725

89. Chung YT, Wu YC, Chen YH: Postoperative pulmonary edema, transfusion-related?--a case report. *Acta Anaesthesiol Sin* 2003; 41:43-46
90. Cina CS, Bruin G: Acute normovolemic hemodilution (ANH) in surgery of the thoraco-abdominal aorta. A cohort study to evaluate coagulation parameters and blood products utilization. *J Cardiovasc Surg* 1999; 40:37-43
91. Cina CS, It SC, Clase CM, Bruin G: A cohort study of coagulation parameters and the use of blood products in surgery of the thoracic and thoracoabdominal aorta. *J Vasc Surg* 2001; 33:462-468
92. Claeys MA, Vermeersch N, Haentjens P: Reduction of blood loss with tranexamic acid in primary total hip replacement surgery. *Acta Chir Belg* 2007; 107:397-401
93. Clagett GP, Valentine RJ, Jackson MR, Mathison C, Kakish HB, Bengtson TD: A randomized trial of intraoperative autotransfusion during aortic surgery. *J Vasc Surg* 1999; 29:22-30
94. Clagett GP, Valentine RJ, Myers SI, Chervu A, Heller J: Does desmopressin improve hemostasis and reduce blood loss from aortic surgery? A randomized, double-blind study. *J Vasc Surg* 1995; 22:223-229
95. Clavé A, Fazilleau F, Dumser D, Lacroix J: Efficacy of tranexamic acid on blood loss after primary cementless total hip replacement with rivaroxaban thromboprophylaxis: A case-control study in 70 patients. *Orthop Traumatol Surg Res* 2012; 98:484-490
96. Clugston PA, Fitzpatrick DG, Kester DA, Folet B, Germann E: Autologous blood use in reduction mammoplasty: is it justified? *Plast Reconstr Surg* 1995; 95:824-828
97. Coakley M, Reddy K, Mackie I, Mallett S: Transfusion triggers in orthotopic liver transplantation: a comparison of the thromboelastometry analyzer, the thromboelastogram, and conventional coagulation tests. *J Cardiothorac Vasc Anesth* 2006; 20:548-553
98. Codispoti M, Mankad PS: Significant merits of a fibrin sealant in the presence of coagulopathy following paediatric cardiac surgery: randomised controlled trial. *Eur J Cardiothorac Surg* 2002; 22:200-205
99. Coffey A, Pittmam J, Halbrook H, Fehrenbacher J, Beckman D, Hormuth D: The use of tranexamic acid to reduce postoperative bleeding following cardiac surgery: a double-blind randomized trial. *Am Surg* 1995; 61:566-568
100. Cohn SM, Pearl RG, Acosta SM, Nowlin MU, Hernandez A, Guta C, Michalek JE: A prospective randomized pilot study of near-infrared spectroscopy-directed restricted fluid therapy versus standard fluid therapy in patients undergoing elective colorectal surgery. *Am Surg* 2010; 76:1384-1392
101. Connelly NR, Kiessling BM, Brull SJ: Does tranexamic acid decrease bleeding in patients undergoing cardiopulmonary bypass? *Yale J Biol Med* 1994; 67:265-268
102. Connor JP, Morris PC, Alagoz T, Anderson B, Bottles K, Buller RE: Intraoperative autologous blood collection and autotransfusion in the surgical management of early cancers of the uterine cervix. *Obstet Gynecol* 1995; 86:373-378
103. Consten ECJ, Henny CP, Eijnsman L, Dongelmans DA, van Oers MHJ: The routine use of fresh frozen plasma in operations with cardiopulmonary bypass is not justified. *J Thorac Cardiovasc Surg* 1996; 112:162-167
104. Copley LA, Richards BS, Safavi FZ, Newton PO: Hemodilution as a method to reduce transfusion requirements in adolescent spine fusion surgery. *Spine* 1999; 24:219-222

105. Crescenti A, Borghi G, Bignami E, Bertarelli G, Landoni G, Casiraghi GM, Briganti A, Montorsi F, Rigatti P, Zangrillo A: Intraoperative use of tranexamic acid to reduce transfusion rate in patients undergoing radical retropubic prostatectomy: double blind, randomised, placebo controlled trial. *BMJ* 2011; 343:d5701. doi
106. Cuenca J, Garcia-Erce JA, Martinez F, Cardona R, Perez-Serrano L, Munoz M: Preoperative haematinics and transfusion protocol reduce the need for transfusion after total knee replacement. *Int J Surg* 2007; 5:89-94
107. Cuenca J, Garcia-Erce JA, Martinez F, Perez-Serrano L, Herrera A, Munoz M: Perioperative intravenous iron, with or without erythropoietin, plus restrictive transfusion protocol reduce the need for allogeneic blood after knee replacement surgery. *Transfusion* 2006; 46:1112-1119
108. Daane CR, Golab HD, Meeder JH, Wijers MJ, Bogers AJ: Processing and transfusion of residual cardiopulmonary bypass volume: effects on haemostasis, complement activation, postoperative blood loss and transfusion volume. *Perfusion* 2003; 18:115-121
109. Daily PO, Lamphere JA, Dembitsky WP, Adamson RM, Dans NF: Effect of prophylactic epsilon-aminocaproic acid on blood loss and transfusion requirements in patients undergoing first-time coronary artery bypass grafting. A randomized, prospective, double-blind study. *J Thorac Cardiovasc Surg* 1994; 108:99-106
110. Dalmau A, Sabate A, Acosta F, Garcia-Huete L, Koo M, Sansano T, Rafecas A, Figueras J, Jaurrieta E, Parrilla P: Tranexamic acid reduces red cell transfusion better than epsilon-aminocaproic acid or placebo in liver transplantation. *Anesth Analg* 2000; 91:29-34
111. Dalrymple-Hay MJ, Dawkins S, Pack L, Deakin CD, Sheppard S, Ohri SK, Haw MP, Livesey SA, Monro JL: Autotransfusion decreases blood usage following cardiac surgery - a prospective randomized trial. *Cardiovasc Surg* 2002; 9:184-187
112. Damgaard S, Steinbruchel DA: Autotransfusion with cell saver for off-pump coronary artery bypass surgery: a randomized trial. *Scand Cardiovasc J* 2006; 40:194-198
113. Dave SP, Greenstein AJ, Sachar DB, Harpaz N, Aledort L: Bleeding diathesis in amyloidosis with renal insufficiency associated with Crohn's disease: response to desmopressin. *Am J Gastroenterol* 2002; 97:187-189
114. Davidson SJ, McGrowder D, Roughton M, Kelleher AA: Can ROTEM thromboelastometry predict postoperative bleeding after cardiac surgery? *J Cardiothorac Vasc Anesth* 2008; 22:655-661
115. de Andrade JR, Jove M, Landon G, Frei D, Guilfoyle M, Young DC: Baseline hemoglobin as a predictor of risk of transfusion and response to Epoetin alfa in orthopedic surgery patients. *Am J Orthop* 1996; 25:533-542
116. de Pree C, Mermillod B, Hoffmeyer P, Beris P: Recombinant human erythropoietin as adjuvant treatment for autologous blood donation in elective surgery with large blood needs ( $\geq 5$  units): a randomized study. *Transfusion* 1997; 37:708-714
117. De Santo LS, Amarelli C, Della Corte A, Scardone M, Bancone C, Carozza A, Grassia MG, Romano G: Blood transfusion after on-pump coronary artery bypass grafting: focus on modifiable risk factors. *Eur J Cardiothorac Surg* 2013; 43:359-366
118. del Pozo Pozo AI, Jimenez-Yuste V, Villar A, Quintana M, Hernandez-Navarro F: Successful thyroidectomy in a patient with Hermansky-Pudlak syndrome treated with recombinant activated factor VII and platelet concentrates. *Blood Coagul Fibrinolysis* 2002; 13:551-553



119. del Trujillo MM, Carrero A, Muñoz M: The utility of the perioperative autologous transfusion system OrthoPAT in total hip replacement surgery: a prospective study. *Arch Orthop Trauma Surg* 2008; 128:1031-1038
120. Despotis GJ, Levine V, Filos KS, Santoro SA, Joist JH, Spitznagel E, Goodnough LT: Evaluation of a new point-of-care test that measures PAF-mediated acceleration of coagulation in cardiac surgical patients. *Anesthesiology* 1996; 85:1311-1323
121. Despotis GJ, Levine V, Saleem R, Spitznagel E, Joist JH: Use of point-of-care test in identification of patients who can benefit from desmopressin during cardiac surgery: a randomised controlled trial. *Lancet* 1999; 354:106-110
122. Devereaux PJ, Mrkobrada M, Sessler DI, Leslie K, Alonso-Coello P, Kurz A, Villar JC, Sigamani A, Biccari BM, Meyhoff CS, Parlow JL, Guyatt G, Robinson A, Garg AX, Rodseth RN, Botto F, Lurati Buse G, Xavier D, Chan MT, Tiboni M, Cook D, Kumar PA, Forget P, Malaga G, Fleischmann E, Amir M, Eikelboom J, Mizera R, Torres D, Wang CY, VanHelder T, Paniagua P, Berwanger O, Srinathan S, Graham M, Pasin L, Le Manach Y, Gao P, Pogue J, Whitlock R, Lamy A, Kearon C, Baigent C, Chow C, Pettit S, Chrolavicius S, Yusuf S: Aspirin in patients undergoing noncardiac surgery. *N Engl J Med* 2014; 370:1494-1503
123. Dexter F, Ledolter J, Davis E, Witkowski TA, Herman JH, Epstein RH: Systematic criteria for type and screen based on procedure's probability of erythrocyte transfusion. *Anesthesiology* 2012 116:768-778
124. Dich-Nielsen JO, Rajan RM, Jensen JJ: An anaphylactoid-like reaction following infusion of salvaged unwashed drain blood. *Can J Anaesth* 1998; 45:189
125. Diprose P, Herbertson MJ, O'Shaughnessy D, Gill RS: Activated recombinant factor VII after cardiopulmonary bypass reduces allogeneic transfusion in complex non-coronary cardiac surgery: Randomized double-blind placebo-controlled pilot study. *Br J Anaesth* 2005; 95:596-602
126. Djaiani G, Fedorko L, Borger MA, Green R, Carroll J, Marcon M, Karski J: Continuous-flow cell saver reduces cognitive decline in elderly patients after coronary bypass surgery. *Circulation* 2007; 116:1888-1895
127. Dotan ZA, Mor Y, Leibovitch I, Varon D, Golomb J, Duvdevani M, Ramon J: The efficacy and safety of perioperative low molecular weight heparin substitution in patients on chronic oral anticoagulant therapy undergoing transurethral prostatectomy for bladder outlet obstruction. *J Urol* 2002; 168:610-613
128. Dryden PJ, O'Connor JP, Jamieson WR, Reid I, Ansley D, Sadeghi H, Burr LH, Munro AI, Merrick PM: Tranexamic acid reduces blood loss and transfusion in reoperative cardiac surgery. *Can J Anaesth* 1997; 44:934-941
129. Dzik WH, Andersen JK, O'Neill EM, Assmann SF, Kalish LA, Stowell CP: A prospective, randomized clinical trial of universal WBC reduction. *Transfusion* 2002; 42:1114-1122
130. Edgren G, Kamper-Jørgensen M, Eloranta S, Rostgaard K, Custer B, Ullum H, Murphy EL, Busch MP, Reilly M, Melbye M, Hjalgrim H, Nyrén O: Duration of red blood cell storage and survival of transfused patients (CME). *Transfusion* 2010; 50:1185-1195
131. Ekback G, Axelsson K, Rytberg L, Edlund B, Kjellberg J, Weckstrom J, Carlsson O, Schott U: Tranexamic acid reduces blood loss in total hip replacement surgery. *Anesth Analg* 2000; 91:1124-1130
132. Ekback G, Schott U, Axelsson K, Carlberg M: Perioperative autotransfusion and functional coagulation analysis in total hip replacement. *Acta Anaesthes Scand* 1995; 39:390-395

133. Engel JM, Hohaus T, Ruwoldt R, Menges T, Jurgensen I, Hempelmann G: Regional hemostatic status and blood requirements after total knee arthroplasty with and without tranexamic acid or aprotinin. *Anesth Analg* 2001; 92:775-780
134. Engstrom KG, Appelblad M, Brorsson, B: Mechanisms behind operating room blood transfusions in coronary artery bypass graft surgery patients with insignificant bleeding. *J Cardiothorac Vasc Anes* 2002; 16:539-544
135. Ereth MH, Nuttall GA, Klindworth JT, MacVeigh I, Santrach PJ, Orszulak TA, Harmsen WS, Oliver W Jr: Does the platelet-activated clotting test (Hemo-STATUS) predict blood loss and platelet dysfunction associated with cardiopulmonary bypass? *Anesth Analg* 1997; 85:257-264
136. Ereth MH, Nuttall GA, Santrach PJ, Klindworth JT, Oliver WC Jr, Schaff HV: The relation between the platelet-activated clotting test (HemoSTATUS) and blood loss after cardiopulmonary bypass. *Anesthesiology* 1998; 88:962-969
137. Estella NM, Berry DL, Baker BW, Wali AT, Belfort MA: Normovolemic hemodilution before cesarean hysterectomy for placenta percreta. *Obstet Gynecol* 1997; 90:669-670
138. Etchason J, Petz L, Keeler E, Calhoun L, Kleinman S, Snider C, Fink A, Brook R: The cost effectiveness of preoperative autologous blood donations. *N Engl J Med* 1995; 332:719-724
139. Faber P, Reid C, El-Shafei H, Falase B, DeAnde A Jr, Mazer CD: Recombinant factor VIIa in the management of postoperative bleeding after repair for inadvertently thrombolysed acute type A aortic Dissection. *J Cardiovasc Vasc Anesth* 2006; 20:736-741
140. Faris PM, Ritter MA, Abels RI: The effects of recombinant human erythropoietin on perioperative transfusion requirements in patients having a major orthopaedic operation. The American Erythropoietin Study Group. *J Bone Joint Surg Am* 1996; 78:62-72
141. Feagan BG, Wong CJ, Kirkley A, Johnston DW, Smith FC, Whitsitt P, Wheeler SL, Lau CY: Erythropoietin with iron supplementation to prevent allogeneic blood transfusion in total hip joint arthroplasty. A randomized, controlled trial. *Ann Intern Med* 2000; 133:845-854
142. Fenger-Eriksen C, Jensen TM, Kristensen BS, Jensen KM, Tønnesen E, Ingerslev J, Sørensen B: Fibrinogen substitution improves whole blood clot firmness after dilution with hydroxyethyl starch in bleeding patients undergoing radical cystectomy: a randomized, placebo-controlled clinical trial. *J Thromb Haemost* 2009; 7:795-802
143. Ferraz-Neto BH, Silva ED, Afonso RC, Gregory FH, Meira-Filho SP, Macedo CP, Parra OM, Leitao RM, Saad WA: Acute normovolemic intraoperative hemodilution in liver transplantation. *Transplant Proc* 1999; 31:3069-3070
144. Figueras J, Llado L, Miro M, Ramos E, Torras J, Fabregat J, Serrano T: Application of fibrin glue sealant after hepatectomy does not seem justified: results of a randomized study in 300 patients. *Ann Surg* 2007; 245:536-542
145. Firanescu CE, Martens EJ, Schönberger JP, Soliman Hamad MA, van Straten AH: Postoperative blood loss in patients undergoing coronary artery bypass surgery after preoperative treatment with clopidogrel. A prospective randomized controlled study. *Eur J Cardiothorac Surg* 2009; 36:856-862
146. Fischer M, Matsuo K, Gonen M, Grant F, Dematteo RP, D'Angelica MI, Mascarenhas J, Brennan MF, Allen PJ, Blumgart LH, Jarnagin WR: Relationship between intraoperative fluid administration and perioperative outcome after pancreaticoduodenectomy: results of

- a prospective randomized trial of acute normovolemic hemodilution compared with standard intraoperative management. *Ann Surg* 2010; 252:952-958
147. Florentino-Pineda I, Blakemore LC, Thompson GH, Poe-Kochert C, Adler P, Tripi P: The Effect of  $\epsilon$ -aminocaproic acid on perioperative blood loss in patients with idiopathic scoliosis undergoing posterior spinal fusion: a preliminary prospective study. *Spine* 2001; 26:1147-1151
  148. Florentino-Pineda I, Thompson GH, Poe-Kochert C, Huang RP, Haber LL, Blakemore LC: The effect of Amicar on perioperative blood loss in idiopathic scoliosis: The results of a prospective, randomized double-blind study. *Spine* 2004; 29:233-238
  149. Ford SM, Unsworth-White MJ, Aziz T, Tooze JA, van Besouw JP, Bevan DH, Treasure T: Platelet pheresis is not a useful adjunct to blood-sparing strategies in cardiac surgery. *J Cardiothorac Vasc Anesth* 2002; 16:321-329
  150. Fox JS, Amaranath L, Hoeltge GA, Andrish JT: Autologous blood transfusion and intraoperative cell salvage in a patient with homozygous sickle cell disease. *Cleve Clin J Med* 1994; 61:137-140
  151. Franchini M, de Gironcoli M, Lippi G, Manzato F, Aprili G, Gandini G: Prophylactic use of desmopressin in surgery of six patients with symptomatic heterozygous factor XI deficiency. *Haematologica* 2000; 85:106-107
  152. Frank SM, Rothschild JA, Masear CG, Rivers RJ, Merritt WT, Savage WJ, Ness PM: Optimizing preoperative blood ordering with data acquired from an anesthesia information management system. *Anesthesiology* 2013; 118:1286-1297
  153. Friederich PW, Henny CP, Messelink EJ, Geerdink MG, Keller T, Kurth KH, Buller HR, Levi M: Effect of recombinant activated factor VII on perioperative blood loss in patients undergoing retropubic prostatectomy: a double-blind placebo-controlled randomised trial. *Lancet* 2003; 361:201-205
  154. Garcia-Huete L, Domenech P, Sabate A, Martinez-Brotons F, Jaurrieta E, Figueras J: The prophylactic effect of aprotinin on intraoperative bleeding in liver transplantation: a randomized clinical study. *Hepatology* 1997; 26:1143-1148
  155. Gardner A, Gibbs N, Evans C, Bell R: Relative cost of autologous red cell salvage versus allogeneic red cell transfusion during abdominal aortic aneurysm repair. *Anaesth Intensive Care* 2000; 28:646-649
  156. Garrido-Martín P, Nassar-Mansur MI, de la Llana-Ducrós R, Virgos-Aller TM, Rodríguez Fortunez PM, Ávalos-Pinto R, Jimenez-Sosa A, Martínez-Sanz R: The effect of intravenous and oral iron administration on perioperative anaemia and transfusion requirements in patients undergoing elective cardiac surgery: a randomized clinical trial. *Interact Cardiovasc Thorac Surg* 2012; 15:1013-1018
  157. Gaudino M, Luciani N, Piancone FL, Bruno P, Rossi M, Schiavello R, Possati G: Perioperative management of a patient with Werlhof disease undergoing myocardial revascularization. *J Cardiovasc Surg (Torino)* 1999; 40:227-228
  158. Gelb AB, Roth RI, Levin J, London MJ, Noall RA, Hauck WW, Cloutier M, Verrier E, Mangano DT: Changes in blood coagulation during and following cardiopulmonary bypass. *Am J Clin Pathol* 1996; 106:87-99
  159. Gelsomino S, Lorusso R, Romagnoli S, Bevilacqua S, Dr Cicco G, Bille G, Stefano P, Gensini GF: Treatment of refractory bleeding after cardiac operations with low-dose recombinant activated factor VII (NovoSeven): A propensity score analysis. *Eur J Cardiothorac Surg* 2008; 33:64-71

160. Gerrah R, Shargal Y, Elami A: Impaired oxygenation and increased hemolysis after cardiopulmonary bypass in patients with glucose-6-phosphate dehydrogenase deficiency. *Ann Thorac Surg* 2003; 76:523-527
161. Gilbert JB, Malkowicz SB, Wein AJ: Cell saver and radical retropubic prostatectomy: analysis of cost-effectiveness. *Urology* 1995; 46:542-544
162. Gill R, Herbertson M, Vuysteke A, Skov P, von Heymann C, Mythen M, Sellke F, Booth R, Schmidt TA: Safety and efficacy of recombinant activated factor VII. A randomized placebo-controlled trial in the setting of bleeding after cardiac surgery. *Circulation* 2009; 320:21-27
163. Giurati G, Zanetta G, Caspani L, Landon F, Pellegrino A, Baglioni S, Mangioni C: Autologous blood transfusion in patients undergoing radical hysterectomy. *Br J Obstet Gynaecol* 1995; 102:64-65
164. Glazier DB, Ciocca RG, Gosin JS, Murphy DP, Graham AM: Elective aortic surgery with minimal banked blood. *Am Surg* 1998; 64:171-174
165. Goel P, Pannu H, Mohan D, Arora R: Efficacy of cell saver in reducing homologous blood transfusions during OPCAB surgery: a prospective randomized trial. *Transfus Med* 2007; 17:285-289
166. Goh M, Kleer CG, Kielczewski P, Wojno KJ, Kim K, Oesterling JE: Autologous blood donation prior to anatomical radical retropubic prostatectomy: is it necessary? *Urology* 1997; 49:569-573
167. Gombotz H, Gries M, Sipurzynski S, Fruhwald S, Rehak P: Preoperative treatment with recombinant human erythropoietin or predeposit of autologous blood in women undergoing primary hip replacement. *Acta Anaesthesiol Scand* 2000; 44:737-742
168. Goobie SM, Meier PM, Pereira LM, McGowan FX, Prescilla RP, Scharp LA, Rogers GF, Proctor MR, Meara JG, Soriano SG, Zurakowski D, Sethna NF: Efficacy of tranexamic acid in pediatric craniostomy surgery: a double-blind, placebo-controlled trial. *Anesthesiology* 2011; 114:862-871
169. Good L, Peterson E, Lisander B: Tranexamic acid decreases external blood loss but not hidden blood loss in total knee replacement. *Br J Anaesth* 2003; 90:596-599
170. Goodnough LT, Monk TG, Sicard G, Satterfield SA, Allen B, Anderson CB, Thompson RW, Flye W, Martin K: Intraoperative salvage in patients undergoing elective abdominal aortic aneurysm repair: an analysis of cost and benefit. *J Vasc Surg* 1996; 24:213-218
171. Goodnough LT, Price TH, Friedman KD, Johnston M, Ciavarella D, Khan N, Sacher R, Vogler WR, Wissel M, Abels RI: A phase III trial of recombinant human erythropoietin therapy in non-anemic orthopedic patients subjected to aggressive removal of blood for autologous use: Dose, response, toxicity, and efficacy. *Transfusion* 1994; 34:66-71
172. Gorlinger K, Dirkmann D, Hanke AA, Kamler M, Kottenberg E, Thielmann M, Jakob H, Peters J: First-line therapy with coagulation factor concentrates combined with point-of-care coagulation testing is associated with decreased allogeneic blood transfusion in cardiovascular surgery: A retrospective, single-center cohort study. *Anesthesiology* 2011; 115:1179-1191
173. Gouvêa G, Diaz R, Auler L, Toledo R, Soluri A, Haack AG, de Oliveira MM, Vilanova B, Machado VR, Martinho JM: Perioperative coagulation profile in living liver donors as assessed by rotational thromboelastometry. *Liver Transpl* 2010; 16:387-392

174. Govinda R, Kasuya Y, Bala E, Mahboobi R, Devarajan J, Sessler DI, Akça O: Early postoperative subcutaneous tissue oxygen predicts surgical site infection. *Anesth Analg* 2010; 111:946-952
175. Gravlee GP, Arora S, Lavendar SW, Mills SA, Hudspeth AS, Cordell AR, James RL, Brockschmidt JK, Stuart JJ: Predictive value of blood clotting tests in cardiac surgical patients. *Ann Thorac Surg* 1994; 58:216-221
176. Gray CL, Amling CL, Polston GR, Powell CR, Kane CJ: Intraoperative cell salvage in radical retropubic prostatectomy. *Urology* 2001; 58:740-745
177. Grosvenor D, Goyal V, Goodman S: Efficacy of postoperative blood salvage following total hip arthroplasty in patients with and without deposited autologous units. *J Bone Joint Surg Am* 2000; 82-A:951-954
178. Grover M, Talwalkar S, Casbard A, Boralessa H, Contreras M, Boralessa H, Brett S, Goldhill DR, Soni N: Silent myocardial ischaemia and haemoglobin concentration: a randomized controlled trial of transfusion strategy in lower limb arthroplasty. *Vox Sang* 2006; 90:105-112
179. Guo JR, Yu J, Jin XJ, Du JM, Guo W, Yuan XH: Effects of acute normovolemic hemodilution on perioperative coagulation and fibrinolysis in elderly patients undergoing hepatic carcinectomy. *Chin Med Sci J* 2010; 25:146-150
180. Gutierrez MC, Goodnough LT, Druzin M, Butwick AJ: Postpartum hemorrhage treated with a massive transfusion protocol at a tertiary obstetric center: a retrospective study. *Int J Obstet Anesth* 2012; 21:230-235
181. Haas T, Spielmann N, Mauch J, Madjdpour C, Speer O, Schmutz M, Weiss M: Comparison of thromboelastometry (ROTEM®) with standard plasmatic coagulation testing in paediatric surgery. *Br J Anaesth* 2012; 108:36-41
182. Hajjar LA, Vincent JL, Galas FR, Nakamura RE, Silva CM, Santos MH, Fukushima J, Kalil Filho R, Sierra DB, Lopes NH, Mauad T, Roquim AC, Sundin MR, Leão WC, Almeida JP, Pomerantzeff PM, Dallan LO, Jatene FB, Stolf NA, Auler JO Jr: Transfusion requirements after cardiac surgery: the TRACS randomized controlled trial. *JAMA* 2010; 304:1559-1567
183. Hallak M, Sharon AS, Diukman R, Auslender R, Abramovici H: Supplementing iron intravenously in pregnancy. A way to avoid blood transfusions. *J Reprod Med* 1997; 42:99-103
184. Hanke AA, Joch C, Görlinger K: Long-term safety and efficacy of a pasteurized nanofiltrated prothrombin complex concentrate (Beriplex P/N): a pharmacovigilance study. *Br J Anaesth* 2013; 110:764-772
185. Hardy JF, Belisle S, Dupont C, Harel F, Robitaille D, Roy M, Gagnon L: Prophylactic tranexamic acid and epsilon-aminocaproic acid for primary myocardial revascularization. *Ann Thorac Surg* 1998; 65:371-376
186. Harley BJ, Beaupre LA, Jones CA, Cinats JG, Guenther CR: The effect of epsilon aminocaproic acid on blood loss in patients who undergo primary total hip replacement: a pilot study. *Can J Surg* 2002; 45:185-190
187. Harwin SF, Issa K, Naziri Q, Johnson AJ, Mont MA: Results of primary total knee arthroplasty in Jehovah's Witness patients. *J Arthroplasty* 2013; 28:49-55
188. Harwin SF, Pivec R, Johnson AJ, Naziri Q, Mont MA: Revision total hip arthroplasty in Jehovah's Witnesses. *Orthopedics* 2012; 35:e1145-e1151

189. Hayashi J, Kumon K, Takanashi S, Kawashima Y, Eguchi S, Takaku F, Yamamura H: Subcutaneous administration of recombinant human erythropoietin before cardiac surgery: a double-blind, multicenter trial in Japan. *Transfusion* 1994; 34:142-146
190. Haynes SL, Torella F, Wong JC, Dalrymple K, James M, McCollum CN: Economic evaluation of a randomized clinical trial of haemodilution with cell salvage in aortic surgery. *Br J Surg* 2002; 89:731-736
191. Healy WJ, Pfeifer BA, Kurtz SR, Johnson C, Johnson W, Johnston R, Sanders D, Karpman R, Hallack GN, Valeri CR: Evaluation of autologous shed blood for autotransfusion after orthopaedic surgery. *Clin Orthop* 1994; 299:53-59
192. Hebert PC, Chin-Yee I, Fergusson D, Blajchman M, Martineau R, Clinch J, Olberg B: A pilot trial evaluating the clinical effects of prolonged storage of red cells. *Anesth Analg* 2005; 100:1433-1438
193. Hedstrom M, Flordal PA, Ahl T, Svensson J, Dalen N: Autologous blood in hip replacement. No effect on blood loss but less increase of plasminogen activator inhibitor in a randomized series of 80 patients. *Acta Orthop Scand* 1996; 67:317-320
194. Heiss MM, Mempel W, Delanoff C, Jauch KW, Gabka C, Mempel M, Dieterich HJ, Eissner HJ, Schildberg FW: Blood transfusion-modulated tumor recurrence: first results of a randomized study of autologous versus allogeneic blood transfusion in colorectal cancer surgery. *J Clin Oncol* 1994; 12:1859-1867
195. Heiss MM, Tarabichi A, Delanoff C, Allgayer H, Jauch KW, Hernandez-Richter T, Mempel W, Beck KG, Schildberg FW, Messmer K: Perisurgical erythropoietin application in anemic patients with colorectal cancer: A double-blind randomized study. *Surgery* 1996; 119:523-527
196. Helm RE, Klemperer JD, Rosengart TK, Gold JP, Peterson P, DeBois W, Altorki NK, Lang S, Thomas S, Isom OW, Krieger KH: Intraoperative autologous blood donation preserves red cell mass but does not decrease postoperative bleeding. *Ann Thorac Surg* 1996; 62:1431-1441
197. Hendriks HG, Meijer K, de Wolf JT, Klompemaker IJ, Porte RJ, de Kam PJ, Hagens AJ, Melsen T, Slooff MJ, van der Meer J: Reduced transfusion requirements by recombinant factor VIIa in orthotopic liver transplantation: a pilot study. *Transplantation* 2001;71:402-405
198. Hendriks HG, van der Meer J, Klompemaker IJ, Choudhury N, Hagens JA, Porte RJ, de Kam PJ, Slooff MJ, de Wolf JT: Blood loss in orthotopic liver transplantation: a retrospective analysis of transfusion requirements and the effects of autotransfusion of cell saver blood in 164 consecutive patients. *Blood Coagul Fibrinolysis* 2000; 11:S87-S93
199. Herbstreit F, Winter EM, Peters J, Hartmann M: Monitoring of haemostasis in liver transplantation: comparison of laboratory based and point of care tests. *Anaesthesia* 2010; 65:44-49
200. Herregods L, Moerman A, Foubert L, Den Blauwen N, Mortier E, Poelaert J, Struys M: Limited intentional normovolemic hemodilution: ST-segment changes and use of homologous blood products in patients with left main coronary artery stenosis. *J Cardiothorac Vasc Anesth* 1997; 11:18-23
201. Hiippala S, Strid L, Wennerstrand M, Arvela V, Mantyla S, Ylinen J, Niemela H: Tranexamic acid (Cyklokapron) reduces perioperative blood loss associated with total knee arthroplasty. *Br J Anaesth* 1995; 74:534-537

202. Hiippala ST, Strid LJ, Wennerstrand MI, Arvela JV, Niemela HM, Mantyla SK, Kuisma RP, Ylinen JE: Tranexamic acid radically decreases blood loss and transfusions associated with total knee arthroplasty. *Anesth Analg* 1997; 84:839-844
203. Hobisch-Hagen P, Schobersberger W, Falkensammer J, Luz G, Innerhofer P, Frischhut B, Puschendorf B, Mair J: No release of cardiac troponin I during major orthopedic surgery after acute normovolemic hemodilution. *Acta Anaesthesiol Scand* 1998; 42:799-804
204. Hohn L, Schweizer A, Licker M, Morel DR: Absence of beneficial effect of acute normovolemic hemodilution combined with aprotinin on allogeneic blood transfusion requirements in cardiac surgery. *Anesthesiology* 2002; 96:276-282
205. Horlocker TT, Nuttall GA, Dekutoski MB, Bryant SC: The accuracy of coagulation tests during spinal fusion and instrumentation. *Anesth Analg* 2001; 93:33-38
206. Houbiers JG, Brand A, van de Watering LM, Hermans J, Verwey PJ, Bijnen AB, Pahlplatz P, Eeftinck Schattenkerk M, Wobbes T, de Vries JE: Randomised controlled trial comparing transfusion of leukocyte-depleted or buffy-coat-depleted blood in colorectal cancer. *Lancet* 1994; 344:573-578
207. Huber TS, Carlton LC, Irwin PB, Flug RR, Harward TR, Flynn TC, Seeger JM: Intraoperative autologous transfusion during elective infrarenal aortic reconstruction. *J Surg Res* 1997; 67:14-20
208. Husted H, Blond L, Sonne-Holm S, Holm G, Jacobsen TW, Gebuhr P: Tranexamic acid reduces blood loss and blood transfusions in primary total hip arthroplasty: a prospective randomized double-blind study in 40 patients. *Acta Orthop Scand* 2003; 74:665-669
209. Hyllner M, Avall A, Swolin B, Bengtson JP, Bengtsson A: Autologous blood transfusion in radical hysterectomy with and without erythropoietin therapy. *Obstet Gynecol* 2002; 99:757-762
210. Hynes M, Calder P, Scott G: The use of tranexamic acid to reduce blood loss during total knee arthroplasty. *Knee* 2003; 10:375-377
211. Ives CL, Harrison DK, Stansby GS: Prediction of surgical site infections after major surgery using visible and near-infrared spectroscopy. *Adv Exp Med Biol* 2007; 599:37-44
212. Jackson MR, Gillespie DL, Longenecker EG, Goff JM, Fiala LA, O'Donnell SD, Gomperts ED, Navalta LA, Hestlow T, Alving BM: Hemostatic efficacy of fibrin sealant (human) on expanded poly-tetrafluoroethylene carotid patch angioplasty: a randomized clinical trial. *J Vasc Surg* 1999; 30:461-466
213. Jansen AJ, Andreica S, Claeys M, D'Haese J, Camu F, Jochmans K: Use of tranexamic acid for an effective blood conservation strategy after total knee arthroplasty. *Br J Anaesth* 1999; 83:596-601
214. Jares M, Vanek T, Straka Z, Brucek P: Tranexamic acid reduces bleeding after off-pump coronary artery bypass grafting. *J Cardiovasc Surg* 2003; 44:205-208
215. Jarnagin WR, Gonen M, Maithel SK, Fong Y, D'Angelica MI, DeMatteo RP, Grant F, Wuest D, Kundu K, Blumgart LH, Fischer M: A prospective randomized trial of acute normovolemic hemodilution compared to standard intraoperative management in patients undergoing major hepatic resection. *Ann Surg* 2008; 248:360-369
216. Jassar AS, Ford PA, Haber HL, Isidro A, Swain JD, Bavaria JE, Bridges CR: Cardiac surgery in Jehovah's Witness patients: ten-year experience. *Ann Thorac Surg* 2012; 93:19-25
217. Jensen LS, Andersen AJ, Christiansen PM, Hokland P, Juhl CO, Madsen G, Mortensen J, Moller-Nielsen C, Hanberg-Sorensen F, Hokland M: Postoperative infection and natural

- killer cell function following blood transfusion in patients undergoing elective colorectal surgery. *Br J Surg* 1992; 79:513-516
218. Jensen LS, Kissmeyer-Nielsen P, Wolff B, Qvist N: Randomised comparison of leucocyte-depleted versus buffy-coat-poor blood transfusion and complications after colorectal surgery. *Lancet* 1996; 348:841-845
219. Jewell AE, Akowuah EF, Suvarna SK, Braidley P, Hopkinson D, Cooper G: A prospective randomised comparison of cardiotomy suction and cell saver for recycling shed blood during cardiac surgery. *Eur J Cardiothorac Surg* 2003; 23:633-636
220. Jo KI, Shin JW, Choi TY, Park YJ, Youm W, Kim MJ: Eight-year experience of bloodless surgery at a tertiary care hospital in Korea. *Transfusion* 2013; 53:948-954
221. Johansson PI, Eriksen K, Nielsen SL, Rojkjaer R, Alsbjörn B: Recombinant FVIIa decreases perioperative blood transfusion requirement in burn patients undergoing excision and skin grafting--results of a single centre pilot study. *Burns* 2007; 33:435-440
222. Johnson HD, Morgan MS, Utley JR, Leyand SA, Nguyen-Duy T, Crawley DM: Comparative analysis of recovery of cardiopulmonary bypass residual blood: cell saver vs hemoconcentrator. *J Extra Corpor Technol* 1994; 26:194-199
223. Johnson LB, Plotkin JS, Kuo PC: Reduced transfusion requirements during major hepatic resection with use of intraoperative isovolemic hemodilution. *Am J Surg* 1998; 176:608-611
224. Johnson RG, Thurer RL, Kruskall MS, Sirois C, Gervino EV, Critchlow J, Weintraub RM: Comparison of two transfusion strategies after elective operations for myocardial revascularization. *J Thorac Cardiovasc Surg* 1992; 104:307-314
225. Jones HW, Savage L, White C, Goddard R, Lumley H, Kashif F, Gurusany K: Postoperative autologous blood salvage drains - are they useful in primary uncemented hip and knee arthroplasty? A prospective study of 186 cases. *Acta Orthop Belg* 2004; 70:466-473
226. Jordan D, Delphin E, Rose E: Prophylactic epsilon-aminocaproic acid (EACA) administration minimizes blood replacement therapy during cardiac surgery. *Anesth Analg* 1995; 80:827-829
227. Kahraman S, Altunkaya H, Celebioglu B, Kanbak M, Pasaoglu I, Erdem K: The effect of acute normovolemic hemodilution on homologous blood requirements and total estimated red blood cell volume lost. *Acta Anaesthesiol Scand* 1997; 41:614-617
228. Kajikawa M, Nonami T, Kurokawa T, Hashimoto S, Harada A, Nakao A, Takagi H: Autologous blood transfusion for hepatectomy in patients with cirrhosis and hepatocellular carcinoma: use of recombinant human erythropoietin. *Surgery* 1994; 115:727-734
229. Kajja I, Bimenya GS, Eindhoven GB, ten Duis HJ, Sibinga CT: Surgical blood order equation in femoral fracture surgery. *Transfus Med* 2011; 21:7-12
230. Karger R, Bornmann A, Kretschmer V: Limited utility of algorithms predicting blood transfusions. *Blood Transfus* 2013; 11:426-432
231. Karkouti K, Beattie WS, Wijeyesundera DN, Yau TM, McCluskey SA, Ghannam M, Sutton D, van Rensburg A, Karski J: Recombinant factor VIIa for intractable blood loss after cardiac surgery: A propensity score-matched case-control analysis. *Transfusion* 2005; 45:26-34



232. Karkouti K, McCluskey SA, Syed S, Pazaratz C, Poonawala H, Crowther MA: The influence of perioperative coagulation status on postoperative blood loss in complex cardiac surgery: a prospective observational study. *Anesth Analg* 2010; 110:1533-1540
233. Karkouti K, Wijeyesundera D, Yau T, McCluskey S, Chan C, Wong P-Y, Beattie WS: Influence of erythrocyte transfusion on the risk of acute kidney injury after cardiac surgery differs in anemic and nonanemic patients. *Anesthesiology* 2011; 115:523-530
234. Karkouti K, Wijeyesundera DN, Beattie WS: Risk associated with preoperative anemia in cardiac surgery: A multicenter cohort study. *Circulation* 2008; 117:478-484
235. Karlsson M, Ternström L, Hyllner M, Baghaei F, Flinck A, Skrtic S, Jeppsson A: Prophylactic fibrinogen infusion reduces bleeding after coronary artery bypass surgery. A prospective randomised pilot study. *Thromb Haemost* 2009; 102:137-144
236. Karnezis TA, Stulberg SD, Wixson RL, Reilly P: The hemostatic effects of desmopressin on patients who had total joint arthroplasty. A double-blind randomized trial. *J Bone Joint Surg Am* 1994; 76:1545-1550
237. Karski JM, Teasdale SJ, Norman P, Carroll J, VanKessel K, Wong P, Glynn MF: Prevention of bleeding after cardiopulmonary bypass with high-dose tranexamic acid. Double-blind, randomized clinical trial. *J Thorac Cardiovasc Surg* 1995; 110:835-842
238. Kaspar M, Ramsay MA, Nguyen AT, Cogswell M, Hurst G, Ramsay KJ: Continuous small-dose tranexamic acid reduces fibrinolysis but not transfusion requirements during orthotopic liver transplantation. *Anesth Analg* 1997; 85:281-285
239. Kasper SM, Giesecke T, Limpers P, Sabatowski R, Mehlhorn U, Diefenbach C: Failure of autologous fresh frozen plasma to reduce blood loss and transfusion requirements in coronary artery bypass surgery. *Anesthesiology* 2001; 95:81-86
240. Katsaros D, Petricevic M, Snow NJ, Woodhall DD, Van Bergen R: Tranexamic acid reduces postbypass blood use: a double-blinded, prospective, randomized study of 210 patients. *Ann Thorac Surg* 1996; 61:1131-1135
241. Katz K, Tamary H, Lahav J, Soudry M, Cohen IJ: Increased operative bleeding during orthopaedic surgery in patients with type I Gaucher disease and bone involvement. *Bull Hosp Jt Dis* 1999; 58:188-190
242. Kaufmann CR, Dwyer KM, Crews JD, Dols SJ, Trask AL: Usefulness of thrombelastography in assessment of trauma patient coagulation. *J Trauma* 1997; 42:716-722
243. Kemper RR, Menitove JE, Hanto DW: Cost analysis of intraoperative blood salvage during orthotopic liver transplantation. *Liver Transpl Surg* 1997; 3:513-517
244. Kettelhack C, Hones C, Messinger D, Schlag PM: Randomized multicentre trial of the influence of recombinant human erythropoietin on intraoperative and postoperative transfusion need in anaemic patients undergoing right hemicolectomy for carcinoma. *Br J Surg* 1998; 85:63-67
245. Kikura M, Levy JH, Tanaka KA, Ramsay JG: A double-blind, placebo-controlled trial of epsilon-aminocaproic acid for reducing blood loss in coronary artery bypass grafting surgery. *J Am Coll Surg* 2006; 202:216-222
246. Kilgore ML, Pacifico AD: Shed mediastinal blood transfusion after cardiac operations: a cost-effectiveness analysis. *Ann Thorac Surg* 1998; 65:1248-1254
247. Kjaergard HK, Trumbull HR: Vivostat system autologous fibrin sealant: preliminary study in elective coronary bypass grafting. *Ann Thorac Surg* 1998; 66:482-486

248. Klodell CT, Richardson JD, Bergamini TM, Spain DA: Does cell-saver blood administration and free hemoglobin load cause renal dysfunction? *Am Surg* 2001; 67:44-47
249. Kluba T, Fiedler K, Kunze B, Ipach I, Suckel A: Fibrin sealants in orthopaedic surgery practical experiences derived from use of QUIXIL® in total knee arthroplasty. *Arch Orthop Trauma Surg* 2012; 132:1147-1152
250. Kluger R, Olive DJ, Stewart AB, Blyth CM: Epsilon-aminocaproic acid in coronary artery bypass graft surgery: preincision or postheparin? *Anesthesiology* 2003; 99:1263-1269
251. Koch CG, Li L, Duncan AI, Mihaljevic T, Cosgrove DM, Loop FD, Starr NJ, Blackstone EH: Morbidity and mortality risk associated with red blood cell and blood-component transfusion in isolated coronary artery bypass grafting. *Crit Care Med* 2006; 34:1608-1616
252. Koch CG, Li L, Sessler DI, Figueroa P, Hoeltge GA, Mihaljevic T, Blackstone EH: Duration of red-cell storage and complications after cardiac surgery. *N Engl J Med* 2008; 358:1229-1239
253. Kochamba GS, Pfeffer TA, Sintek CF, Khonsari S: Intraoperative autotransfusion reduces blood loss after cardiopulmonary bypass. *Ann Thorac Surg* 1996; 61:900-903
254. Kohli N, Mallipeddi PK, Neff JM, Sze EH, Roat TW: Routine hematocrit after elective gynecologic surgery. *Obstet Gynecol* 2000; 95:847-850
255. Kosmadakis N, Messaris E, Maris A, Katsaragakis S, Leandros E, Konstadoulakis MM, Androulakis G: Perioperative erythropoietin administration in patients with gastrointestinal tract cancer: prospective randomized double-blind study. *Ann Surg* 2003; 237:417-421
256. Krupp NL, Weinstein G, Chalian A, Berlin Jam Wolf P, Weber RS: Validation of a transfusion prediction model in head and neck cancer surgery. *Arch Otolaryngol Head Neck Surg* 2003; 129:1297-1302
257. Kulier A, Levin J, Moser R, Rumpold-Seitlinger G, Tudor IC, Snyder-Ramos SA, Moehnle P, Mangano DT: Impact of preoperative anemia on outcome in patients undergoing coronary artery bypass graft surgery. *Circulation* 2007; 116:471-479
258. LaPar DJ, Crosby IK, Ailawadi G, Ad N, Choi E, Spiess BD, Rich JB, Kasirajan V, Fonner E Jr, Kron IL, Speir AM, Investigators for the Virginia Cardiac Surgery Quality Initiative: Blood product conservation is associated with improved outcomes and reduced costs after cardiac surgery. *J Thorac Cardiovasc Surg* 2013; 145:796-803
259. Laub GW, Dharan M, Riebman JB, Chen C, Moore R, Bailey BM, Fernandez J, Adkins MS, Anderson W, McGrath LB: The impact of intraoperative autotransfusion on cardiac surgery. A prospective randomized double-blind study. *Chest* 1993; 104:686-689
260. Lee GC, Kicza AM, Liu KY, Nyman CB, Kaufman RM, Body SC: Does rotational thromboelastometry (ROTEM) improve prediction of bleeding after cardiac surgery? *Anesth Analg* 2012; 115:499-506
261. Lemay E, Guay J, Cote C, Roy A: Tranexamic acid reduces the need for allogenic red blood cell transfusions in patients undergoing total hip replacement. *Can J Anaesth* 2004; 51:31-37
262. Lemmer JH Jr, Metzdorff MT, Krause AH Jr, Martin MA, Okies JE, Hill JG: Emergency coronary artery bypass graft surgery in abciximab-treated patients. *Ann Thorac Surg* 2000; 69:90-95

263. LeRoux PD, Elliott JP, Winn HR: Blood transfusion during aneurysm surgery. *Neurosurgery* 2001; 49:1068-1075
264. Levrat A, Gros A, Rugeri L, Inaba K, Floccard B, Negrier C, David JS: Evaluation of rotation thrombelastography for the diagnosis of hyperfibrinolysis in trauma patients. *Br J Anaesth* 2008; 100: 792–797
265. Levy O, Martinowitz U, Oran A, Tauber C, Horoszowski H: The use of fibrin tissue adhesive to reduce blood loss and the need for blood transfusion after total knee arthroplasty. A prospective, randomized, multicenter study. *J Bone Joint Surg Am* 1999; 81:1580-1588
266. Lichtman AD, Carullo V, Minhaj M, Karkouti K: Case 6-2007: Massive intraoperative thrombosis and death after recombinant activated factor VII administration. *J Cardiothorac Vasc Anesth* 2007; 21:897-902
267. Licker M, Sierra J, Kalangos A, Panos A, Diaper J, Ellenberger C: Cardioprotective effects of acute normovolemic hemodilution in patients with severe aortic stenosis undergoing valve replacement. *Transfusion* 2007; 47:341-350
268. Lin PC, Hsu CH, Chen WS, Wang JW: Does tranexamic acid save blood in minimally invasive total knee arthroplasty? *Clin Orthop Relat Res* 2011; 469:1995-2002
269. Lin PC, Hsu CH, Huang CC, Chen WS, Wang JW: The blood-saving effect of tranexamic acid in minimally invasive total knee replacement: is an additional pre-operative injection effective? *J Bone Joint Surg Br* 2012; 94:932-936
270. Lindgren L, Yli-Hankala A, Halme L, Koskimies S, Orko R: Transfusion-related acute lung injury (TRALI) after fresh frozen plasma in a patient with coagulopathy. *Acta Anaesthesiol Scand* 1996; 40:641-644
271. Lisander B, Jonsson R, Nordwall A: Combination of blood-saving methods decreases homologous blood requirements in scoliosis surgery. *Anaesth Intensive Care* 1996; 24:555-558
272. Lison S, Spannagl M, Dietrich W: Hemophilia A in cardiac operations: a model of reduced thrombin generation. *Ann Thorac Surg* 2011; 91:1606-1608
273. Llewellyn-Bennett R, Greenwood R, Benson JR, English R, Turner J, Rayter Z, Winters ZE: Randomized clinical trial on the effect of fibrin sealant on latissimus dorsi donor-site seroma formation after breast reconstruction. *Br J Surg* 2012; 99:1381-1388
274. Lodge JP, Jonas S, Jones RM, Olausson M, Mir-Pallardo J, Soefelt S, Garcia-Valdecasas JC, McAlister V, Mirza DF: Efficacy and safety of repeated perioperative doses of recombinant factor VIIa in liver transplantation. *Liver Transpl* 2005; 11:973-979
275. Lodge JP, Jonas S, Oussoultzoglou E, Malagó M, Jayr C, Cherqui D, Anthuber M, Mirza DF, Kuhlman L, Bechstein WO, Díaz JC, Tartiere J, Eyraud D, Fridberg M, Erhardtsen E, Mimoz O: Recombinant coagulation factor VIIa in major liver resection: a randomized, placebo-controlled, double-blind clinical trial. *Anesthesiology* 2005; 102:269-275
276. Lorenz R, Kienast J, Otto U, Kiehl M, Schreiter D, Haertel S, Barthels M: Successful emergency reversal of phenprocoumon anticoagulation with prothrombin complex concentrate: a prospective clinical study. *Blood Coagul Fibrinolysis* 2007; 18:565-570
277. Loubser PG: Use of acute normovolemic hemodilution for blood conservation during off-pump coronary artery bypass surgery. *Can J Anaesth* 2003; 50:620-621

278. Lozano M, Basora M, Peidro L, Merino I, Segur JM, Pereira A, Salazar F, Cid J, Lozano L, Mazzara R, Macule F: Effectiveness and safety of tranexamic acid administration during total knee arthroplasty. *Vox Sang* 2008; 95:39-44
279. MacIvor D, Nelson J, Triulzi D: Impact of intraoperative red blood cell salvage on transfusion requirements and outcomes in radical prostatectomy. *Transfusion* 2009; 49:1431-1434
280. Mackay Z, Shugufta Q, Din M, Guru AA: Hemodilution in complicated high velocity vascular injuries of limbs. *J Cardiovasc Surg* 1996; 37:217-221
281. Magner D, Ouellette JR, Lee JR, Colquhoun S, Lo S, Nissen NN: Pancreaticoduodenectomy after neoadjuvant therapy in a Jehovah's witness with locally advanced pancreatic cancer: case report and approach to avoid transfusion. *Am Surg* 2006; 72:435-437
282. Mahadevan D, Challand C, Clarke A, Keenan J: Maximum surgical blood ordering schedules for revision lower limb arthroplasty. *Arch Orthop Trauma Surg* 2011; 131:663-667
283. Mahla E, Suarez TA, Bliden KP, Rehak P, Metzler H, Sequeira AJ, Cho P, Sell J, Fan J, Antonino MJ, Tantry US, Gurbel PA: Platelet function measurement-based strategy to reduce bleeding and waiting time in clopidogrel-treated patients undergoing coronary artery bypass graft surgery: the timing based on platelet function strategy to reduce clopidogrel-associated bleeding related to CABG (TARGET-CABG) study. *Cardiovasc Interv* 2012; 5:261-269
284. Maniar RN, Kumar G, Singhi T, Nayak RM, Maniar PR: Most effective regimen of tranexamic acid in knee arthroplasty: a prospective randomized controlled study in 240 patients. *Clin Orthop Relat Res* 2012; 470:2605-2612
285. Marcheix B, Carrier M, Martel C, Cossette M, Pellerin M, Bouchard D, Perrault LP: Effect of pericardial blood processing on postoperative inflammation and the complement pathways. *Ann Thorac Surg* 2008; 85:530-535
286. Martin K, Keller E, Gertler R, Tassani P, Wiesner G: Efficiency and safety of preoperative autologous blood donation in cardiac surgery: a matched-pair analysis in 432 patients. *Eur J Cardiothorac Surg* 2010; 37:1396-1401
287. Martinez V, Monsaingeon-Lion A, Cherif K, Judet T, Chauvin M, Fletcher D: Transfusion strategy for primary knee and hip arthroplasty: impact of an algorithm to lower transfusion rates and hospital costs. *Br J Anaesth* 2007; 99:794-800
288. Marty-Ane CH, Alric P, Picot MC, Picard E, Colson P, Mary H: Ruptured abdominal aortic aneurysm: influence of intraoperative management on surgical outcome. *J Vasc Surg* 1995; 22:780-786
289. Mathiasen RA, Cruz RM: Prospective, randomized, controlled clinical trial of a novel matrix hemostatic sealant in children undergoing adenoidectomy. *Otolaryngol Head Neck Surg* 2004; 131:601-605
290. Matot I, Scheinin O, Jurim O, Eid A: Effectiveness of acute normovolemic hemodilution to minimize allogeneic blood transfusion in major liver resections. *Anesthesiology* 2002; 97:794-800
291. Mawatari M, Higo T, Tsutsumi Y, Shigematsu M, Hotokebuchi T: Effectiveness of autologous fibrin tissue adhesive in reducing postoperative blood loss during total hip arthroplasty: a prospective randomised study of 100 cases. *J Orthop Surg (Hong Kong)* 2006; 14:117-121

292. McConnell JS, Shewale S, Munro NA, Shah K, Deakin AH, Kinninmonth AW: Reduction of blood loss in primary hip arthroplasty with tranexamic acid or fibrin spray. *Acta Orthop* 2011; 82:660-663
293. McGill N, O'Shaughnessy D, Pickering R, Herbertson M, Gill R: Mechanical methods of reducing blood transfusion in cardiac surgery: randomised controlled trial. *BMJ* 2002; 324:1299-303
294. McKenny M, Ryan T, Tate H, Graham B, Young VK, Dowd N: Age of transfused blood is not associated with increased postoperative adverse outcome after cardiac surgery. *Br J Anaesth* 2011; 106:643-649
295. Menges T, Wagner RM, Welters I, Ruwoldt R, Boldt J, Hempelmann G: The role of protein C-thrombomodulin system and fibrinolysis during cardiovascular surgery: influence of acute preoperative plasmapheresis. *J Cardiothorac Vasc Anesth* 1996; 10:482-489
296. Menichetti A, Tritapepe L, Ruvolo G, Speziale G, Cogliati A, Di Giovanni C, Pacilli M, Criniti A: Changes in coagulation patterns, blood loss and blood use after cardiopulmonary bypass: aprotinin vs tranexamic acid vs epsilon aminocaproic acid. *J Cardiovasc Surg* 1996; 37:401-407
297. Mercer KG, Spark JI, Berridge DC, Kent PJ, Scott DJ: Randomized clinical trial of intraoperative autotransfusion in surgery for abdominal aortic aneurysm. *Br J Surg* 2004; 91:1443-1448
298. Mercuriali F, Gualtieri G, Sinigaglia L, Inghilleri G, Biffi E, Vinci A, Colotti MT, Barosi G, Lambertengh Delilieri G: Use of recombinant human erythropoietin to assist autologous blood donation by anemic rheumatoid arthritis patients undergoing major orthopedic surgery. *Transfusion* 1994; 34:501-506
299. Mercuriali F, Inghilleri G, Biffi E, Colotti MT, Vinci A, Oriani G: Epoetin alfa in low hematocrit patients to facilitate autologous blood donation in total hip replacement: a randomized, double-blind, placebo-controlled, dose-ranging study. *Acta Haematol* 1998; 100:69-76
300. Miceli A, Duggan SMJ, Aresu G, de Siena PM, Romeo F, Glauber M, Caputo M, Angelini GD: Combined clopidogrel and aspirin treatment up to surgery increases the risk of postoperative myocardial infarction, blood loss and reoperation for bleeding in patients undergoing coronary artery bypass grafting. *Eur J Cardio-Thorac Surg* 2012; 43:722-728
301. Milic DJ, Perisic ZD, Zivic SS, Stanojkovic ZA, Stojkovic AM, Karanovic ND, Krstic NH, Salinger SS: Prevention of pocket related complications with fibrin sealant in patients undergoing pacemaker implantation who are receiving anticoagulant treatment. *Europace* 2005; 7:374-379
302. Mongan PD, Brown RS, Thwaites BK: Tranexamic acid and aprotinin reduce postoperative bleeding and transfusions during primary coronary revascularization. *Anesth Analg* 1998; 87:258-265
303. Monk TG, Goodnough LT, Brecher ME, Pulley DD, Colberg JW, Andriole GL, Catalona WJ: Acute normovolemic hemodilution can replace preoperative autologous blood donation as a standard of care for autologous blood procurement in radical prostatectomy. *Anesth Analg* 1997; 85:953-958
304. Monte S, Lyons G: Peripartum management of a patient with Glanzmann's thrombasthenia using Thrombelastograph. *Br J Anaesth* 2002; 88:734-738

305. Moonen AF, Knoors NT, van Os JJ, Verburg AD, Pilot P: Retransfusion of filtered shed blood in primary total hip and knee arthroplasty: a prospective randomized clinical trial. *Transfusion* 2007; 47:379-384
306. Moraca RJ, Wanamaker KM, Bailey SH, McGregor WE, Benckart DH, Maher TD, Magovern GJ Jr: Strategies and outcomes of cardiac surgery in Jehovah's Witnesses. *J Card Surg* 2011; 26:135-143
307. Moscardo F, Perez F, de la Rubia J, Balerdi B, Lorenzo JI, Senent ML, Aznar I, Carceller S, Sanz MA: Successful treatment of severe intra-abdominal bleeding associated with disseminated intravascular coagulation using recombinant activated factor VII. *Br J Haematol* 2001; 114:174-176
308. Murkin JM, Adams SJ, Novick RJ, Quantz M, Bainbridge D, Iglesias I, Cleland A, Schaefer B, Irwin G, Fox S: Monitoring brain oxygen saturation during coronary bypass surgery: a randomized prospective study. *Anesth Analg* 2007; 104:51-58
309. Murphy GJ, Allen SM, Unsworth-White J, Lewis CT, Dalrymple-Hay MJ: Safety and efficacy of perioperative cell salvage and autotransfusion after coronary artery bypass grafting: a randomized trial. *Ann Thorac Surg* 2004; 77:1553-1559
310. Murphy GJ, Mango E, Lucchetti V, Battaglia F, Catapano D, Rogers CA, Angelini GD: A randomized trial of tranexamic acid in combination with cell salvage plus a meta-analysis of randomized trials evaluating tranexamic acid in off-pump coronary artery bypass grafting. *J Thorac Cardiovasc Surg* 2006; 132:475-480
311. Murphy GJ, Reeves BC, Rogers CA, Rizve SI, Culliford L, Angelini GD: Increased mortality, postoperative morbidity and cost after red blood cell transfusion in patients having cardiac surgery. *Circulation* 2007; 116:2544-2552
312. Murphy GJ, Rogers CS, Lansdowne WB, Channon I, Alwair H, Cohen A, Caputo M, Angelini GD: Safety, efficacy, and cost of intraoperative cell salvage and autotransfusion after off-pump coronary artery bypass surgery: a randomized trial. *J Thorac Cardiovasc Surg* 2005; 130:20-28
313. Murray D, Pennell B, Olson J: Variability of prothrombin time and activated partial thromboplastin time in the diagnosis of increased surgical bleeding. *Transfusion* 1999; 39:56-62
314. Murray DJ, Pennell BJ, Weinstein SL, Olson JD: Packed red cells in acute blood loss: Dilutional coagulopathy as a cause of surgical bleeding. *Anesth Analg* 1995; 80:336-342
315. Musallam KM, Tamim HM, Richards T, Spahn DR, Rosendaal FR, Habbal A, Khreiss M, Dahdaleh FS, Khavandi K, Sfeir PM, Soweid A, Hoballah JJ, Taher AT, Jamali FR: Preoperative anaemia and postoperative outcomes in non-cardiac surgery: a retrospective cohort study. *Lancet* 2011; 378:1396-1407
316. Myers ER, Clarke-Pearson DL, Olt GJ, Soper JT, Berchuck A: Preoperative coagulation testing on a gynecologic oncology service. *Obstet Gynecol* 1994; 83:438-444
317. Mynster T, Nielsen HJ: The impact of storage time of transfused blood on postoperative infectious complications in rectal cancer surgery. *Scand J Gastroenterol* 2000; 35:212-217
318. Neilipovitz DT, Murto K, Hall L, Barrowman NJ, Splinter WM: A randomized trial of tranexamic acid to reduce blood transfusion for scoliosis surgery. *Anesth Analg* 2001; 93:82-87

319. Netzer G, Shah CV, Iwashyna TJ, Lancken PN, Finkel B, Fuchs B, Guo W, Christie JD: Association of RBC transfusion with mortality in patients with acute lung injury. *Chest* 2007; 132:1116-1118
320. Niranjana G, Asimakopoulou G, Karagounis A, Cockerill G, Thompson M, Chandrasekaran V: Effects of cell saver autologous blood transfusion on blood loss and homologous blood transfusion requirements in patients undergoing cardiac surgery on- versus offcardiopulmonary bypass: a randomised trial. *Eur J Cardiothorac Surg* 2006; 30:271-277
321. Notarnicola A, Moretti L, Martucci A, Spinarelli A, Tafuri S, Pesce V, Moretti B: Comparative efficacy of different doses of fibrin sealant to reduce bleeding after total knee arthroplasty. *Blood Coagul Fibrinolysis* 2012; 23:278-284
322. Nuttall GA, Horlocker TT, Santrach PJ, Oliver WC Jr, Dekutoski MB, Bryant S: Use of the surgical blood order equation in spinal instrumentation and fusion surgery. *Spine* 2000; 25:602-605
323. Nuttall GA, Oliver WC Jr, Beynen FM, Dull JJ, Murray MJ, Nichols WL: Intraoperative measurement of activated partial thromboplastin time and prothrombin time by a portable laser photometer in patients following cardiopulmonary bypass. *Cardiothorac Vasc Anesth* 1993; 7:402-409
324. Nuttall GA, Oliver WC, Beynen FM, Santrach PJ, Strickland RA, Murray MJ: Determination of normal versus abnormal activated partial thromboplastin time and prothrombin time after cardiopulmonary bypass. *J Cardiothorac Vasc Anesth* 1995; 9:355-361
325. Nuttall GA, Oliver WC, Ereth MH, Santrach PJ, Bryant SC, Orszulak TA, Schaff HV: Comparison of blood-conservation strategies in cardiac surgery patients at high risk for bleeding. *Anesthesiology* 2000; 92:674-682
326. Nuttall GA, Oliver WC, Ereth MH, Santrach PJ: Coagulation tests predict bleeding after cardiopulmonary bypass. *J Cardiothorac Vasc Anesth* 1997; 11:815-823
327. Nuttall GA, Oliver WC, Santrach PJ, Bryant S, Dearani JA, Schaff HV, Ereth MH: Efficacy of a simple intraoperative transfusion algorithm for nonerythrocyte component utilization after cardiopulmonary bypass. *Anesthesiology* 2001; 94:773-781
328. Nuttall GA, Santrach PJ, Oliver WC Jr, Ereth MH, Horlocker TT, Cabanela ME, Trousdale RT, Bryant S, Currie TW: A prospective randomized trial of the surgical blood order equation for ordering red blood cells for total hip arthroplasty patients. *Transfusion* 1998; 38:828-833
329. Oefelein MG, Colangelo LA, Rademaker AW, McVary KT: Intraoperative blood loss and prognosis in prostate cancer patients undergoing radical retropubic prostatectomy. *J Urol* 1995; 154:442-447
330. Ogawa S, Szlam F, Chen EP, Nishimura T, Kim H, Roback JD, Levy JH, Tanaka KA: A comparative evaluation of rotation thromboelastometry and standard coagulation tests in hemodilution-induced coagulation changes after cardiac surgery. *Transfusion*. 2012; 52:14-22
331. O'Hara PJ, Hertzler NR, Krajewski LP, Cox GS, Beven EG: Reduction in the homologous blood requirement for abdominal aortic aneurysm repair by the use of preadmission autologous blood donation. *Surgery* 1994; 115:69-76

332. Okuyama M, Ikeda K, Shibata T, Tsukahara Y, Kitada M, Shimano T: Preoperative iron supplementation and intraoperative transfusion during colorectal cancer surgery. *Surg Today* 2005; 35:36-40
333. Oliver WC Jr, Santrach PJ, Danielson GK, Nuttall GA, Schroeder DR, Ereth MH: Desmopressin does not reduce bleeding and transfusion requirements in congenital heart operations. *Ann Thorac Surg* 2000; 70:1923-1930
334. Olsfanger D, Fredman B, Goldstein B, Shapiro A, Jedeikin R: Acute normovolaemic haemodilution decreases postoperative allogeneic blood transfusion after total knee replacement. *Br J Anaesth* 1997; 79:317-321
335. Orlov D, McCluskey SA, Selby R, Yip P, Pendergrast J, Karkouti K: Platelet dysfunction as measured by a point-of-care monitor is an independent predictor of high blood loss in cardiac surgery. *Anesth Analg* 2014; 118:257-263
336. Orpen NM, Little C, Walker G, Crawford EJ: Tranexamic acid reduces early post-operative blood loss after total knee arthroplasty: a prospective randomized controlled trial of 29 patients. *Knee* 2006; 13:106-110
337. Oscarsson A, Gupta A, Fredrikson M, Jarhult J, Nystrom M, Pettersson E, Darvish B, Krook H, Swahn E, Eintrei C: To continue or discontinue aspirin in the perioperative period: a randomized, controlled clinical trial. *Br J Anaesth* 2010; 104:305-312
338. Oswald E, Stalzer B, Heitz E, Weiss M, Schmutz M, Strasak A, Innerhofer P, Haas T: Thromboelastometry (ROTEM) in children: age-related reference ranges and correlations with standard coagulation tests. *Br J Anaesth* 2010; 105:827-835
339. Ott DA, Cooley DA: Cardiovascular surgery in Jehovah's Witnesses. Report of 542 operations without blood transfusion. *JAMA* 1977; 232:1256-1258
340. Ozkisacik E, Islamoglu F, Posacioglu H, Yagdi T, Basarir S, Omay SB, Ozbaran M, Buket S: Desmopressin usage in elective cardiac surgery. *J Cardiovasc Surg (Torino)* 2001; 42:741-747
341. Pagani FD, Polito RJ, Bolling SF: Mitral valve reconstruction in sickle cell disease. *Ann Thorac Surg* 1996; 61:1841-1843
342. Palmer JD, Francis DA, Roath OS, Francis JL, Iannotti F: Hyperfibrinolysis during intracranial surgery: effect of high dose aprotinin. *J Neurosurg Psychiatry* 1995; 58:104-106
343. Palmer T, Wahr JA, O'Reilly M, Greenfield ML: Reducing unnecessary cross-matching: a patient-specific blood ordering system is more accurate in predicting who will receive a blood transfusion than the maximum blood ordering system. *Anesth Analg* 2003; 96:369-375
344. Pang G, Donaldson A: Probable right atrial thrombus immediately after recombinant activated factor VII administration. *Br J Anaesth* 2007; 99:221-225
345. Pasic M, D'Ancona G, Unbehaun A, Hetzer R: Bloodless third complex heart operation in a Jehovah's Witness patient with extremely low preoperative haemoglobin level. *Interact Cardiovasc Thorac Surg* 2012; 14:692-693
346. Pellegrino A, Landoni F, Cormio G, Ferrando P, Lissoni A, Scalabrino S, Mangioni C: Effectiveness of autologous blood transfusion in patients undergoing radical hysterectomy. *Ann Chir Gynaecol* 1995; 84:391-394
347. Penta de Peppo A, Pierri MD, Scafuri A, De Paulis R, Colantuono G, Caprara E, Tomai F, Chiariello L: Intraoperative antifibrinolysis and blood-saving techniques in cardiac surgery. Prospective trial of 3 antifibrinolytic drugs. *Tex Heart Inst J* 1995; 22:231-236



348. Pereboom ITA, de Boer MT, Haagsma EB, Hendriks HGD, Lisman T, Porte RJ: Platelet transfusion during liver transplantation is associated with increased postoperative mortality due to acute lung injury. *Anesth Analg* 2009; 108:1083-1091
349. Phillips LE, McLintock C, Pollock W, Gatt S, Popham P, Jankelowitz G, Ogle R, Cameron PA: Recombinant activated Factor VII in obstetric hemorrhage: experiences from the Australian and New Zealand haemostasis registry. *Anesth Analg* 2009; 109:1908-1915
350. Pinkerton PH, Covens A: Autologous blood transfusion in radical hysterectomy. *Transfus Med* 1996; 6:223-225
351. Pirat A, Sargin D, Torgay A, Arslan G: Identification of preoperative predictors of intraoperative blood transfusion requirement in orthotopic liver transplantation. *Transplant Proc* 2002; 34:2153-2155
352. Pivalizza EG: Perioperative use of the Thrombelastograph in patients with inherited bleeding disorders. *J Clin Anesth* 2003; 15:366-370
353. Planinsic RM, van der Meer J, Testa G, Grande L, Candela A, Porte RJ, Ghobrial M, Isoniemi H, Schelde PB, Erhardtsen E, Klintmalm G, Emre S: Safety and efficacy of a single bolus administration of recombinant factor VIIa in liver transplantation due to chronic liver disease. *Liver Transpl* 2005; 11:895-900
354. Pleym H, Stenseth R, Wahba A, Bjella L, Karevold A, Dale O: Single-dose tranexamic acid reduces postoperative bleeding after coronary surgery in patients treated with aspirin until surgery. *Anesth Analg* 2003; 96:923-928
355. Pleym H, Stenseth R, Wahba A, Bjella L, Tromsdal A, Karevold A, Dale O: Prophylactic treatment with desmopressin does not reduce postoperative bleeding after coronary surgery in patients treated with aspirin before surgery. *Anesth Analg* 2004; 98:578-584
356. Podesta A, Carmagnini E, Parodi E, Dottori V, Crivellari R, Barberis L, Audo A, Lijoi A, Passerone G: Elective coronary and valve surgery without blood transfusion in patients treated with recombinant human erythropoietin (epoetin-alpha). *Minerva Cardioangiolo* 2000; 48:341-347
357. Podesta A, Parodi E, Dottori V, Crivellari R, Passerone GC: Epoetin alpha in elective coronary and valve surgery in Jehovah's Witnesses patients. Experience in 45 patients. *Minerva Cardioangio* 2002; 50:125-131
358. Posacioglu H, Apaydin AZ, Islamoglu F, Calkavur T, Yagdi T, Atay Y, Buket S: Adverse effects of cell saver in patients undergoing ruptured abdominal aortic aneurysm repair. *Ann Vasc Surg* 2002; 16:450-455
359. Potter PS, Waters JH, Burger GA, Mraovic B: Application of cell-salvage during cesarean section. *Anesthesiology* 1999; 90:619-621
360. Powell DM, Chang E, Farrior EH: Recovery from deep-plane rhytidectomy following unilateral wound treatment with autologous platelet gel: a pilot study. *Arch Facial Plast Surg* 2001; 3:245-250
361. Price TH, Goodnough LT, Vogler W, Sacher RA, Hellman RM, Johnston MF, Bolgiano DC, Abels RI: The effect of recombinant human erythropoietin on the efficacy of autologous blood donation in patients with low hematocrits: A multicenter, randomized, double-blind controlled trial. *Transfusion* 1996; 36:29-36
362. Prieto MA, Guash S, Mendez JC, Munoz C, Planas A, Reyes G: Does use of cell saver decrease the inflammatory response in cardiac surgery? *Asian Cardiovasc Thorac Ann* 2013; 21:37-42

363. Pugh SC, Wielogorski AK: A comparison of the effects of tranexamic acid and low-dose aprotinin on blood loss and homologous blood usage in patients undergoing cardiac surgery. *J Cardiothorac Vasc Anesth* 1995; 9:240-244
364. Qvist N, Boesby S, Wolff B, Hansen CP: Recombinant human erythropoietin and hemoglobin concentration at operation and during the postoperative period: reduced need for blood transfusions in patients undergoing colorectal surgery--prospective double-blind placebo-controlled study. *World J Surg* 1999; 23:30-35
365. Rahe-Meyer N, Solomon C, Hanke A, Schmidt DS, Knoerzer D, Hochleitner G, Sørensen B, Hagl C, Pichlmaier M: Effects of fibrinogen concentrate as first-line therapy during major aortic replacement surgery: a randomized, placebo-controlled trial. *Anesthesiology* 2013; 118:40-50
366. Rahe-Meyer N, Solomon C, Winterhalter M, Pipenbrock S, Tanaka K, Haverich A, Pichlmaier M: Thrombolastometry-guided administration of fibrogen concentrate for the treatment of excessive intraoperative bleeding in thoracoabdominal aortic aneurysm surgery. *J Thorac Cardiovasc Surg* 2009; 138:694-702
367. Rainaldi MP, Tazzari PL, Scagliarini G, Borghi B, Conte R: Blood salvage during caesarean section. *Br J Anaesth* 1998; 80:195-198
368. Ralph CJ, Sullivan I, Faulds J: Intraoperative cell salvaged blood as part of a blood conservation strategy in Cesarean section: is fetal red cell contamination important? *Br J Anaesth* 2011; 107:404-408
369. Ramnath AN, Naber HR, de Boer A, Leusink JA: No benefit of intraoperative whole blood sequestration and autotransfusion during coronary artery bypass grafting: results of a randomized clinical trial. *J Thorac Cardiovasc Surg* 2003; 125:1432-1437
370. Rannikko A, Pétas A, Taari K: Tranexamic acid in control of primary hemorrhage during transurethral prostatectomy. *Urology* 2004; 64:955-958
371. Raobaikady R, Redman J, Ball JA, Maloney G, Grounds RM: Use of activated recombinant coagulation factor VII in patients undergoing reconstruction surgery for traumatic fracture of pelvis or pelvis and acetabulum: a double-blind, randomized, placebo-controlled trial. *Br J Anaesth* 2005; 94:586-591
372. Rauh MA, Bayers-Thering M, LaButti RS, Krackow KA: Preoperative administration of epoetin alfa to total joint arthroplasty patients. *Orthopedics* 2002; 25:317-320
373. Ray M, Hatcher S, Whitehouse SL, Crawford S, Crawford R: Aprotinin and epsilon aminocaproic acid are effective in reducing blood loss after primary total hip arthroplasty--a prospective randomized double-blind placebo-controlled study. *J Thromb Haemost* 2005; 3:1421-1427
374. Ray MJ, Hales MM, Brown L, O'Brien MF, Stafford EG: Postoperatively administered aprotinin or epsilon aminocaproic acid after cardiopulmonary bypass has limited benefit. *Ann Thorac Surg* 2001; 72:521-526
375. Rebarber A, Lonser R, Jackson S, Copel JA, Sipes S: The safety of intraoperative autologous blood collection and autotransfusion during cesarean section. *Am J Obstet Gynecol* 1998; 179:715-720
376. Reents W, Babin-Ebell J, Misoph MR, Schwarzkopf A, Elert O: Influence of different autotransfusion devices on the quality of salvaged blood. *Ann Thorac Surg* 1999; 68:58-62

377. Rehm M, Orth V, Kreimeier U, Thiel M, Haller M, Brechtelsbauer H, Finsterer U: Changes in intravascular volume during acute normovolemic hemodilution and intraoperative retransfusion in patients with radical hysterectomy. *Anesthesiology* 2000; 92:657-664
378. Reich DL, Yanakakis MJ, Vela-Cantos FP, DePerio M, Jacobs E: Comparison of bedside coagulation monitoring tests with standard laboratory tests in patients after cardiac surgery. *Anesth Analg* 1999; 88:312-319
379. Reinhöfer M, Brauer M, Franke U, Barz D, Marx G, Lösche W: The value of rotation thromboelastometry to monitor disturbed perioperative haemostasis and bleeding risk in patients with cardiopulmonary bypass. *Blood Coagul Fibrinolysis* 2008; 19:212-219
380. Riou B, Arock M, Guerrero M, Ramos M, Thoreux P, Guillosson JJ, Roy-Camille R, Viars P: Haematological effects of postoperative autotransfusion in spinal surgery. *Acta Anaesthesiol Scand* 1994; 38:336-341
381. Roberts WA, Kirkley SA, Newby M: A cost comparison of allogeneic and preoperatively or intraoperatively donated autologous blood. *Anesth Analg* 1996; 83:129-133
382. Rocha E, Hidalgo F, Llorens R, Melero JM, Arroyo JL, Paramo JA. Randomized study of aprotinin and DDAVP to reduce postoperative bleeding after cardiopulmonary bypass surgery. *Circulation* 1994; 90:921-927
383. Rollo VJ, Hozack WJ, Rothman RH, Chao W, Eng KO: Prospective randomized evaluation of blood salvage techniques for primary total hip arthroplasty. *J Arthroplasty* 1995; 10:532-539
384. Rousou J, Levitsky S, Gonzalez-Lavin L, Cosgrove D, Magilligan D, Weldon C, Hiebert C, Hess P, Joyce L, Bergsland J, Gazzaniga A: Randomized clinical trial of fibrin sealant in patients undergoing re sternotomy or reoperation after cardiac operations. A multicenter study. *J Thorac Cardiovasc Surg* 1989; 97:194-203
385. Rousou JA, Engelman RM, Flack JE 3rd, Deaton DW, Owen SG: Tranexamic acid significantly reduces blood loss associated with coronary revascularization. *Ann Thorac Surg* 1995; 59:671-675
386. Royston D, von Kier S: Reduced haemostatic factor transfusion using heparinase-modified thrombelastography during cardiopulmonary bypass. *Br J Anaesth* 2001; 86:575-578
387. Rubens FD, Boodhwani M, Mesana T, Wozny D, Wells G, Nathan HJ: The cardiotomy trial: a randomized, double-blind study to assess the effect of processing of shed blood during cardiopulmonary bypass on transfusion and neurocognitive function. *Circulation* 2007; 116: I-89-I-97.
388. Rzyman W, Dziadziuszko R, Skokowski J, Wilimski R, Raiter A, Szymanowska A, Jassem J: The influence of blood transfusion on survival in operated non-small cell lung cancer patients. *J Thorac Cardiovasc Surg* 2003; 126:755-760
389. Saager L, Turan A, Dalton JE, Figueroa PI, Sessler DI, Kurz A: Erythrocyte storage duration is not associated with increased mortality in noncardiac surgical patients: A retrospective analysis of 6,994 patients. *Anesthesiology* 2013; 118:51-58
390. Sabatini L, Trecci A, Imarisio D, Uslenghi MD, Bianco G, Scagnelli R: Fibrin tissue adhesive reduces postoperative blood loss in total knee arthroplasty. *J Orthop Traumatol* 2012; 13:145-151
391. Safwat AM, Bush R, Prevec W, Reitan JA: Intraoperative use of platelet-plasmapheresis in vascular surgery. *J Clin Anesth* 2002; 14:10-14

392. Saha SP, Muluk S, Schenk W 3rd, Burks SG, Grigorian A, Ploder B, Presch I, Pavlova BG, Hantak E: Use of fibrin sealant as a hemostatic agent in expanded polytetrafluoroethylene graft placement surgery. *Ann Vasc Surg* 2011; 25:813-822
393. Saha SP, Muluk S, Schenk W 3rd, Dennis JW, Ploder B, Grigorian A, Presch I, Goppelt A: A prospective randomized study comparing fibrin sealant to manual compression for the treatment of anastomotic suture-hole bleeding in expanded polytetrafluoroethylene grafts. *J Vasc Surg* 2012; 56:134-141
394. Sandrelli L, Pardini A, Lorusso R, Sala ML, Licenziati M, Alfieri O: Impact of autologous blood predonation on a comprehensive blood conservation program. *Ann Thorac Surg* 1995; 59:730-755
395. Sarkanović ML, Gvozdenović L, Savić D, Ilić MP, Jovanović G: Autologous blood transfusion in total knee replacement surgery. *Vojnosanit Pregl* 2013; 70:274-278
396. Schaden E, Kimberger O, Kraincuk P, Baron DM, Metnitz PG, Kozek-Langenecker S: Perioperative treatment algorithm for bleeding burn patients reduces allogeneic blood product requirements. *Br J Anaesth* 2012; 109:376-381
397. Schick KS, Fertmann JM, Jauch KW, Hoffmann JN: Prothrombin complex concentrate in surgical patients: retrospective evaluation of vitamin K antagonist reversal and treatment of severe bleeding. *Crit Care* 2009; 13:R191
398. Schmied H, Schiferer A, Sessler DI, Meznik C: The effects of red-cell scavenging, hemodilution, and active warming on allogeneic blood requirements in patients undergoing hip or knee arthroplasty. *Anesth Analg* 1998; 86:387-391
399. Scott SN, Boeve TJ, McCulloch TM, Fitzpatrick KA, Karnell LH: The effects of epoetin alfa on transfusion requirements in head and neck cancer patients: a prospective, randomized, placebo-controlled study. *Laryngoscope* 2002; 112:1221-1229
400. Sebastian C, Romero R, Olalla E, Ferrer C, Garcia-Vallejo JJ, Munoz M: Postoperative blood salvage and reinfusion in spinal surgery: blood quality, effectiveness and impact on patient blood parameters. *Eur Spine J* 2000; 9:458-465
401. Segal N, Puterman M, Rotem E, Niv A, Kaplan D, Kraus M, Brenner H, Nash M, Tal A, Leiberman A: A prospective randomized double-blind trial of fibrin glue for reducing pain and bleeding after tonsillectomy. *Int J Pediatr Otorhinolaryngol* 2008; 72:469-473
402. Senanayake E, Komber M, Nassef A, Massey N, Cooper G: Effective cerebral protection using near-infrared spectroscopy monitoring with antegrade cerebral perfusion during aortic surgery. *J Card Surg* 2012; 27:211-216
403. Serrano FJ, Monux G, Aroca M: Should the cell saver autotransfusion system be routinely used in elective aortic surgery? *Ann Vasc Surg* 2000; 14:663-668
404. Shao YF, Yang JM, Chau GY, Sirivatanauksorn Y, Zhong SX, Erhardtsen E, Nivatvongs S, Lee PH: Safety and hemostatic effects of recombinant activated factor VII in cirrhotic patients undergoing partial hepatectomy; a multicenter, randomized, double-blind, placebo-controlled trial. *Am J Surg* 2006; 191:245-249
405. Shapiro GS, Boachie-Adjei O, Dhawlikar SH, Maier LS: The use of Epoetin alfa in complex spine deformity surgery. *Spine* 2002; 27:2067-2071
406. Shenolikar A, Wareham K, Newington D, Thomas D, Hughes J, Downes M: Cell salvage auto transfusion in total knee replacement surgery. *Transfus Med* 1997; 7:277-280
407. Sheridan DP, Card RT, Pinilla JC, Harding SM, Thomson DJ, Gauthier L, Drotar D: Use of desmopressin acetate to reduce blood transfusion requirements during cardiac surgery in

- patients with acetylsalicylic-acid-induced platelet dysfunction. *Can J Surg* 1994; 37:33-36
408. Shields LE, Smalarz K, Reffigee L, Mugg S, Burdumy TJ, Propst M: Comprehensive maternal hemorrhage protocols improve patient safety and reduce utilization of blood products. *Am J Obstet Gynecol* 2011; 205: 368.e1-368.e8
  409. Shim JK, Choi YS, Oh YJ, Bang SO, Yoo KJ, Kwak YL: Effects of preoperative aspirin and clopidogrel therapy on perioperative blood loss and blood transfusion requirements in patients undergoing off-pump coronary artery bypass graft surgery. *J Thorac Cardiovasc Surg* 2007; 134:59-64
  410. Shore-Lesserson L, Manspeizer HE, DePerio M, Francis S, Vela-Cantos F, Ergin MA: Thromboelastography-guided transfusion algorithm reduces transfusions in complex cardiac surgery. *Anesth Analg* 1999; 88:312-319
  411. Shore-Lesserson L, Reich DL, DePerio M, Silvey G: Autologous platelet-rich plasmapheresis: risk versus benefit in repeat cardiac operations. *Anesth Analg* 1995; 81:229-235
  412. Shore-Lesserson L, Reich DL, Vela-Cantos F, Ammar T, Ergin MA: Tranexamic acid reduces transfusions and mediastinal drainage in repeat cardiac surgery. *Anesth Analg* 1996; 83:18-26
  413. Shulman G, Grecula MJ, Hadjipavlou AG: Intraoperative autotransfusion in hip arthroplasty. *Clin Orthop* 2002; 396:119-30
  414. Shulman G, Solanki DR, Hadjipavlou A: Augmented autologous transfusions in major reconstructive spine surgery. *J Clin Apheresis* 1998; 13:62-68
  415. Simmons JW, White CE, Eastridge BJ, Mace JE, Wade CE, Blackburne LH: Impact of policy change on US Army combat transfusion practices. *J Trauma* 2010; 69 (Suppl 1):S75-S80
  416. Simonian PT, Robinson RP: Titanium contamination of recycled Cell Saver blood in revision hip arthroplasty. *J Arthroplasty* 1995; 10:83-86
  417. Singh J, Ballal MS, Mitchell P, Denn PG: Effects of tranexamic acid on blood loss during total hip arthroplasty. *J Orthop Surg (Hong Kong)* 2010; 18:282-286
  418. Slappendel R, Huvers FC, Benraad B, Novakova I, van Hellenmond G: Use of recombinant factor VIIa to reduce postoperative bleeding after total hip arthroplasty in a patient with cirrhosis and thrombocytopenia. *Anesthesiology* 2002; 96:1525-1527
  419. Slaughter TF, Faghieh F, Greenberg CS, Leslie JB, Sladen RN: The effects of epsilon-aminocaproic acid on fibrinolysis and thrombin generation during cardiac surgery. *Anesth Analg* 1997; 85:1221-1226
  420. Sokolic V, Bukovic D, Fures R, Zadro M, Scuric I, Colak F, Celovec K, Bocan A: Recombinant factor VIIa (rFVIIa) is effective at massive bleeding after caesarean section--a case report. *Coll Antropol* 2002; 26 (Suppl):155-157
  421. So-Osman C, Nelissen R, Te Slaa R, Coene L, Brand R, Brand A: A randomized comparison of transfusion triggers in elective orthopaedic surgery using leucocyte-depleted red blood cells. *Vox Sang* 2010; 98:56-64
  422. Sowade O, Warnke H, Scigalla P, Sowade B, Franke W, Messinger D, Gross J: Avoidance of allogeneic blood transfusions by treatment with epoetin beta (recombinant human erythropoietin) in patients undergoing open-heart surgery. *Blood* 1997; 89:411-418

423. Sowade O, Ziemer S, Sowade B, Franke W, Messinger D, Ziebell E, Scigalla P, Warnke H: The effect of preoperative recombinant human erythropoietin therapy on platelets and hemostasis in patients undergoing cardiac surgery. *J Lab Clin Med* 1997; 129:376-383
424. Spalding GJ, Hartrumpf M, Sierig T, Oesberg N, Kirschke CG, Albes JM: Cost reduction of perioperative coagulation management in cardiac surgery: value of bedside thrombelastography (ROTEM). *Eur J Cardiothorac Surg* 2007; 31:1052-1057
425. Spark JI, Chetter IC, Kester RC, Scott DJ: Allogeneic versus autologous blood during abdominal aortic aneurysm surgery. *Eur J Vasc Endovasc Surg* 1997; 14:482-486
426. Spiess BD, Gillies BSA, Chandler, W, Verrier E: Changes in transfusion therapy and reexploration rate after institution of a blood management program in cardiac surgical patients. *J Cardiothorac Vasc Anesth* 1995; 9:168-173
427. Sreeram GM, Welsby IJ, Sharma AD, Philips-Bute B, Smith PK, Slaughter TF: Infectious complications after cardiac surgery: lack of association with FFP of platelet transfusions. *J Cardiothorac Vasc Anesth* 2005; 19:430-434
428. Stancheva A, Spassov L, Tzatchev K: Correlation between rotation thrombelastometry ROTEM analysis and standard haemostatic parameters during liver transplantation. *Clin Lab* 2011; 57:407-413
429. Stoeckli SJ, Moe KS, Huber A, Schmid S: A prospective randomized double-blind trial of fibrin glue for pain and bleeding after tonsillectomy. *Laryngoscope* 1999; 109:652-655
430. Stover EP, Siegel LC, Hood PA, O'Riordan GE, McKenna TR: Platelet-rich plasma sequestration, with therapeutic platelet yields, reduces allogeneic transfusion in complex cardiac surgery. *Anesth Analg* 2000; 90:509-516
431. Stowell CP, Jones SC, Enny C, Langholff W, Leitz G: An open-label, randomized, parallel-group study of perioperative epoetin alfa versus standard of care for blood conservation in major elective spinal surgery: safety analysis. *Spine (Phila Pa 1976)* 2009; 34:2479-2485
432. Stoye A, Chapin JW, Botha J, Grant W: Bloodless liver transplantation in a Jehovah's Witness. *Int Anesthesiol Clin* 2011; 49:108-115
433. Stratmann G, Russell IA, Merrick SH: Use of recombinant factor VIIa as a rescue treatment for intractable bleeding following repeat aortic arch repair. *Ann Thorac Surg* 2003; 76:2094-2097
434. Strumper D, Weber EW, Gielen-Wijffels S, Van Drumpt R, Bulstra S, Slappendel R, Durieux ME, Marcus MAE: Clinical efficacy of postoperative autologous transfusion of filtered shed blood in hip and knee arthroplasty. *Transfusion* 2004; 44:1567-1571
435. Stuklis RG, O'Shaughnessy DF, Ohri SK: Novel approach to bleeding in patients undergoing cardiac surgery with liver dysfunction. *Eur J Cardiothorac Surg* 2001; 19:219-220
436. Subramanian A, Sagar S, Kumar S, Agrawal D, Albert V, Misra MC: Maximum surgical blood ording schedule in a tertiary trauma center in northern India. *J Emerg Trauma Shock* 2012; 5:312-327
437. Sucker C, Sabel M, Stummer W, Zotz RB, Scharf RE, Gerhardt A: Recombinant factor VIIa for the prophylaxis of perioperative hemorrhage in a patient with congenital factor XI deficiency undergoing brain tumor neurosurgery. *Clin Appl Thromb Hemost* 2008; 14:472-475

438. Sutton SW, Hunley EK, Duncan MA, Rodriguez R, Meyers TP: Sick cell disease and aortic valve replacement: use of cardiopulmonary bypass, partial exchange transfusion, platelet sequestration, and continuous hemofiltration. *Tex Heart Inst J* 1999; 26:283-288
439. Svartholm E, Ammerhagen V, Lanne T: Treatment of bleeding in severe necrotizing pancreatitis with recombinant factor VIIa. *Anesthesiology* 2002; 96:1528
440. Szalay D, Wong D, Lindsay T: Impact of red cell salvage on transfusion requirements during elective abdominal aortic aneurysm repair. *Ann Vasc Surg* 1999; 13:576-581
441. Taghaddomi RJ, Mirzaee A, Attar AS, Shirdel A: Tranexamic acid reduces blood loss in off-pump coronary artery bypass surgery. *J Cardiothorac Vasc Anesth* 2009; 23:312-315
442. Tanaka N, Sakahashi H, Sato E, Hirose K, Ishima T, Ishii S: Timing of the administration of tranexamic acid for maximum reduction in blood loss in arthroplasty of the knee. *J Bone Jt Surg Br* 2001; 83:702-705
443. Tartter PI, Mohandas K, Azar P, Endres J, Kaplan J, Spivack M: Randomized trial comparing packed red blood cell transfusion with and without leukocyte depletion for gastrointestinal surgery. *Am J Surg* 1998; 176:462-466
444. Tawes RL Jr, Duvall TB: Is the "salvaged-cell syndrome" myth or reality? *Am J Surg* 1996; 172:172-174
445. Temeck BK, Bachenheimer LC, Katz NM, Coughlin SS, Wallace RB: Desmopressin acetate in cardiac surgery: a double-blind, randomized study. *South Med J* 1994; 87:611-615
446. Terada N, Arai Y, Matsuta Y, Maekawa S, Okubo K, Ogura K, Matsuda N, Yonei A: Acute normovolemic hemodilution for radical prostatectomy: can it replace preoperative autologous blood transfusion? *Int J Urol* 2001; 8:149-152
447. Terai A, Terada N, Yoshimura K, Ichioka K, Ueda N, Utsunomiya N, Kohei N, Arai Y, Yonei A: Use of acute normovolemic hemodilution in patients undergoing radical prostatectomy. *Urology* 2005; 65:1152-1156
448. Thomas D, Wareham K, Cohen D, Hutchings H: Autologous blood transfusion in total knee replacement surgery. *Br J Anaesth* 2001; 86:669-673
449. Tirosh-Wagner T, Strauss T, Rubinshtein M, Tamarin I, Mishaly D, Paret G, Kenet G: Point of care testing in children undergoing cardiopulmonary bypass. *Pediatr Blood Cancer* 2011; 56:794-798
450. Titlestad IL, Ebbesen LS, Ainsworth AP, Lillevang ST, Qvist N, Georgsen J: Leukocyte-depletion of blood components does not significantly reduce the risk of infectious complications. *Int J Colorectal Dis* 2001; 16:147-153
451. Tobias JD: Synthetic factor VIIa to treat dilutional coagulopathy during posterior spinal fusion in two children. *Anesthesiology* 2002; 96:1522-1525
452. Toy P, Gajic O, Bacchetti P, Looney MR, Gropper MA, Hubmayr R, Lowell CA, Norris PJ, Murphy EL, Weiskopf RB, Wilson G, Koenigsberg M, Lee D, Schuller R, Wu P, Grimes B, Gandhi MJ, Winters JL, Mair D, Hirschler N, Rosen RS, Matthat MA: Transfusion-related acute lung injury: incidence and risk factors. *Blood* 2012; 119:1757-1767
453. Tritapepe L, De Santix V, Vitale D, Nencini C, Pellegrini F, Landoni G, Toscano F, Miraldi F, Pirtropaoli P: Recombinant activated factor VII for refractory bleeding after acute aortic dissection surgery: A propensity score analysis. *Crit Care Med* 2007:1685-1690

454. Triulzi DJ, Gilmor GD, Ness PM, Baumgartner WA, Schultheis LW: Efficacy of autologous fresh whole blood or platelet-rich plasma in adult cardiac surgery. *Transfusion* 1995; 35:627-634
455. Troianos CA, Sypula RW, Lucas DM, D'Amico F, Mathie TB, Desai M, Pasqual RT, Pellegrini RV, Newfeld ML: The effect of prophylactic epsilon-aminocaproic acid on bleeding, transfusions, platelet function, and fibrinolysis during coronary artery bypass grafting. *Anesthesiology* 1999; 91:430-435
456. Trubel W, Gunen E, Wuppinger G, Tschernko E, Gunen-Frank A, Staudacher M, Polterauer P: Recovery of intraoperatively shed blood in aortoiliac surgery: comparison of cell washing with simple filtration. *Thorac Cardiovasc Surg* 1995; 43:165-170
457. Trzebicki J, Flakiewicz E, Kosieradzki M, Blaszczyk B, Kołacz M, Jureczko L, Pacholczyk M, Chmura A, Lagiewska B, Lisik W, Wasiak D, Kosson D, Kwiatkowski A, Lazowski T: The use of thromboelastometry in the assessment of hemostasis during orthotopic liver transplantation reduces the demand for blood products. *Ann Transplant* 2010; 15:19-24
458. Tsuji Y, Kambayashi J, Shiba E, Sakon M, Kawasaki T, Mori T: Effect of recombinant human erythropoietin on anaemia after gastrectomy: a pilot study. *Eur J Surg* 1995; 161:29-33
459. Tuman KJ, Spiess BD, McCarthy RJ, Ivankovich AD: Comparison of viscoelastic measures of coagulation after cardiopulmonary bypass. *Anesth Analg* 1989; 69:69-75
460. Uchino H, Nakamura T, Kuroda S, Houkin K, Murata J, Saito H: Intraoperative dual monitoring during carotid endarterectomy using motor evoked potentials and near-infrared spectroscopy. *World Neurosurg* 2012; 78:651-657
461. Uen WC, Chou YH, Liu CC, Lin SM, Chen TJ: Successful resection of sigmoid colon cancer in a patient with factor XI deficiency. *J Formos Med Assoc* 1998; 97:283-285
462. Urban MK, Beckman J, Gordon M, Urquhart B, Boachie-Adjei O: The efficacy of antifibrinolytics in the reduction of blood loss during complex adult reconstructive spine surgery. *Spine* 2001; 26:1152-1156
463. Vaiman M, Eviatar E, Shlamkovich N, Segal S: Effect of modern fibrin glue on bleeding after tonsillectomy and adenoidectomy. *Ann Otol Rhinol Laryngol* 2003; 112:410-414
464. Vaislic CD, Dalibon N, Ponzio O, Ba M, Jugan E, Lagneau F, Abbas P, Olliver Y, Gaillard D, Baget F, Sportiche M, Chedid A, Chaoul G, Maribas P, Dupuy C, Robine B, Kasanin N, Michon H, Ruat JM, Habis M, Bouharaoua T: Outcomes in cardiac surgery in 500 consecutive Jehovah's Witness patients: 21 year experience. *J Cardiothorac Surg* 2012; 7:95
465. Vamvakas EC, Carven JH: Length of storage of transfused red cells and postoperative morbidity in patents undergoing coronary artery bypass graft surgery. *Transfusion* 2000; 40:101-109
466. Vamvakas EC, Carven JH: Transfusion and postoperative pneumonia in coronary artery bypass graft surgery: effect of the length of storage of transfused red cells. *Transfusion* 1999; 39:701-710
467. Van de Garde EMW, Bras LJ, Heijmen RH, Knibbe CAJ, Van Dongen EPA, Wiltink EHH, Biesma DH: Low-dose recombinant factor VIIa in the management of uncontrolled postoperative hemorrhage in cardiac surgery patients. *J Cardiothorac Vasc Anesth* 2006; 20:573-575



468. van de Watering L, Lorinser J, Versteegh M, Westendord R, Brand A: Effects of storage time of red blood cell transfusions on the prognosis of coronary artery bypass graft patients. *Transfusion* 2006; 46:1712-1718
469. van de Watering LM, Hermans J, Houbiers JG, van den Broek PJ, Bouter H, Boer H, Harvey MS, Huysmans HA, Brand A: Beneficial effects of leukocyte depletion of transfused blood on postoperative complications in patients undergoing cardiac surgery: a randomized clinical trial. *Circulation* 1998; 97:562-568
470. Van Klei WA, Moons KG, Leyssius AT, Knape JT, Rutten CL, Grobee DE: A reduction in type and screen: Preoperative prediction of RBC transfusions in surgery procedures with intermediate transfusion risks. *Br J Anaesth* 2001; 87:250-257
471. van Straten AH, Soliman Hamad MA, van Zundert AA, Mertens EJ, ter Woorst JF, de Wolf AM, Scharnhorst V: Effect of duration of red cell storage on early and late mortality after coronary artery bypass grafting. *J Thorac Cardiovasc Surg* 2011; 141:231-237
472. Vanek T, Jares M, Fajt R, Straka Z, Jirasek K, Kolesar M, Brucek P, Maly M: Fibrinolytic inhibitors in off-pump coronary surgery: a prospective, randomized, double-blind TAP study (tranexamic acid, aprotinin, placebo). *Eur J Cardiothorac Surg* 2005; 28:563-568
473. Veien M, Sorensen JV, Madsen F, Juelsgaard P: Tranexamic acid given intraoperatively reduces blood loss after total knee replacement: a randomized, controlled study. *Acta Anaesthesiol Scand* 2002; 46:1206-1211
474. Vichinsky EP, Haberkern CM, Neumayr L, Earles AN, Black D, Koshy M, Pegelow C, Abboud M, Ohene-Frempong K, Iyer RV: A comparison of conservative and aggressive transfusion regimens in the perioperative management of sickle cell disease. The Preoperative Transfusion in Sickle Cell Disease Study Group. *N Engl J Med* 1995; 333:206-213
475. von Heymann C, Hotz H, Konertz W, Kox WJ, Spies C: Successful treatment of refractory bleeding with recombinant factor VIIa after redo coronary artery bypass graft surgery. *J Cardiothorac Vasc Anesth* 2002; 16:615-616
476. von Heymann C, Redlich U, Jain U, Kastrup M, Schroeder T, Sander M, Grosse J, Ziemer S, Koscielny J, Fonertz WF, Wernecke KD, Spies C: Recombinant activated factor VII for refractory bleeding after cardiac surgery - A retrospective analysis of safety and efficacy. *Crit Care Med* 2005; 33:2241-2246
477. Wajon P, Gibson J, Calcroft R, Hughes C, Thrift B: Intraoperative plateletpheresis and autologous platelet gel do not reduce chest tube drainage or allogeneic blood transfusion after reoperative coronary artery bypass graft. *Anesth Analg* 2001; 93:536-542
478. Waldow T, Krutzsch D, Wils M, Plötze K, Matschke K: Low dose aprotinin and low dose tranexamic acid in elective cardiac surgery with cardiopulmonary bypass. *Clin Hemorheol Microcirc* 2009; 42:269-277
479. Wallis JP, Chapman CE, Orr KE, Clark SC, Forty JR: Effect of WBC reduction of transfused RBCs on postoperative infection rates in cardiac surgery. *Transfusion* 2002; 42:1127-1134
480. Walpoth B, Galliker B, Spirig P, Haeberli A, Rosenmund A, Althaus U, Nydegger UE: Use of epoetin alfa in autologous blood donation programs for patients scheduled for elective cardiac surgery. *Semin Hematol* 1996; 33(2 Suppl 2):75-76
481. Walsh TS, Palmer J, Watson D, Biggin K, Seretny M, Davidson H, Harkness M, Hay A: Multicentre cohort study of red blood cell use for revision hip arthroplasty and factors

- associated with greater risk of allogeneic blood transfusion. *Br J Anaesth* 2012; 108:63-71
482. Wang GJ, Goldthwaite CA Jr, Burks S, Crawford R, Spotnitz WD: Fibrin sealant reduces perioperative blood loss in total hip replacement. *J Long Term Eff Med Implants* 2003; 13:399-411
  483. Wang GJ, Hungerford DS, Savory CG, Rosenberg AG, Mont MA, Burks SG, Mayers SL, Spotnitz WD: Use of fibrin sealant to reduce bloody drainage and hemoglobin loss after total knee arthroplasty: a brief note on a randomized prospective trial. *J Bone Joint Surg Am* 2001; 83-A:1503-1505
  484. Wang JS, Lin CY, Hung WT, O'Connor MF, Thisted RA, Lee BK, Karp RB, Yang MW: Thromboelastogram fails to predict postoperative hemorrhage in cardiac patients. *Ann Thorac Surg* 1992; 53:435-439
  485. Wang MJ, Chen TL, Huang CH, Chao A, Chu SH: Difference in efficacy of the Cell Saver in coronary bypass grafting surgery and cardiac valvular reoperations. *J Formos Med Assoc* 1994; 93:117-121
  486. Wang SC, Shieh JF, Chang KY, Chu YC, Liu CS, Loong CC, Chan KH, Mandell S, Tsou MY: Thromboelastography-guided transfusion decreases intraoperative blood transfusion during orthotopic liver transplantation: randomized clinical trial. *Transplant Proc* 2010; 42:2590-2593
  487. Warkentin TE, Margetts P, Connolly SJ, Lamy A, Ricci C, Eikelboom JW: Recombinant factor VIIa (rFVIIa) and hemodialysis to manage dabigatran-associated postcardiac surgery bleeding. *Blood* 2012; 119:2172-2174
  488. Waters JH, Biscotti C, Potter PS, Phillipson E: Amniotic fluid removal during cell salvage in the cesarean section patient. *Anesthesiology* 2000; 92:1531-1536
  489. Waters JH, Lee JS, Klein E, O'Hara J, Zippe C, Potter PS: Preoperative autologous donation versus cell salvage in the avoidance of allogeneic transfusion in patients undergoing radical retropubic prostatectomy. *Anesth Analg* 2004 ;98:537-542
  490. Waters JH, Lukauskiene E, Anderson ME: Intraoperative blood salvage during cesarean delivery in a patient with beta thalassemia intermedia. *Anesth Analg* 2003; 97:1808-1809
  491. Waters JH, Sprung J: Errors during intraoperative cell salvage because of inappropriate wash solutions. *Anesth Analg* 2001; 93:1483-1485
  492. Weber CF, Gorlinger K, Meininger D, Herrmann E, Bingold T, Moritz A, Cohn LH, Zacharowski K: Point-of-care testing: a prospective, randomized clinical trial of efficacy in coagulopathic cardiac surgery patients. *Anesthesiology* 2012; 117:531-547
  493. Weber CF, Jámbor C, Strasser C, Moritz A, Papadopoulos N, Zacharowski K, Meininger D: Normovolemic modified ultrafiltration is associated with better preserved platelet function and less postoperative blood loss in patients undergoing complex cardiac surgery: a randomized and controlled study. *J Thorac Cardiovasc Surg* 2011; 141:1298-1304
  494. Weber EW, Slappendel R, Hémon Y, Mähler S, Dalén T, Rouwet E, van Os J, Vosmaer A, van der Ark P: Effects of epoetin alfa on blood transfusions and postoperative recovery in orthopaedic surgery. *Eur J Anaesthesiol* 2005; 22:249-257
  495. Wei M, Jian K, Guo Z, Wang L, Jiang D, Zhang L, Tarkka M: Tranexamic acid reduces postoperative bleeding in off-pump coronary artery bypass grafting. *Scand Cardiovasc J* 2006; 40:105-109

496. Weisbach V, Skoda P, Rippel R, Lauer G, Glaser A, Zingsem J, Zimmermann R, Eckstein R: Oral or intravenous iron as an adjuvant to autologous blood donation in elective surgery: a randomized, controlled study. *Transfusion* 1999; 39:465-472
497. Weltert L, D'Alessandro S, Nardella S, Girola F, Bellisario A, Maselli D, De Paulis R: Preoperative very short-term, high-dose erythropoietin administration diminishes blood transfusion rate in off-pump coronary artery bypass: a randomized blind controlled study. *J Thorac Cardiovasc Surg* 2010; 139:621-626
498. White MC, Pryn SJ, Monk CR: Thrombogenic side-effects of recombinant factor VIIa after use in coronary artery bypass surgery. *Anaesth Intensive Care* 2006; 34:664-667
499. Whitson BA, Huddleston SJ, Savik K, Shumway SJ: Bloodless cardiac surgery is associated with decreased morbidity and mortality. *J Card Surg* 2007; 22:373-378
500. Wilhelmi M, Franke U, Cohnert T, Weber P, Kaukemuller J, Fischer S, Wahlers T, Haverich A: Coronary artery bypass grafting surgery without the routine application of blood products: Is it feasible? *Eur J Cardiothorac Surg* 2001; 19:657-661
501. Williams GD, Bratton SL, Riley EC, Ramamoorthy C: Coagulation tests during cardiopulmonary bypass correlate with blood loss in children undergoing cardiac surgery. *J Cardiothorac Vasc Anesth* 1999; 13:398-404
502. Wolowczyk L, Lewis DR, Nevin M, Smith FC, Baird RN, Lamont PM: The effect of acute normovolaemic haemodilution on blood transfusion requirements in abdominal aortic aneurysm repair. *Eur J Vasc Endovasc Surg* 2001; 22:361-364
503. Wolowczyk L, Nevin M, Smith FC, Baird RN, Lamont PM: Haemodilutional effect of standard fluid management limits the effectiveness of acute normovolaemic haemodilution in AAA surgery--results of a pilot trial. *Eur J Vasc Endovasc Surg* 2003; 26:405-411
504. Wong AY, Irwin MG, Hui TW, Fung SK, Fan ST, Ma ES: Desmopressin does not decrease blood loss and transfusion requirements in patients undergoing hepatectomy. *Can J Anaesth* 2003; 50:14-20
505. Wong CJ, Vandervoort MK, Vandervoort SL, Donner A, Zou G, MacDonald JK, Freedman J, Karkouti K, MacDonald SJ, Feagan BG: A cluster-randomized controlled trial of a blood conservation algorithm in patients undergoing total hip joint arthroplasty. *Transfusion* 2007; 47:832-841
506. Wong JC, Torella F, Haynes SL, Dalrymple K, Mortimer AJ, McCollum CN; ATIS Investigators: Autologous versus allogeneic transfusion in aortic surgery: a multicenter randomized clinical trial. *Ann Surg* 2002; 235:145-151
507. Wong Y: Use of prothrombin complex concentrate for vitamin K antagonist reversal before surgical treatment of intracranial hemorrhage. *Clin Med Insights Case Rep* 2011; 4:1-6
508. Wu CC, Ho WM, Cheng SB, Yeh DC, Wen MC, Liu TJ, P'eng FK: Perioperative parenteral tranexamic acid in liver tumor resection: a prospective randomized trial toward a "blood transfusion"-free hepatectomy. *Ann Surg* 2006; 243:173-180
509. Wurnig C, Schatz K, Noske H, Hemon Y, Dahlberg G, Josefsson G, Milbrink J, Hamard C; Collaborative Study Group: Subcutaneous low-dose epoetin beta for the avoidance of transfusion in patients scheduled for elective surgery not eligible for autologous blood donation. *Eur Surg Res* 2001; 33:303-310
510. Yaffee DW, Smith DE, Ursomanno PA, Hill FT, Galloway AC, DeAnda A, Grossi EA: Management of blood transfusion in aortic valve surgery: impact of a blood conservation strategy. *Ann Thorac Surg* 2014; 97:95-101

511. Yamada T, Yamashita Y, Terai Y, Ueki M: Intraoperative blood salvage in abdominal uterine myomectomy. *Int J Gynaecol Obstet* 1997; 56:141-145
512. Yamasaki S, Masuhara K, Fuji T: Tranexamic acid reduces blood loss after cementless total hip arthroplasty: Prospective randomized study in 40 cases. *Int Orthop* 2004; 28:69-73
513. Yap CH, Lau L, Krishnaswamy M, Gaskell M, Yui M: Age of transfused red cells and early outcomes after cardiac surgery. *Ann Thorac Surg* 2008; 86:554-559
514. Yasuda H, Ohto H, Yamaguchi O, Sakuma S, Suzuki T, Mita M, Tsuneyama H, Uchikawa M: Three episodes of delayed hemolytic transfusion reactions due to multiple red cell antibodies, anti-Di, anti-Jk and anti-E. *Transfus Sci* 2000; 23:107-112
515. Yokomuro M, Ebine K, Shiroma K, Tamura S, Kumabe S, Ohtuki M, Suzuki H, Uchida S: Safety and efficacy of autologous platelet transfusion in cardiac surgery: comparison of cryopreservation, blood collection on the day before surgery, and blood collection during surgery. *Cryobiology* 1999; 38:236-242
516. Yoo YC, Shim JK, Kim JC, Jo YY, Lee JH, Kwak YL: Effect of single recombinant human erythropoietin injection on transfusion requirements in preoperatively anemic patients undergoing valvular heart surgery. *Anesthesiology* 2011; 115:929-937
517. Yu L, Gu T, Song L, Shi E, Fang Q, Wang C, Zhao J: Fibrin sealant provides superior hemostasis for sternotomy compared with bone wax. *Ann Thorac Surg* 2012; 93:641-644
518. Zabeeda D, Medalion B, Sverdlov M, Ezra S, Schachner A, Ezri T, Cohen AJ: Tranexamic acid reduces bleeding and the need for blood transfusion in primary myocardial revascularization. *Ann Thorac Surg* 2002; 74:733-738
519. Zufferey PJ, Miquet M, Quenet S, Martin P, Adam P, Albaladejo P, Mismetti P, Molliex S: Tranexamic acid in hip fracture surgery: a randomized controlled trial. *Br J Anaesth* 2010; 104:23-30
520. Zupancic Salek S, Sokolic V, Viskovic T, Sanjug J, Simic M, Kastelan M: Successful use of recombinant factor VIIa for massive bleeding after caesarean section due to HELLP syndrome. *Acta Haematol* 2002; 108:162-163