

Supplementary Material

Neurophysiologic Correlates of Ketamine Sedation and Anesthesia: a High-Density Electroencephalography Study in Healthy Volunteers

Phillip E. Vlisides, M.D.; Tarik Bel-Bahar, Ph.D.; UnCheol Lee, Ph.D.; Duan Li, Ph.D.;
Hyoungkyu Kim, Ph.D.; Ellen Janke, M.D.; Vijay Tarnal, M.D.; Adrian B. Pichurko, M.D.; Amy
M. McKinney, M.A.; Bryan S. Kunkler, B.S.; Paul Picton, M.B. Ch.B., M.R.C.P., F.R.C.A.;
George A. Mashour, M.D., Ph.D.

Department of Anesthesiology, University of Michigan Medical School

(PEV, TB, UL, DL, HKK, EJ, VT, AM, BK, PP, GAM)

Center for Consciousness Science, University of Michigan Medical School

(PEV, TB, UL, HKK, DL, GAM)


Department of Anesthesiology, Northwestern University Feinberg School of Medicine

(AP)

Neuroscience Graduate Program, University of Michigan Medical School

(GAM)

Corresponding Author: George A. Mashour, M.D., Ph.D. at gmashour@med.umich.edu



Supplemental Digital Content 1: 128-channel EEG Hydrocel Geodesic Sensor Net

(Electrical Geodesics, Inc.) utilized in this study with international 10-10 electrode system equivalents. Blue shading represents the eight frontal channels centered on Fz used for frontal channel analyses.