

## 2022 American Society of Anesthesiologists Practice Guidelines for Management of the Difficult Airway: Supplement 1 - Bibliography in Alphabetical Order

1. Abbasijahromi A, Sanie Jahromi MS, Farzaneh M, Javadpour S, Montaseri MA, Kalani N: Success in difficult airway managements with video laryngoscope after two failures in Intubation with Macintosh and McCoy Laryngoscope: A case report. *Iran Red Cresc Med J* 2017; 19
2. Abdel Naim HE, Mohamed SAR, Soaida SM, Eltrabily HHA: The importance of neck circumference to thyromental distance ratio (NC/TM) as a predictor of difficult intubation in obstructive sleep apnea (OSA) patients. *Egypt J Anaesth* 2019; 30:219-25
3. Abdelhady BS, Elrabiey MA, Abd Elrahman AH, Mohamed EE: Ultrasonography versus conventional methods (Mallampati score and thyromental distance) for prediction of difficult airway in adult patients. *Egypt J Anaesth* 2020; 36:83-9
4. Abdellatif AA, Ali MA: GlideScope videolaryngoscope versus flexible fiberoptic bronchoscope for awake intubation of morbidly obese patient with predicted difficult intubation. *Middle East J Anaesthesiol* 2014; 22:385-92
5. Abraham S, Himarani J, Mary Nancy S, Shanmugasundaram S, Krishnakumar Raja VB: Ultrasound as an assessment method in predicting difficult intubation: A prospective clinical study. *J Maxillofac Oral Surg* 2018; 17:563-9
6. Abramson SI, Holmes AA, Hagberg CA: Awake insertion of the Bonfils retromolar intubation fiberscope in five patients with anticipated difficult airways. *Anesth Analg* 2008; 106:1215-7
7. Abrons RO, Vansickle RA, Ouanes JP: Seldinger technique for nasal intubation: a case series. *J Clin Anesth* 2016; 34:609-11
8. Acar HV, Uysal HY, CeyhanKaya A, Ceyhan A, Dikmen B: Does the STOP-Bang, an obstructive sleep apnea screening tool, predict difficult intubation? *Eur Rev Med Pharmacol Sci* 2014; 18:1869-74
9. Adi O, Fong CP, Sum KM, Ahmad AH: Usage of airway ultrasound as an assessment and prediction tool of a difficult airway management. *Am J Emerg Med* 2020
10. Ads A, Auerbach F, Ryan K, El-Ganzouri AR: Air-Q laryngeal airway for rescue and tracheal intubation. *J Clin Anesth* 2016; 32:108-11
11. Agro F, Brimacombe J, Brain AI, Marchionni L, Cataldo R: The intubating laryngeal mask for maxillo-facial trauma. *Eur J Anaesthesiol* 1999; 16:263-4
12. Agrò F, Totonelli A, Gherardi S: Planned lightwand intubation in a patient with a known difficult airway. *Can J Anaesth* 2004; 51:1051-2
13. Ahmadi K, Ebrahimi M, Hashemian AM, Sarshar S, Rahimi-Movaghar V: GlideScope video laryngoscope for difficult intubation in emergency patients: A quasi-randomized controlled trial. *Acta Med Iran* 2015; 53:738-42
14. Ahsan B, Kamali G, Nessleri K: The laryngeal mask airway for difficult airway in temporomandibular joint ankylosis- A case report. *Middle East J Anaesthesiol* 2012; 21:639-42
15. Akhlaghi M, Abedinzadeh M, Ahmadi A, Heidari Z: Predicting difficult laryngoscopy and intubation with laryngoscopic exam test: A new method. *Acta Medica Iranica* 2017; 55:453-8
16. Aktas S, Atalay YO, Tugrul M: Predictive value of bedside tests for difficult intubations. *Eur Rev Med Pharmacol Sci* 2015; 19:1595-9
17. Al Harbi M, Thomas J, Khalil Hassan N, Said Hassanin N, Wannous S, Abouras C, Al Harthi A, Dimitrou V: Anesthetic management of advanced stage ludwig's angina: A case report and review with emphasis on compromised airway management. *Middle East J Anaesthesiol* 2016; 23:665-73
18. Alain TI, Drissa BS, Flavien K, Serge I, Idriss T: Video-laryngoscopy introduction in a Sub-Saharan national teaching hospital: luxury or necessity? *Pan African Med J* 2015; 22:381

19. Alessandri F, Antenucci G, Piervincenzi E, Buonopane C, Bellucci R, Andreoli C, Alunni Fegatelli D, Ranieri MV, Bilotta F: Ultrasound as a new tool in the assessment of airway difficulties: An observational study. *Eur J Anaesthesiol* 2019; 36:509-15
20. Ali MA, Qamar-ul-Hoda M, Samad K: Comparison of upper lip bite test with Mallampati test in the prediction of difficult intubation at a tertiary care hospital of Pakistan. *J Pak Med Assoc* 2012; 62:1012-5
21. Ali QE, Amir SH, Ahmed S: A comparative evaluation of King Vision video laryngoscope (channelled blade), McCoy, and Macintosh laryngoscopes for tracheal intubation in patients with immobilized cervical spine. *Sri Lankan J Anaesthesiol* 2017; 25:70
22. Ali QE, Amir SH, Siddiqui OA, Pal K: King vision video laryngoscope: A suitable device for severe ankylosing spondylitis. *Egypt J Anaesth* 2017; 33:129-31
23. Ali QE, Amir, SH, Siddiqui OA, Nadeem A, Azhar A Z: Airtraq® optical laryngoscope for tracheal intubation in patients with severe ankylosing spondylitis: A report of two cases. *Indian J Anaesth* 2012; 56:165-167
24. Ali QE, Siddiqui OA, Amir SH, Azhar AZ, Ali K: Airtraq® Optical laryngoscope for tracheal intubation in a patient with giant lipoma at the nape: A case report. *Revista Brasileira de Anestesiologia* 2012; 62:736-740
25. Allencherril JP, Joseph L: Soft palate trauma induced during GlideScope intubation. *J Clin Anesth* 2016; 35:278-80
26. Aloqab S, Chandrashekhariah M, Shah V, Adeel S: Difficult airway management: Burning no bridges. *Sri Lankan J Anaesthesiol* 2019; 27:169-71
27. Altun D, Demir G, Ayhan A, Türköz A: Successful anesthetic and airway management in Coffin-Siris syndrome with congenital heart disease: Case report. *Egypt J Anaesth* 2016; 32:593-6
28. Altun D, Kara H, Bozbora E, Ali A, Dinc T, Sonmez S, Buget M, Aydemir L, Basaran B, Tugrul M, Camci E: The role of indirect laryngoscopy, clinical and ultrasonographic assessment in prediction of difficult airway. *Laryngoscope* 2021; 131:E555-60
29. Alvis BD, King AB, Hester D, Hughes CG, Higgins MS: Randomized controlled pilot trial of the rigid and flexing laryngoscope versus the fiberoptic bronchoscope for intubation of potentially difficult airway. *Minerva Anesthesiol* 2015; 81:946-50
30. Amaniti A, Papakonstantinou P, Gkinas D, Dalakakis I, Papapostolou E, Nikopoulou A, Tsatali M, Zarogoulidis P, Sapalidis K, Kosmidis C, Koulouris C, Giannakidis D, Romanidis K, Oikonomou P, Michalopoulos N, Ioannidis A, Tsakiridis K, Vagionas A, Kesisoglou I, Grosomanidis V: Comparison of laryngoscopic views between C-MAC TM and conventional laryngoscopy in patients with multiple preoperative prognostic criteria of difficult intubation: An observational cross-sectional study. *Medicina* 2019; 55:27
31. Amata AO: Difficult airway management and suspected malignant hyperthermia in a child with Cri Du Chat Syndrome. *Saudi J Anaesth* 2019; 13:81-3
32. Ambesh SP, Singh N, Rao PB, Gupta D, Singh PK, Singh U: A combination of the modified Mallampati score, thyromental distance, anatomical abnormality, and cervical mobility (M-TAC) predicts difficult laryngoscopy better than Mallampati classification. *Acta Anaesthesiol Taiwan* 2013; 51:58-62
33. Andruszkiewicz P, Wojtczak J, Sobczyk D, Stach O, Kowalik I: Effectiveness and validity of sonographic upper airway evaluation to predict difficult laryngoscopy. *J Ultrasound Med* 2016; 35:2243-52
34. Aoyama K, Nagaoka E, Takenaka I, Kadoya T: The McCoy laryngoscope expands the laryngeal aperture in patients with difficult intubation. *Anesthesiology* 2000; 92:1855-6
35. Arevalo-Ludena J, Arcas-Bellas JJ, Alvarez-Rementeria R, Alameda LE: Fiberoptic-guided intubation after insertion of the i-gel airway device in spontaneously breathing patients with difficult airway predicted: a prospective observational study. *J Clin Anesth* 2016; 35:287-92
36. Asai T, Hirose T, Shingu K: Failed tracheal intubation using a laryngoscope and intubating laryngeal mask. *Can J Anaesth* 2000; 47:325-8

37. Asai T, Liu EH, Matsumoto S, Hirabayashi Y, Seo N, Suzuki A, Toi T, Yasumoto K, Okuda Y: Use of the Pentax-AWS in 293 patients with difficult airways. *Anesthesiology* 2009; 110:898-904
38. Asai T, Matsumoto H, Shingu K: Awake tracheal intubation through the intubating laryngeal mask. *Can J Anaesth* 1999; 46:182-4
39. Asai T, Matsumoto S, Shingu K: Use of the McCoy laryngoscope or fingers to facilitate fibrescope-aided tracheal intubation. *Anaesthesia* 1998; 53:903-5
40. Asai T, Nagata A, Shingu K: Awake tracheal intubation through the laryngeal mask in neonates with upper airway obstruction. *Paediatr Anaesth* 2008; 18:77-80
41. Asai T, Shingu K: Tracheal intubation through the intubating laryngeal mask in a patient with a fixed flexed neck and deviated larynx. *Anaesthesia* 1998; 53:1199-201
42. Asai T, Shingu K: Tracheal intubation through the intubating laryngeal mask in patients with unstable necks. *Acta Anaesthesiol Scand* 2001; 45:818-22
43. Asai T: Intubating laryngeal mask airway after failed insertion of a Classic laryngeal mask. *Anaesthesia* 2006; 61:303
44. Asai T: Pentax-AWS videolaryngoscope for awake nasal intubation in patients with unstable necks. *Br J Anaesth* 2010; 104:108-11
45. Asghar A, Shamim F, Aman A: Fiberoptic intubation in a paediatric patient with severe temporomandibular joint (TMJ) ankylosis. *J Coll Physicians Surg Pak* 2012; 22:783-5
46. Ayoub E, Tohme J, Lutfallah AA, Jabbour H, Chalhoub V, Naccache N: Intractable course of a submandibular abscess following difficult endotracheal intubation a case report. *J Med Libanais* 2019; 67:103-6
47. Aziz MF, Brambrink AM, Healy DW, Willett AW, Shanks A, Tremper T, Jameson L, Ragheb J, Biggs DA, Paganelli WC, Rao J, Epps JL, Colquhoun DA, Bakke P, Kheterpal S: Success of intubation rescue techniques after failed direct laryngoscopy in adults: A retrospective comparative analysis from the multicenter perioperative outcomes group. *Anesthesiology* 2016; 125:656-66
48. Aziz MF, Dillman D, Fu R, Brambrink AM: Comparative effectiveness of the C-MAC video laryngoscope versus direct laryngoscopy in the setting of the predicted difficult airway. *Anesthesiology* 2012; 116:629-36
49. Aziz MF, Healy D, Kheterpal S, Fu RF, Dillman D, Brambrink AM: Routine clinical practice effectiveness of the Glidescope in difficult airway management: an analysis of 2,004 Glidescope intubations, complications, and failures from two institutions. *Anesthesiology* 2011; 114:34-41
50. Badhe VK, Deogaonkar SG, Tambe MV, Singla A, Shidhaye RV: Clinical comparison of five different predictor tests for difficult intubation. *Anaesth Pain Intensive Care* 2014; 18:31-7
51. Badheka J, Doshi P, Vyas A, Kacha N, Parmar V: Comparison of upper lip bite test and ratio of height to thyromental distance with other airway assessment tests for predicting difficult endotracheal intubation. *Indian J Crit Care Med* 2016; 20:3-8
52. Bae HM, Yoon JR, Yoo JH, Han YJ, Park YJ: A vallecular cyst in a patient with deep neck infection causing difficult airway management. *Korean J Anesthesiol* 2014; 67:S21-2
53. Baig MM, Khan FH: To compare the accuracy of prayer's sign and Mallampatti test in predicting difficult intubation in diabetic patients. *J Pak Med Assoc* 2014; 64:879-83
54. Bamgbade OA: The use of intubating lightwand in difficult airway patients with limited management options. *Niger Postgrad Med J* 2017; 24:187-90
55. Barch B, Rastatter J, Jagannathan N: Difficult pediatric airway management using the intubating laryngeal airway. *Int J Pediatr Otorhinolaryngol* 2012; 76:1579-82
56. Beilin B, Yardeni IZ, Smolyarenko V, Zeidel A, Ram E, Mayburd E: Comparison of the Flexiblade levering laryngoscope with the English Macintosh laryngoscope in patients with a poor laryngoscopic view. *Anaesthesia* 2005; 60:400-5

57. Berthelsen P, Prytz S, Jacobsen E: Two-stage fiberoptic nasotracheal intubation in infants: a new approach to difficult pediatric intubation. *Anesthesiology* 1985; 63:457-8
58. Beshey BN, Helmy TA, Asaad HS, Ibrahim EEDM: Emergency percutaneous tracheotomy in failed intubation. *Egypt J Chest Dis Tuberc* 2014; 63:939-45
59. Bhaktavar J, Gupta P: Comparison of acromio-axillo-suprasternal notch index with upper lip bite test and ratio of height to thyromental distance for prediction of difficult intubation: A prospective study. *Sri Lankan J Anaesthesiol* 2020; 28:119-24
60. Bhat R, Mane RS, Patil MC, Suresh SN: Fiberoptic intubation through laryngeal mask airway for management of difficult airway in a child with Klippel-Feil syndrome. *Saudi J Anaesth* 2014; 8:412-4
61. Bhatnagar S, Mishra S, Jha RR, Singhal AK, Bhatnagar N: The LMA Fastrach facilitates fiberoptic intubation in oral cancer patients. *Can J Anaesth* 2005; 52:641-5
62. Bhola N, Jadhav A, Borle R, Khemka G, Ajani AA: Awake endotracheal retrograde intubation in restricted mouth opening: a 'J'-tipped guide wire technique--A retrospective study. *Oral Maxillofac Surg* 2014; 18:393-6
63. Binar M, Arslan F, Aydin U: Another cause of difficult airway in an elderly patient: Tongue-base abscess. *Gerodontology* 2018; 35:155-8
64. Bindra A, Prabhakar H, Bithal PK, Singh GP, Chowdhury T: Predicting difficult laryngoscopy in acromegalic patients undergoing surgery for excision of pituitary tumors: A comparison of extended Mallampati score with modified Mallampati classification. *J Anaesthesiol Clin Pharmacol* 2013; 29:187-90
65. Bittar D: Respiratory obstruction associated with induction of general anesthesia in a patient with mediastinal Hodgkin's disease. *Anesth Analg* 1975; 54:399-403
66. Blanco G, Melman E, Cuairan V, Moyao D, Ortiz-Monasterio F: Fiberoptic nasal intubation in children with anticipated and unanticipated difficult intubation. *Paediatr Anaesth* 2001; 11:49-53
67. Borland LM, Casselbrant M: The Bullard laryngoscope. A new indirect oral laryngoscope (pediatric version). *Anesth Analg* 1990; 70:105-8
68. Bouroche G, Motamed C, de Guibert JM, Hartl D, Bourgain JL: Rescue transtracheal jet ventilation during difficult intubation in patients with upper airway cancer. *Anaesth Crit Care Pain Med* 2018; 37:539-44
69. Brenman S, Gupta S, Tseeng S: Successful retrograde intubation after failed fiberoptic intubation and percutaneous cricothyrotomy. *J Emerg Med* 2017; 53:550-3
70. Brimacombe J, Keller C: Awake fiberoptic-guided insertion of the ProSeal laryngeal mask airway. *Anaesthesia* 2002; 57:719
71. Brimacombe JR: Difficult airway management with the intubating laryngeal mask. *Anesth Analg* 1997; 85:1173-5
72. Brodsky JB, Lemmens HJ, Brock-Utne JG, Vierra M, Saidman LJ: Morbid obesity and tracheal intubation. *Anesth Analg* 2002; 94:732-6
73. Bruserud Ø, Wendelbo Ø, Vetti N, Goplen FK, Johansen S, Reikvam H: Critical upper airway obstruction as the first symptom of acute myeloid leukemia - An anesthesiologic reminder. *Clin Prac* 2020; 10:34-6
74. Buckland RW, Pedley J: Lingual thyroid--a threat to the airway. *Anaesthesia* 2000; 55:1103-5
75. Budania L S, Eeshwar MV, Reddy VC, Gaude Y K, Anand S, Sharma A: Epidermolysis Bullosa – An anaesthetic challenge. *Sri Lankan J Anaesthesiol* 2019; 27:163-165
76. Burjek NE, Nishisaki A, Fiadjoe JE, Adams HD, Peeples KN, Raman VT, Olomu PN, Kovatsis PG, Jagannathan N, Hunyady A, Bosenberg A, Tham S, Low D, Hopkins P, Glover C, Olutoye O, Szmuk P, McCloskey J, Dalesio N, Koka R, Greenberg R, Watkins S, Patel V, Reynolds P, Matuszczak M, Jain R, Khalil S, Polaner D, Zieg J, Szolnoki J, Sathyamoorthy K, Taicher B, Riveros Perez NR, Bhattacharya S, Bhalla T, Stricker P, Lockman J, Galvez J, Rehman M, Von Ungern-Sternberg B, Sommerfield D, Soneru C,

- Chiao F, Richtsfeld M, Belani K, Sarmiento L, Mireles S, Bilen Rosas G, Park R, Peyton J, Pe DICI: Videolaryngoscopy versus fiber-optic Intubation through a supraglottic airway in children with a difficult airway: An analysis from the Multicenter Pediatric Difficult Intubation Registry. *Anesthesiology* 2017; 127:432-40
77. Butler PJ, Dhara SS: Prediction of difficult laryngoscopy: an assessment of the thyromental distance and Mallampati predictive tests. *Anaesth Intensive Care* 1992; 20:139-42
  78. Capistrano-Baruh E, Wenig B, Steinberg L, Stegnjajic A, Baruh S: Laryngeal web: A cause of difficult endotracheal intubation. *Anesthesiology* 1982; 57:123-5
  79. Cattano D, Killoran PV, Cai C, Katsiampoura AD, Corso RM, Hagberg CA: Difficult mask ventilation in general surgical population: Observation of risk factors and predictors. *F1000Res* 2014; 3
  80. Cattano D, Panicucci E, Paolicchi A, Forfori F, Giunta F, Hagberg C: Risk factors assessment of the difficult airway: an italian survey of 1956 patients. *Anesth Analg* 2004; 99:1774-9
  81. Cavus E, Neumann T, Doerges V, Moeller T, Scharf E, Wagner K, Bein B, Serocki G: First clinical evaluation of the C-MAC D-Blade videolaryngoscope during routine and difficult intubation. *Anesth Analg* 2011; 112:382-5
  82. Çelik F, Tokgoz O, Dogan E, Guzel A, Ciftci T, Tufek A: Retrograde intubation in the patient with cystic tumor located at the base of tongue. *Middle East J Anaesthesiol* 2013; 22:333-6
  83. Chae JS, Woo JH, Kim CH, Chun EH, Baik HJ, Choi MH: Endotracheal intubation using McGrath videolaryngoscope in Klippel-Feil syndrome. *EWHA Med J* 2018; 41:86-9
  84. Chan CS: Anaesthetic management during repair of tracheo-oesophageal fistula. *Anaesthesia* 1984; 39:158-60
  85. Chan SMM, Wong WY, Lam SKT, Wong OF, Law WSS, Shiu WYY, Mak PYE: Use of ultrasound to predict difficult intubation in Chinese population by assessing the ratio of the pre-epiglottis space distance and the distance between epiglottis and vocal folds. *Hong Kong J Emerg Med* 2018; 25:152-9
  86. Char DS, Gipp M, Boltz MG, Williams GD: Case report: airway and concurrent hemodynamic management in a neonate with oculo-auriculo-vertebral (Goldenhar) syndrome, severe cervical scoliosis, interrupted aortic arch, multiple ventricular septal defects, and an unstable cervical spine. *Paediatr Anaesth* 2012; 22:932-4
  87. Chaudhuri S, Duggappa AK, Mathew S, Venkatesh S: Safe intubation in Morquio-Brailsford syndrome: A challenge for the anesthesiologist. *J Anaesthesiol Clin Pharmacol* 2013; 29:258-61
  88. Cheng T, Wang LK, Wu HY, Yang XD, Zhang X, Jiao L: Shikani optical stylet for awake nasal intubation in patients undergoing head and neck surgery. *Laryngoscope* 2021; 131:319-25
  89. Chhina AK, Jain R, Gautam PL, Garg J, Singh N, Grewal A: Formulation of a multivariate predictive model for difficult intubation: A double blinded prospective study. *J Anaesthesiol Clin Pharmacol* 2018; 34:62-7
  90. Chisholm DG, Calder I: Experience with the McCoy laryngoscope in difficult laryngoscopy. *Anaesthesia* 1997; 52:906-908
  91. Choi CG, Yang KH, Jung JK, Han JU, Lee CS, Cha YD, Song JH: Endotracheal intubation using i-gel(R) and lightwand in a patient with difficult airway: a case report. *Korean J Anesthesiol* 2015; 68:501-4
  92. Choi EK, Kim JE, Soh SR, Kim CK, Park WK: Usefulness of a Cook(R) airway exchange catheter in laryngeal mask airway-guided fiberoptic intubation in a neonate with Pierre Robin syndrome -A case report. *Korean J Anesthesiol* 2013; 64:168-71

93. Choi GS, Park SI, Lee EH, Yoon SH: Awake Glidescope® intubation in a patient with a huge and fixed supraglottic mass -A case report. *Korean J Anesthesiol* 2010; 59 Suppl:S26-9
94. Choi JW, Kim JA, Kim HK, Oh MS, Kim DK: Chest anteroposterior diameter affects difficulty of laryngoscopy for non-morbidly obese patients. *J Anesth* 2013; 27:563-8
95. Chung MY, Park B, Seo J, Kim CJ: Successful airway management with combined use of McGrath((R)) MAC video laryngoscope and fiberoptic bronchoscope in a severe obese patient with huge goiter -a case report. *Korean J Anesthesiol* 2018; 71:232-6
96. Ciccozzi A, Angeletti C, Guetti C, Papola R, Angeletti PM, Paladini A, Varrassi G, Marinangeli F: GlideScope and Frova introducer for difficult airway management. *Case Rep Anesthesiol* 2013; 2013:717928
97. Cobas MA, Martin ND, Barkin HB: Two lost airways and one unexpected problem: undiagnosed tracheal stenosis in a morbidly obese patient. *J Clin Anesth* 2016; 35:225-7
98. Cohn AI, McGraw SR, King WH: Awake intubation of the adult trachea using the Bullard laryngoscope. *Can J Anaesth* 1995; 42:246-8
99. Cohn AI, Zornow MH: Awake endotracheal intubation in patients with cervical spine disease: a comparison of the Bullard laryngoscope and the fiberoptic bronchoscope. *Anesth Analg* 1995; 81:1283-6
100. Cook TM, Brooks TS, Van der Westhuizen J, Clarke M: The Proseal LMA is a useful rescue device during failed rapid sequence intubation: two additional cases. *Can J Anaesth* 2005; 52:630-3
101. Cook TM, Silsby J, Simpson TP: Airway rescue in acute upper airway obstruction using a ProSeal Laryngeal mask airway and an Aintree Catheter: A review of the ProSeal laryngeal mask airway in the management of the difficult airway. *Anaesthesia* 2005; 60:1129-1136
102. Coonan TJ, Hope CE, Howes WJ, Holness RO, MacInnis EL: Ankylosis of the temporomandibular joint after temporal craniotomy: a cause of difficult intubation. *Can Anaesth Soc J* 1985; 32:158-60
103. Cooper RM: Use of a new videolaryngoscope (GlideScope) in the management of a difficult airway. *Can J Anaesth* 2003; 50:611-3
104. Cordovani D, Russell T, Wee W, Suen A, Cooper RM: Measurement of forces applied using a Macintosh direct laryngoscope compared with a Glidescope video laryngoscope in patients with predictors of difficult laryngoscopy: A randomised controlled trial. *Eur J Anaesthesiol* 2019; 36:221-6
105. Correll LR, Jin C, Park MS, Webber AM: Urgent complex intraoperative reintubation in a known difficult airway after endotracheal tube damage: A case report. *A A Pract* 2019; 13:4-6
106. Corso RM, Petrini F, Buccioli M, Nanni O, Carretta E, Trolio A, De Nuzzo D, Pigna A, Di Giacinto I, Agnoletti V, Gambale G: Clinical utility of preoperative screening with STOP-Bang questionnaire in elective surgery. *Minerva Anesthesiol* 2014; 80:877-84
107. Črnjar K, Kralik S, Kerovec Sorić I, Bekavac I, Barčot Z, Butković D: Difficult airway management – an old challenge: A case report. *Paediatr Croatica* 2019; 63:38-41
108. Cumpston PH: Fiberoptic intubation under general anaesthesia--a simple method using an endotracheal tube as a conduit. *Anaesth Intens Care* 2009; 37:296-300
109. Curran C: Case report: Oropharyngeal injuries with Glidescope® usage in two obese patients. *Southern African Journal of Anaesthesia and Analgesia* 2015; 22:30–32
110. Dada OF, Faponle AF, Adenekan AT: Description and comparison of the performance of the upper lip bite test, the ratio of height to thyromental distance and other methods of preoperative airway assessment in a Nigerian population – a pilot study. *South Afr J Anaesth Analg* 2019; 25:6-12

111. Dalal PG, Coleman M, Horst M, Rocourt D, Ladda RL, Janicki PK: Case Report: Genetic analysis and anesthetic management of a child with Niemann-Pick disease Type A. *F1000Res* 2015; 4
112. Dalal RJ, Pai H, Pandya S: Difficult airway with HELLP syndrome. *Int J Infertil Fetal Med* 2012; 3:65-7
113. Dangelser G, Dincq AS, Lawson G, Collard E: Case report: Severe laryngeal hemorrhage after withdrawal of a size 5 l-gel in elective surgery. *Acta Anaesthesiol Belg* 2009; 60:255-7
114. Dar S, Khan MS, Iqbal F, Nazeer T, Hussain R: Comparison of upper lip bite test (ULBT) with mallampati classification, regarding assessment of difficult intubation. *Pak J Med Health Sci* 2017; 11:767-9
115. Dargin JM, Emlet LL, Guyette FX: The effect of body mass index on intubation success rates and complications during emergency airway management. *Intern Emerg Med* 2013; 8:75-82
116. Daum RE, Jones DJ: Fiberoptic intubation in Klippel-Feil syndrome. *Anaesthesia* 1988; 43:18-21
117. De Cassai A, Papaccio F, Betteto G, Schiavolin C, Iacobone M, Carron M: Prediction of difficult tracheal intubations in thyroid surgery. Predictive value of neck circumference to thyromental distance ratio. *PLOS ONE* 2019; 14:e0212976
118. Degler SM, Dowling RD, Sucherman DR, Leighton BL: Awake intubation using an intubating laryngeal mask airway in a parturient with spina bifida. *Int J Obstet Anesth* 2005; 14:77-8
119. Dey S, Ninu M, Yunus M, Syiemiong N: Fiberoptic guided retrograde intubation in an anticipated difficult airway: revival of an antiquated technique. *J Clin Diagn Res* 2016; 10:UD06-07
120. Dhanger S, Adinarayanan S, Vinayagam S, Kumar MP: l-gel assisted fiberoptic intubation in a child with Morquio's syndrome. *Saudi J Anaesth* 2015; 9:217-9
121. Dimitriou V, Voyagis GS, Brimacombe JR: Flexible lightwand-guided tracheal intubation with the intubating laryngeal mask Fastrach in adults after unpredicted failed laryngoscope-guided tracheal intubation. *Anesthesiology* 2002; 96:296-9
122. Dimitriou V, Voyagis GS, Douma A: Unexpected resistance during tracheal tube insertion through the intubating laryngeal mask. *Eur J Anaesthesiol* 1999; 16:419-20
123. Dimitriou VK, Zogogiannis ID, Liotiri DG: Awake tracheal intubation using the Airtraq laryngoscope: a case series. *Acta Anaesthesiol Scand* 2009; 53:964-7
124. Dimitriou, V, El Kouny A, Al Harbi M, Wambi F, Tawfeeq N, Tanweer A, Al Atassi A, Geldhof G: Airtraq® 8 optical laryngoscope for tracheal intubation in a patient with an uncommon giant lipoma on the posterior aspect of neck and additional risk factors of anticipated difficult airway: A case report. *Middle East J Anaesthesiol* 2015; 23(3):355-8
125. Divekar VM, Kothari MD, Kamdar BM: Anaesthesia in Turner's syndrome. *Can Anaesth Soc J* 1983; 30:417-8
126. D'Mello J, Pagedar R, Butani M, Kurkal P, Pandey K: Use of the intubating laryngeal mask airway in a case of ankylosing spondylitis for coronary artery bypass grafting. *Eur J Anaesthesiol* 2002; 19:298-302
127. Dohi S, Inomata S, Tanaka M, Ishizawa Y, Matsumiya N: End-tidal carbon dioxide monitoring during awake blind nasotracheal intubation. *J Clin Anesth* 1990; 2:415-9
128. Dohrn N, Sommer T, Bisgaard J, Ronholm E, Larsen JF: Difficult tracheal intubation in obese gastric bypass patients. *Obes Surg* 2016; 26:2640-7
129. Domino KB, Posner KL, Caplan RA, Cheney FW: Airway injury during anesthesia: a closed claims analysis. *Anesthesiology* 1999; 91:1703-11
130. Dong Y, Li G, Wu W, Su R, Shao Y: Lightwand-guided nasotracheal intubation in oromaxillofacial surgery patients with anticipated difficult airways: a comparison with blind nasal intubation. *Int J Oral Maxillofac Surg* 2013; 42:1049-53

131. Driver BE, Prekker ME, Klein LR, Reardon RF, Miner JR, Fagerstrom ET, Cleghorn MR, McGill JW, Cole JB: Effect of use of a bougie vs endotracheal tube and stylet on first-attempt intubation success among patients with difficult airways undergoing emergency intubation: A randomized clinical trial. *JAMA* 2018; 319:2179-89
132. Earl DS, Shinde S, Bullen KE, Carter JA: Novel use of capnography during an awake fiberoptic intubation. *Anaesthesia* 2002; 57:194-5
133. Eiamcharoenwit J, Itthisompaiboon N, Limpawattana P, Suwanpratheap A, Siriussawakul A: The performance of neck circumference and other airway assessment tests for the prediction of difficult intubation in obese parturients undergoing cesarean delivery. *Int J Obstet Anesth* 2017; 31:45-50
134. Eipe N, Fossey S, Kingwell SP: Airway management in cervical spine ankylosing spondylitis: Between a rock and a hard place. *Indian J Anaesth* 2013; 57:592-5
135. El Kholy J, Mohamed NN: A case report of successful awake fiberoptic intubation in a child with severe airway burn. *Egypt J Anaesth* 2013; 29:171-4
136. Ellard L, Brown DH, Wong DT: Extubation of a difficult airway after thyroidectomy: Use of a flexible bronchoscope via the LMA-ClassicTM. *Can J Anaesth* 2012; 59:53-7
137. El-Radaideh K, Al-Maaitah A, Hassoun M, Al-Issa A, Attaieh G: Prediction of difficult intubation depending on two different methods of airway assessment: A prospective study. *Middle East J Anaesthesiol* 2018; 25:181-8
138. El-Radaideh K, Dheeb E, Shbool H, Garaibeh S, Bataineh A, Khraise W, El-Radaideh B: Evaluation of different airway tests to determine difficult intubation in apparently normal adult patients: undergoing surgical procedures. *Pat Safe Surg* 2020; 14
139. El-Tahan MR, Doyle DJ, Khidr AM, Abdulshafi M, Regal MA, Othman MS: Use of the King Vision video laryngoscope to facilitate fiberoptic intubation in critical tracheal stenosis proves superior to the GlideScope(R). *Can J Anaesth* 2014; 61:213-4
140. Etemadi SH, Bahrami A, Farahmand AM, Zamani MM: Sitting nasal intubation with fiberoptic in an elective mandible surgery under general anesthesia. *Anesth Pain Med* 2015; 5:e29299
141. Etezadi F, Ahangari A, Shokri H, Najafi A, Khajavi MR, Daghigh M, Moharari RS: Thyromental height: A new clinical test for prediction of difficult laryngoscopy. *Anesth Analg* 2013; 117:1347-51
142. Ezri T, Medalion B, Weisenberg M, Szmuk P, Warters RD, Charuzi I: Increased body mass index per se is not a predictor of difficult laryngoscopy. *Can J Anaesth* 2003; 50:179-83
143. Ezri T, Waintrob R, Avelansky Y, Izakson A, Dayan K, Shimonov M: Pre-selection of primary intubation technique is associated with a low incidence of difficult intubation in patients with a BMI of 35 kg/m(2) or higher. *Rom J Anaesth Intensive Care* 2018; 25:25-30
144. Ezri T, Warters RD, Szmuk P, Saad-Eddin H, Geva D, Katz J, Hagberg C: The incidence of class "zero" airway and the impact of Mallampati score, age, sex, and body mass index on prediction of laryngoscopy grade. *Anesth Analg* 2001; 93:1073-5
145. Fabregat-Lopez J: Successful pre-emptive emergency management of a compromised airway with a Proseal Laryngeal Mask Airway followed by tracheostomy. *Minerva Anesthesiol* 2012; 78:619-21
146. Falcetta S, Cavallo S, Gabbanelli V, Pelaia P, Sorbello M, Zdravkovic I, Donati A: Evaluation of two neck ultrasound measurements as predictors of difficult direct laryngoscopy: A prospective observational study. *Eur J Anaesthesiol* 2018; 35:605-12
147. Ferrari LR, Bedford RF: General anesthesia prior to treatment of anterior mediastinal masses in pediatric cancer patients. *Anesthesiology* 1990; 72:991-5
148. Ferson DZ, Rosenblatt WH, Johansen MJ, Osborn I, Ovassapian A: Use of the intubating LMA-Fastrach in 254 patients with difficult-to-manage airways. *Anesthesiology* 2001; 95:1175-81



149. Fetterman D, Dubovoy A, Reay M: Unforeseen esophageal misplacement of airway exchange catheter leading to gastric perforation. *Anesthesiology* 2006; 104:1111-2
150. Fitzmaurice BC, Lambert BG: Failed fiberoptic intubation in a child with epidermolysis bullosa, rescued with combined use of the Glidescope(R). *Paediatr Anaesth* 2016; 26:455-6
151. Frawley G, Espenell A, Howe P, Shand J, Heggie A: Anesthetic implications of infants with mandibular hypoplasia treated with mandibular distraction osteogenesis. *Paediatr Anaesth* 2013; 23:342-8
152. Fuchs G, Schwarz G, Baumgartner A, Kaltenbock F, Voit-Augustin H, Planinz W: Fiberoptic intubation in 327 neurosurgical patients with lesions of the cervical spine. *J Neurosurg Anesthesiol* 1999; 11:11-6
153. Fukutome T, Amaha K, Nakazawa K, Kawamura T, Noguchi H: Tracheal intubation through the intubating laryngeal mask airway (LMA-Fastrach) in patients with difficult airways. *Anaesth Intens Care* 1998; 26:387-91
154. Fulkerson JS, Moore HM, Lowe RF, Anderson TS, Lucas LL, Reed JW: Airway sonography fails to detect difficult laryngoscopy in an adult Veteran surgical population. *Trends Anaesth Crit Care* 2019; 29:26-34
155. Furutani K, Kodera Y, Hiruma M, Ishii H, Baba H: Difficult tracheal intubation in a patient with maternal uniparental disomy 14. *JA Clin Rep* 2016; 2:25
156. Gaitini L, Fradis M, Croitoru M, Somri M, Vaida S: The intubating laryngeal mask. *Otolaryngol Head Neck Surg* 2002; 126:87-8
157. Galderisi A, De Bernardo G, Lorenzon E, Trevisanuto D: i-gel: a new supraglottic device for effective resuscitation of a very low birthweight infant with Cornelia de Lange syndrome. *BMJ Case Reports* 2015; 25:25
158. Gallagher DM, Hyler RL, Epker BN: Hemifacial microsomia: an anesthetic airway problem. *Oral Surg Oral Med Oral Pathol* 1980; 49:2-4
159. Gandhe RU, Bhave CP, Kakde AS, Sathe KA: Unanticipated difficulty in an anticipated difficult airway in the neurointervention suite: A case report. *J Neuroanaesth Crit Care* 2018; 5:190-2
160. Gaszynska E, Wiczorek A, Gaszynski T: Awake endotracheal intubation in patients with severely restricted mouth opening- alternative devices to fiberscope: Series of cases and literature review. *Cent Eur J Med* 2014; 9:768-72
161. Gaszynski T, Gaszynska E, Szewczyk T: Dexmedetomidine for awake intubation and an opioid-free general anesthesia in a superobese patient with suspected difficult intubation. *Drug Des Devel Ther* 2014; 8:909-12
162. Gaszynski T, Gaszynska E: The Clarus Video System stylet for awake intubation in a very difficult urgent intubation. *Anesthesiol Intensive Ther* 2013; 45:153-4
163. Gaszyński T: The use of the C-MAC videolaryngoscope for awake intubation in patients with a predicted extremely difficult airway: case series. *Ther Clin Risk Manag* 2018; 14:539-542
164. Gautam PL, Kaur M, Singh RJ, Gupta S: Large mediastinal tumor in a neonate: an anesthetic challenge. *J Anesth* 2012; 26:124-7
165. Gemma M, Buratti L, Di Santo D, Calvi MR, Ravizza A, Bondi S, Bussi M, Beretta L: Pre-operative transnasal endoscopy as a predictor of difficult airway: A prospective cohort study. *Eur J Anaesthesiol* 2020; 37:98-104
166. Ghabach MB, Roupheal MAA, Roumoulian CE, Helou MR: Airway management in a patient with Le Fort III fracture. *Saudi J Anaesth* 2014; 8:128-30
167. Ghaly RF, Candido KD, Sauer R, Knezevic NN: Anesthetic management during Cesarean section in a woman with residual Arnold-Chiari malformation Type I, cervical kyphosis, and syringomyelia. *Surg Neurol Int* 2012; 3:26
168. Ghanem MT, Ahmed FI: GlideScope versus McCoy laryngoscope: Intubation profile for cervically unstable patients in critical care setting. *Egypt J Anaesth* 2017; 33:103-6

169. Gist RS, Miller DW, Warren T: A difficult airway in a patient with nephrogenic sclerosing fibrosis. *Anesth Analg* 2010; 110:555-7
170. Godley M, Reddy AR: Use of LMA for awake intubation for caesarean section. *Can J Anaesth* 1996; 43:299-302
171. Goni-Zaballa M, Perez-Ferrer A, Charco-Mora P: Difficult airway in a pediatric patient with Klippel-Feil syndrome and an unexpected lingual tonsil. *Minerva Anesthesiol* 2012; 78:254-7
172. González-Giraldo D, Largo-Pineda CE, Zamudio-Burbano MA: Successful rescue with videolaryngoscopy after failed fibroscopy in anticipated difficult airway: Case series. *Colomb J Anesthesiol* 2020; 48:96-9
173. Gorbach MS: Management of the challenging airway with the Bullard laryngoscope. *J Clin Anesth* 1991; 3:473-7
174. Graciano AL, Tamburro R, Thompson AE, Fiadjoe J, Nadkarni VM, Nishisaki A: Incidence and associated factors of difficult tracheal intubations in pediatric ICUs: a report from National Emergency Airway Registry for Children: NEAR4KIDS. *Intensive Care Med* 2014; 40:1659-69
175. Grass B, Simma L, Reinehr M, Zimmermann U, Gysin C, Henze G, Cannizzaro V: Two case reports of unexpected tracheal agenesis in the neonate: 3 C's beyond algorithms for difficult airway management. *BMC Pediatr* 2017; 17:49
176. Greif R, Kleine-Brueggene M, Theiler L: Awake tracheal intubation using the Sensascope in 13 patients with an anticipated difficult airway. *Anaesthesia* 2010; 65:525-8
177. Grimes D, Macleod I, Taylor T, O'Connor M, Sidebottom A: Computed tomography as an aid to planning intubation in the difficult airway. *Br J Oral Maxillofac Surg* 2016; 54:80-2
178. Guardo LA, Gala CC, Calvo AR, Aguirre AM, Arango NA, Fernandez EG, Chiroso CR, Poveda DS, Alvarez AD: Known difficult airway in a patient with pheochromocytoma: a case report. *J Clin Anesth* 2016; 35:411-414
179. Guerrero-Dominguez R, Lopez-Herrera-Rodriguez D, Benitez-Linero I, Ontanilla A: Anesthetic management for surgery of esophagus atresia in a newborn with Goldenhar's syndrome. *Brazilian J Anesthesiol* 2015; 65:298-301
180. Guo Y, Feng Y, Liang H, Zhang R, Cai X, Pan X: Role of flexible fiberoptic laryngoscopy in predicting difficult intubation. *Minerva Anesthesiol* 2018; 84:337-45
181. Gupta A, Gupta N: Anterior laryngeal web leading to unanticipated difficult tracheal intubation in a neonate diagnosed and managed successfully with CMAC video laryngoscope: A case report. *A A Pract* 2018; 10:28-30
182. Gupta N, Pandia MP, Prabhakar H, Chauhan M: Video laryngoscopy added fiberoptic intubation in a patient with difficult airway. *J Anaesthesiol Clin Pharmacol* 2013; 29:283-4
183. Gupta N, Rath GP, Bala R, Reddy BK, Chaturvedi A: Anesthetic management in children with Hurler's syndrome undergoing emergency ventriculoperitoneal shunt surgery. *Saudi J Anaesth* 2012; 6:178-80
184. Gupta N, Rath GP, Prabhakar H: Clinical evaluation of C-MAC videolaryngoscope with or without use of stylet for endotracheal intubation in patients with cervical spine immobilization. *J Anesth* 2013; 27:663-70
185. Gupta S, Pareek S, Dulara SC: Comparison of two methods for predicting difficult intubation in obstetric patients. *Middle East J Anaesthesiol* 2003; 17:275-85
186. Gurusurthy T, Shailaja S, Kishan S, Stephen M: Management of an anticipated difficult airway in Hurler's syndrome. *J Anaesthesiol Clin Pharmacol* 2014; 30:558-61
187. Hajjar WM, Alsubaie N, Nouh TA, Al-Nassar SA: Is it safe to use frova airway intubating device during tracheal intubation in difficult airway patient with multiple and chest trauma? *Saudi J Anaesth* 2016; 10:477-9
188. Han TH, Teissler H, Han RJ, Gaines JD, Nguyen TQ: Managing difficult airway in patients with post-burn mentosternal and circumoral scar contractures. *Int J Burns Trauma* 2012; 2:80-5

189. Han Y, Fang J, Zhang H, Xu M, Guo X: Anterior neck soft tissue thickness for airway evaluation measured by MRI in patients with cervical spondylosis: Prospective cohort study. *BMJ Open* 2019; 9
190. Han YZ, Tian Y, Xu M, Ni C, Li M, Wang J, Guo XY: Neck circumference to inter-incisor gap ratio: a new predictor of difficult laryngoscopy in cervical spondylosis patients. *BMC Anesthesiol* 2017; 17:55
191. Han YZ, Tian Y, Zhang H, Zhao YQ, Xu M, Guo XY: Radiologic indicators for prediction of difficult laryngoscopy in patients with cervical spondylosis. *Acta Anaesthesiol Scand* 2018; 62:474-82
192. Hanna SF, Mikat-Stevens M, Loo J, Uppal R, Jellish WS, Adams M: Awake tracheal intubation in anticipated difficult airways: LMA Fastrach vs flexible bronchoscope: A pilot study. *J Clin Anesth* 2017; 37:31-7
193. Hansda U, Agarwal J, Patra C, Ganjoo P: Extradural hematoma surgery in a child with Hutchinson-Gilford progeria syndrome: Perioperative concerns. *J Pediatr Neurosci* 2013; 8:165-7
194. Hariharan U, Shah SB, Naithani BK: Difficult intubation due to outgrowth between the epiglottic fold and the vocal cords: C-MAC™ to our rescue! *Sri Lankan J Anaesthesiol* 2015; 23
195. Harjai M, Bhaskar P, Saxena S, Rastogi S, Singh PK: Is RHTMD a new predictor for assessment of difficult intubation in non-obese patients? *J Evol Med Dent Sci* 2018; 7:4957-60
196. Hashim KV, Thomas M: Sensitivity of palm print sign in prediction of difficult laryngoscopy in diabetes: A comparison with other airway indices. *Indian J Anaesth* 2014; 58:298-302
197. Hazarika H, Saxena A, Meshram P, Kumar Bhargava A: A randomized controlled trial comparing C Mac D Blade and Macintosh laryngoscope for nasotracheal intubation in patients undergoing surgeries for head and neck cancer. *Saudi J Anaesth* 2018; 12:35-41
198. Heinrich S, Birkholz T, Ihmsen H, Irouschek A, Ackermann A, Schmidt J: Incidence and predictors of difficult laryngoscopy in 11,219 pediatric anesthesia procedures. *Paediatr Anaesth* 2012; 22:729-36
199. Heinrich S, Birkholz T, Irouschek A, Ackermann A, Schmidt J: Incidences and predictors of difficult laryngoscopy in adult patients undergoing general anesthesia: A single-center analysis of 102,305 cases. *J Anesth* 2013; 27:815-21
200. Hill CM: Death following dental clearance in a patient suffering from ankylosing spondylitis - a case report with discussion on management of such problems. *Br J Oral Surg* 1980; 18:73-6
201. Hilton G, Mihm F, Butwick A: Anesthetic management of a parturient with VACTERL association undergoing Cesarean delivery. *Can J Anaesth* 2013; 60:570-6
202. Hiraakawa M, Nishihara T, Nakanishi K, Kitamura S, Fujii S, Ikemune K, Dote K, Takasaki Y, Yorozuya T: Perioperative management of a patient with Coffin-Lowry syndrome complicated by severe obesity: A case report and literature review. *Medicine* 2017; 96:e9026
203. Hirmanpour A, Safavi M, Honarmand A, Jabalameli M, Banisadr G: The predictive value of the ratio of neck circumference to thyromental distance in comparison with four predictive tests for difficult laryngoscopy in obstetric patients scheduled for caesarean delivery. *Adv Biomed Res* 2014; 3:200
204. Hodgson RE, Pillay TK: Awake percutaneous tracheostomy as an alternative to open emergency tracheostomy in a threatened airway. *South Afr J Anaesth Analg* 2017; 23:23-8
205. Holzman RS, Nargoizian CD, Florence FB: Lightwand intubation in children with abnormal upper airways. *Anesthesiology* 1988; 69:784-7
206. Honarmand A, Safavi M, Ansari N: A comparison of between hyomental distance ratios, ratio of height to thyromental, modified Mallampati classification test and upper lip bite

- test in predicting difficult laryngoscopy of patients undergoing general anesthesia. *Adv Biomed Res* 2014; 3:166
207. Honarmand A, Safavi M, Yaraghi A, Attari M, Khazaei M, Zamani M: Comparison of five methods in predicting difficult laryngoscopy: Neck circumference, neck circumference to thyromental distance ratio, the ratio of height to thyromental distance, upper lip bite test and Mallampati test. *Adv Biomed Res* 2015; 4:122
  208. Hsin ST, Chen CH, Juan CH, Tseng KW, Oh CH, Tsou MY, Tsai SK: A modified method for intubation of a patient with ankylosing spondylitis using intubating laryngeal mask airway (LMA-Fastrach)--a case report. *Acta Anaesthesiol Sin* 2001; 39:179-82
  209. Huang L, Wang J, Chen S, Fang X: Study and reflection on anesthesia for tracheobronchopathia osteochondroplastica. *J Int Med Res* 2020; 48
  210. Huang R-C, Hsu C-H, Chuang Y-S, Chan W-H, Wu Z-F, Cherng C-H, Kuo C-Y: Successful nasotracheal intubation in a patient with distorted airway anatomy by combined use of flexible fiberoptic bronchoscope and trachway. *J Med Sci* 2014; 34:95-7
  211. Hung OR, Pytka S, Morris I, Murphy M, Stewart RD: Lightwand intubation: II--Clinical trial of a new lightwand for tracheal intubation in patients with difficult airways. *Can J Anaesth* 1995; 42:826-30
  212. Ikram M, Mahboob S: Anesthetic challenges in a large multinodular thyroidectomy at a peripheral hospital. *Anaesth Pain Intens Care* 2019; 23:311-3
  213. Indiveri L, Mohamed AN, Milner A: Branchio-otic syndrome: An opportunity to reassess the paediatric anaesthetists' approach to the difficult syndromic airway. *South Afr J Anaesth Analg* 2019; 25:27-30
  214. Iseli TA, Iseli CE, Golden JB, Jones VL, Boudreaux AM, Boyce JR, Weeks DM, Carroll WR: Outcomes of intubation in difficult airways due to head and neck pathology. *Ear Nose Throat J* 2012; 91:E1-5
  215. Jagannathan N, Sequera-Ramos L, Sohn L, Wallis B, Shertzer A, Schaldenbrand K: Elective use of supraglottic airway devices for primary airway management in children with difficult airways. *Br J Anaesth* 2014; 112:742-8
  216. Jain A, Naithani M: Infant with unanticipated difficult airway - Trachlight to the rescue. *J Anaesthesiol Clin Pharmacol* 2012; 28:361-3
  217. Jain M, Gupta A, Garg M, Rastogi B, Chauhan H: Innovative lighted stylet--succeeds where conventional lighted stylet fails. *Middle East J Anaesthesiol* 2009; 20:447-50
  218. Jain N, Das S, Kanchi M: Thyromental height test for prediction of difficult laryngoscopy in patients undergoing coronary artery bypass graft surgical procedure. *Ann Card Anaesth* 2017; 20:207-11
  219. Jamuna T, Rao MI, Naveen: Comparative study between MMT classification and ULBT for predicting difficulty during endotracheal intubation. *J Evol Med Dent Sci* 2018; 7:1428-33
  220. Jarraya A, Choura D, Mejdoub Y, Kammoun M, Grati F, Kolsi K: New predictors of difficult intubation in obstetric patients: A prospective observational study. *Trends Anaesth Crit Care* 2019; 24:22-5
  221. Jeong H, Chae M, Seo H, Yi JW, Kang JM, Lee BJ: Face-to-face intubation using a lightwand in a patient with severe thoracolumbar kyphosis: a case report. *BMC Anesthesiol* 2018; 18:92
  222. Joo H, Rose K: Fastrach--a new intubating laryngeal mask airway: successful use in patients with difficult airways. *Can J Anaesth* 1998; 45:253-6
  223. Joo HS, Kapoor S, Rose DK, Naik VN: The intubating laryngeal mask airway after induction of general anesthesia versus awake fiberoptic intubation in patients with difficult airways. *Anesth Analg* 2001; 92:1342-6
  224. Jungbauer A, Schumann M, Brunkhorst V, Borgers A, Groeben H: Expected difficult tracheal intubation: a prospective comparison of direct laryngoscopy and video laryngoscopy in 200 patients. *Br J Anaesth* 2009; 102:546-50

225. Juvin P, Lavaut E, Dupont H, Lefevre P, Demetriou M, Dumoulin JL, Desmonts JM: Difficult tracheal intubation is more common in obese than in lean patients. *Anesth Analg* 2003; 97:595-600
226. Kakizaki R, Bunya N, Uemura S, Narimatsu E: Successful difficult airway management with emergent venovenous extracorporeal membrane oxygenation in a patient with severe tracheal deformity: a case report. *Acute Med Surg* 2020; 7
227. Kalra S: Unanticipated difficult intubation in a child with Beals-Hecht syndrome presenting for emergency surgery. *Sri Lankan J Anaesthesiol* 2015; 23:27-8
228. Kamal G, Gupta A, Batla S, Gupta N: Anaesthetic management of a child with stone man syndrome: Look before you leap!. *Indian J Anaesth* 2017; 61:266-268
229. Kanase NV, Gandhi S, Singh AK: Predicting difficult intubation through upper lip bite test and modified mallampati classification. *J Cardiovasc Dis Res* 2020; 11:36-9
230. Kanemaru H, Tsurumaki T, Kurata S, Tanaka Y, Yoshikawa H, Sato Y, Kodama Y, Suda A, Yamada Y, Seo K: Endotracheal intubation complicated by a palatal tooth in a patient with Treacher Collins Syndrome. *Anesth Prog* 2019; 66:42-3
231. Kaniyil S, Anandan K, Thomas S: Ratio of height to thyromental distance as a predictor of difficult laryngoscopy: A prospective observational study. *J Anaesthesiol Clin Pharmacol* 2018; 34:485-9
232. Kannan S, Chestnutt N, McBride G: Intubating LMA guided awake fiberoptic intubation in severe maxillo-facial injury. *Can J Anaesth* 2000; 47:989-91
233. Karami A, Malak A, Vatankhan P, Saravi ZF: Predicting factors of difficult intubation in obese patients undergoing bariatric surgeries. *Res J Med Sci* 2016; 10:565-7
234. Kariya N, Nishi S, Minami W, Funao T, Mori M, Nishikawa K, Asada A: Airway problems related to laryngeal mask airway use associated with an undiagnosed epiglottic cyst. *Anaesth Intensive Care* 2004; 32:268-70
235. Karsli C, Armstrong J, John J: A comparison between the GlideScope video laryngoscope and direct laryngoscope in paediatric patients with difficult airways - a pilot study. *Anaesthesia* 2010; 65:353-357
236. Katsiampoura AD, Killoran PV, Corso RM, Cai C, Hagberg CA, Cattano D: Laryngeal mask placement in a teaching institution: Analysis of difficult placements. *F1000Res* 2015; 4
237. Kaufmann J, Laschat M, Engelhardt T, Hellmich M, Wappler F: Tracheal intubation with the Bonfils fiberscope in the difficult pediatric airway: A comparison with fiberoptic intubation. *Paediatr Anaesth* 2015; 25:372-8
238. Kaur J, Swami AC, Kumar A, Lata S: Anesthetic management of a child with Hunter's syndrome. *J Anaesthesiol Clin Pharmacol* 2012; 28:255-7
239. Kawai T, Shimozato K, Ochiai S: Elongated styloid process as a cause of difficult intubation. *J Oral Maxillofac Surg* 1990; 48:1225-8
240. Keenan MA, Stiles CM, Kaufman RL: Acquired laryngeal deviation associated with cervical spine disease in erosive polyarticular arthritis. Use of the fiberoptic bronchoscope in rheumatoid disease. *Anesthesiology* 1983; 58:441-9
241. Kelly FE, Gatward JJ, Howes BW, Gould TH: Kinking of intubating laryngeal mask tracheal tubes following fiberoptic intubation. *Anaesthesia* 2010; 65:653
242. Kezo A, Patel RD, Mathkar S, Butada S: Use of a Macintosh blade in extrahepatic portal vein obstruction with difficult intubation: two case reports. *J Med Case Rep* 2016; 10:245
243. Khan MU: Endotracheal intubation in patients with unstable cervical spine using LMA-Fastrach and gum elastic bogie. *J Coll Physicians Surg Pak* 2014; 24:4-7
244. Khan ZH, Arbabi S: Diagnostic value of the upper lip bite test in predicting difficulty in intubation with head and neck landmarks obtained from lateral neck X-ray. *Indian J Anaesth* 2013; 57:381-6

245. Khan ZH, Eskandari S, Yekaninejad MS: A comparison of the Mallampati test in supine and upright positions with and without phonation in predicting difficult laryngoscopy and intubation: A prospective study. *J Anaesthesiol Clin Pharmacol* 2015; 31:207-11
246. Khatiwada S, Bhattarai B, Pokharel K, Acharya R: Prediction of difficult airway among patients requiring endotracheal intubation in a tertiary care hospital in eastern Nepal. *J Nepal Med Assoc* 2017; 56:314-8
247. Kidani DC, Shah NK: The use of a laryngeal mask airway after a prolonged suspension laryngoscopy to preserve a vocal cord fat graft. *Anesth Analg* 2007; 105:1753-4
248. Kihara S, Watanabe S, Brimacombe J, Taguchi N, Yaguchi Y, Yamasaki Y: Segmental cervical spine movement with the intubating laryngeal mask during manual in-line stabilization in patients with cervical pathology undergoing cervical spine surgery. *Anesth Analg* 2000; 91:195-200
249. Kilicaslan A, Topal A, Tavlan A, Erol A, Otelcioglu S: Effectiveness of the C-MAC video laryngoscope in the management of unexpected failed intubations. *Braz J Anesthesiol* 2014; 64:62-5
250. Kim JC, Ki Y, Kim J, Ahn SW: Ethnic considerations in the upper lip bite test: The reliability and validity of the upper lip bite test in predicting difficult laryngoscopy in Koreans. *BMC Anesthesiol* 2019; 19
251. Kim JS, Park SY, Min SK, Kim JH, Lee SY, Moon BK: Awake nasotracheal intubation using fiberoptic bronchoscope in a pediatric patient with Freeman-Sheldon syndrome. *Paediatr Anaesth* 2005; 15:790-2
252. Kim SM, Kim HJ: Successful advancement of endotracheal tube with combined fiberoptic bronchoscopy and videolaryngoscopy in a patient with a huge goiter. *SAGE Open Med Case Rep* 2020; 8
253. Kim Y, Kim JE, Jeong DH, Lee J: Combined use of a McGrath(R) MAC video laryngoscope and Frova Intubating Introducer in a patient with Pierre Robin syndrome: A case report. *Korean J Anesthesiol* 2014; 66:310-3
254. Kim YL, Seo DM, Shim KS, Kim EJ, Lee JH, Lee SG, Ban JS: Successful tracheal intubation using fiberoptic bronchoscope via an I-gel supraglottic airway in a pediatric patient with Goldenhar syndrome -A case report. *Korean J Anesthesiol* 2013; 65:61-5
255. Kitahata LM: Airway difficulties associated with anaesthesia in acromegaly. Three case reports. *Br J Anaesth* 1971; 43:1187-90
256. Kleeman PP, Jantzen JP, Bonfils P: The ultra-thin bronchoscope in management of the difficult paediatric airway. *Can J Anaesth* 1987; 34:606-8
257. Kleine-Bruegeney M, Theiler L, Urwyler N, Vogt A, Greif R: Randomized trial comparing the i-gel™ and Magill tracheal tube with the single-use ILMA™ and ILMA™ tracheal tube for fiberoptic-guided intubation in anaesthetized patients with a predicted difficult airway. *Br J Anaesth* 2011; 107:251-257
258. Kolarkar P, Badwaik G, Watve A, Abhishek K, Bhangale N, Bhalerao A, Gupta GJ, Giri A: Upper lip bite test: a novel test of predicting difficulty in intubation. *J Evol Med Dent Sci* 2015; 4:4149-56
259. Kondo I, Kobayashi H, Suga Y, Suzuki A, Kiyama S, Uezono S: Effect of availability of video laryngoscopy on the use of fiberoptic intubation in school-aged children with microtia. *Paediatr Anaesth* 2017; 27:1115-1119
260. Konwar C, Baruah ND, Saikia P, Chakrabartty A: A prospective study of the usefulness of upper lip bite test in combination with sternomental distance, thyro-mental distance and inter-incisor distance as predictor of ease of laryngoscopy. *J Evol Med Dent Sci* 2015; 4:16286-9
261. Kothandan H, Ho VK, Chan YM, Wong T: Difficult intubation in a patient with vallecular cyst. *Singapore Med J* 2013; 54:e62-5

262. Koundal V, Rana S, Thakur R, Chauhan V, Ekke S, Kumar M: The usefulness of point of care ultrasound (POCUS) in preanaesthetic airway assessment. *Indian J Anaesth* 2019; 63:1022-8
263. Kovac AL: Use of the Augustine stylet anticipating difficult tracheal intubation in Treacher-Collins syndrome. *J Clin Anesth* 1992; 4:409-12
264. Kovacs G, Law AJ, Petrie D: Awake fiberoptic intubation using an optical stylet in an anticipated difficult airway. *Ann Emerg Med* 2007; 49:81-3
265. Krafft P, Fitzgerald R, Pernerstorfer T, Kapral S, Weinstabl C: A new device for blind oral intubation in routine and difficult airway management. *Eur J Anaesthesiol* 1994; 11:207-12
266. Krishna HM, Shetty N, Vinodhadevi V: Airtraq in difficult/failed intubation. *Sri Lankan J Anaesthesiol* 2013; 21:32-35
267. Krishna R, Shenoy TV, Goneppanavar U: Airway management in an infant with congenital trismus: The role of retrograde intubation. *South Afr J Anaesth Analg* 2012; 18:267-9
268. Krishnan PL, Thiessen BH: Use of the Bonfils intubating fibroscope in a baby with a severely compromised airway. *Paediatr Anaesth* 2013; 23:670-2
269. Kukreja V, Khurana S, Kohli G, Sadhu R: Treacher Collins syndrome: Case report of a patient with a difficult airway. *J Clin Diagn Res* 2012; 6:1089-91
270. Kulka PJ, Tryba M, Zenz M: Difficult airway management in a patient with severe aortic stenosis, coronary artery disease, and heart failure. *J Clin Anesth* 2002; 14:150-3
271. Kumar D, Gombar S, Ahuja V, Malhotra A, Gupta S: GlideScope versus D-blade for tracheal intubation in cervical spine patients: A randomised controlled trial. *Indian J Anaesth* 2019; 63:544-550
272. Kurnutala LN, Sandhu G, Bergese SD: Fiberoptic nasopharyngoscopy for evaluating a potentially difficult airway in a patient with elevated intracranial pressure. *J Clin Anesth* 2016; 34:336-8
273. Kwon YS, Lee CA, Park S, Ha SO, Sim YS, Baek MS: Incidence and outcomes of cricothyrotomy in the "cannot intubate, cannot oxygenate" situation. *Medicine* 2019; 98:e17713
274. Lai HY, Chen IH, Chen A, Hwang FY, Lee Y: The use of the GlideScope for tracheal intubation in patients with ankylosing spondylitis. *Br J Anaesth* 2006; 97:419-22
275. Lakhe G, Poudel H, Adhikari KM: Assessment of airway parameters for predicting difficult laryngoscopy and intubation in a tertiary center in western Nepal. *J Nepal Health Res Counc* 2020; 17:516-20
276. Lange M, Frommer M, Redel A, Trautner H, Hampel J, Kranke P, Kehl F, Scholtz LU, Roewer N: Comparison of the Glidescope and Airtraq optical laryngoscopes in patients undergoing direct microlaryngoscopy. *Anaesthesia* 2009; 64:323-8
277. Langeron O, Masso E, Huraux C, Guggiari M, Bianchi A, Coriat P, Riou B: Prediction of difficult mask ventilation. *Anesthesiology* 2000; 92:1229-36
278. Langeron O, Semjen F, Bourgain JL, Marsac A, Cros AM: Comparison of the intubating laryngeal mask airway with the fiberoptic intubation in anticipated difficult airway management. *Anesthesiology* 2001; 94:968-72
279. Larson SM, Parks DH: Managing the difficult airway in patients with burns of the head and neck. *J Burn Care Rehabil* 1988; 9:55-6
280. Łasińska-Kowara M, Karwacki Z, Łapiński M: Complications after using the Airtraq laryngoscope for a predicted difficult intubation. *Anaesth Intensiv Ther* 2013; 45:35-37
281. Leary JA: Mechanical failure of the McCoy laryngoscope during a difficult intubation. *Anaesthesia* 2001; 56:88-9
282. Lee HC, Andree RA: Cervical spondylosis and difficult intubation. *Anesth Analg* 1979; 58:434-5
283. Lee HC, Kim MK, Kim YH, Park HP: Radiographic predictors of difficult laryngoscopy in acromegaly patients. *J Neurosurg Anesthesiol* 2019; 31:50-6

284. Lee JH, Jung HC, Shim JH, Lee C: Comparison of the rate of successful endotracheal intubation between the "sniffing" and "ramped" positions in patients with an expected difficult intubation: a prospective randomized study. *Korean J Anesthesiol* 2015; 68:116-21
285. Lee JJ, Lim BG, Lee MK, Kong MH, Kim KJ, Lee JY: Fiberoptic intubation through a laryngeal mask airway as a management of difficult airway due to the fusion of the entire cervical spine - A report of two cases. *Korean J Anesthesiol* 2012; 62:272-6
286. Lee KH, Kang ES, Jung JW, Park JH, Choi YG: Use of the i-gel supraglottic airway device in a patient with subglottic stenosis -a case report. *Korean J Anesthesiol* 2013; 65:254-6
287. Lee SM, Lim H: McGrath® videolaryngoscopy in an awake patient with a huge dangling vocal papilloma: a case report. *J Int Med Res* 2019; 47:3416-20
288. Lenhardt R, Burkhart MT, Brock GN, Kanchi-Kandadai S, Sharma R, Akca O: Is video laryngoscope-assisted flexible tracheoscope intubation feasible for patients with predicted difficult airway? A prospective, randomized clinical trial. *Anesth Analg* 2014; 118:1259-65
289. Leoni A, Arlati S, Ghisi D, Verweij M, Lugani D, Ghisi P, Cappelleri G, Cedrati V, El Tantawi Ali Alsheraei A, Pocar M, Ceriani V, Aldegheri G: Difficult mask ventilation in obese patients: Analysis of predictive factors. *Minerva Anesthesiol* 2014; 80:149-57
290. Levin R, Kissoon N, Froese N: Fiberoptic and videoscopic indirect intubation techniques for intubation in children. *Pediatr Emerg Care* 2009; 25:473-9
291. Levitt MW, Collison JM: Difficult endotracheal intubation in a patient with pseudoxanthoma elasticum. *Anaesth Intensive Care* 1982; 10:62-4
292. Li Q, Xie P, Zha B, Wu Z, Wei H: Supraglottic jet oxygenation and ventilation saved a patient with 'cannot intubate and cannot ventilate' emergency difficult airway. *J Anesth* 2017; 31:144-7
293. Liang H, Hou Y, Wei H, Feng Y: Supraglottic jet oxygenation and ventilation assisted fiberoptic intubation in a paralyzed patient with morbid obesity and obstructive sleep apnea: A case report. *BMC Anesthesiol* 2019; 19
294. Liaskou C, Vouzounerakis E, Moirasgenti M, Trikoupi A, Staikou C: Anatomic features of the neck as predictive markers of difficult direct laryngoscopy in men and women: A prospective study. *Indian J Anaesth* 2014; 58:176-82
295. Liew G, Leong XF, Wong T: Awake tracheal intubation in a patient with a supraglottic mass with the Bonfils fibroscope after failed attempts with a flexible fibroscope. *Singapore Med J* 2015; 56:e139-41
296. Liew GHC, Wong TGL, Lu A, Kothandan H: Combined use of the glidescope and flexible fibroscope as a rescue technique in a difficult airway. *Proceed Singapore Health* 2015; 24:117-20
297. Lillie EM, Harding L, Thomas M: A new twist in the pediatric difficult airway. *Paediatr Anaesth* 2015; 25:428-30
298. Lim WY, Wong P: Awake supraglottic airway guided flexible bronchoscopic intubation in patients with anticipated difficult airways: A case series and narrative review. *Korean J Anesthesiol* 2019; 72:548-57
299. Lin YT, Lee YS, Jeng MJ, Chen WY, Tsao PC, Chan IC, Soong WJ: Flexible bronchoscopic findings and the relationship to repeated extubation failure in critical children. *J Chin Med Assoc* 2018; 81:804-10
300. Liu L, Yue H, Li J: Comparison of three tracheal intubation techniques in thyroid tumor patients with a difficult airway: A randomized controlled trial. *Med Princ Pract* 2014; 23:448-52
301. Liu Z, Zhao L, Jia Q, Liang S, Yang X: A comparison of disposcope endoscope and fiberoptic bronchoscope during awake orotracheal intubation in obese patients with anticipated difficult airway: A prospective, randomized and controlled clinical trial. *Int J Clin Exp Med* 2018; 11:9654-9661



302. Loftus PA, Ow TJ, Siegel B, Tassler AB, Smith RV, Schiff BA: Risk factors for perioperative airway difficulty and evaluation of intubation approaches among patients with benign goiter. *Ann Otol Rhinol Laryngol* 2014; 123:279-85
303. Ludena JA, Bellas JJA, Alvarez-Rementería R, Muñoz LE: Fiberoptic-guided intubation after awake insertion of the I-gel™ supraglottic device in a patient with predicted difficult airway. *J Anaesthesiol Clin Pharmacol* 2017; 33:560-1
304. Lyons G: Failed intubation. Six years' experience in a teaching maternity unit. *Anaesthesia* 1985; 40:759-62
305. Mackie AM, Watson CB: Anaesthesia and mediastinal masses. A case report and review of the literature. *Anaesthesia* 1984; 39:899-903
306. Madan K, Shrestha P, Garg R, Hadda V, Mohan A, Guleria R: Bronchoscopic management of critical central airway obstruction by thyroid cancer: Combination airway stenting using tracheal and inverted-Y carinal self-expanding metallic stents. *Lung India* 2017; 34:202-5
307. Maddali MM, Al-Zaabi HAM, Al-Aamri ISS, Arora NR, Panchatcharam SM: Preoperative predictors of poor laryngoscope views in pediatric population undergoing cardiac catheterization. *Ann Card Anaesth* 2018; 21:376-81
308. Magalhaes E, Oliveira Marques F, Sousa Goveia C, Araujo Ladeira LC, Lagares J: Use of simple clinical predictors on preoperative diagnosis of difficult endotracheal intubation in obese patients. *Braz J Anesthesiol* 2013; 63:262-6
309. Mahmoodpoor A, Soleimanpour H, Golzari SE, Nejabatian A, Pournak T, Amani M, Hajmohammadi S, Hosseinzadeh H, Esfanjani RM: Determination of the diagnostic value of the modified mallampati score, upper lip bite test and facial angle in predicting difficult intubation: A prospective descriptive study. *J Clin Anesth* 2017; 37:99-102
310. Mahmoodpoor A, Soleimanpour H, Shahsavari Nia KS, Rahimi Panahi J, Afhami M, Samad EJ, Majani K: Sensitivity of palm print, modified mallampati score and 3-3-2 rule in prediction of difficult intubation. *Int J Prev Med* 2013; 4:1063-9
311. Mahrous RSS, Ahmed AMM: The Shikani Optical Stylet as an alternative to awake fiberoptic intubation in patients at risk of secondary cervical spine injury: A randomized controlled trial. *J Neurosurg Anesthesiol* 2018; 30:354-8
312. Maktabi MA, Hoffman H, Funk G, From RP: Laryngeal trauma during awake fiberoptic intubation. *Anesth Analg* 2002; 95:1112-4
313. Malik MA, Subramaniam R, Maharaj CH, Harte BH, Laffey JG: Randomized controlled trial of the Pentax AWS, Glidescope, and Macintosh laryngoscopes in predicted difficult intubation. *Br J Anaesth* 2009; 103:761-8
314. Mallampati SR, Gatt SP, Gugino LD, Desai SP, Waraksa B, Freiburger D, Liu PL: A clinical sign to predict difficult tracheal intubation: a prospective study. *Can Anaesth Soc J* 1985; 32:429-34
315. Malpas G, Hung O, Gilchrist A, Wong C, Kent B, Hirsch GM, Hart RD: The use of extracorporeal membrane oxygenation in the anticipated difficult airway: A case report and systematic review. *Can J Anaesth* 2018; 65:685-97
316. Manabe Y, Iwamoto S, Miyawaki H, Seo K, Sugiyama K: Mallampati classification without tongue protrusion can predict difficult tracheal intubation more accurately than the traditional Mallampati classification. *Oral Sci Int* 2014; 11:52-5
317. Mansano AM, Modolo NS, Silva LM, Ganem EM, Braz LG, Knabe Ade C, Freitas FM: Bedside tests to predict laryngoscopic difficulty in pediatric patients. *Int J Pediatr Otorhinolaryngol* 2016; 83:63-8
318. Marcotegui Barber I, Bilbao Ares A, Azcona Salvatierra A, Carrascosa Gil A, Hualde Algarra A, Salvador Bravo M: Lingual tonsillar hypertrophy, an unknown enemy: A case report. *Colomb J Anesthesiol* 2019; 47:245-8
319. Markham TH, Nwokolo OO, Guzman-Reyes S, Medina-Rivera G, Gumbert SD, Cai C, Burnett T, Syed TA, Hagberg CA: A comparison of the King Vision® and Glidescope®

- video intubation systems in patients at risk for difficult intubation. *Trends Anaesth Crit Care* 2019; 28:27-35
320. Mathangi K, Mathews J, Mathangi CD: Assessment of perioperative difficult airway among undiagnosed obstructive sleep apnoea patients undergoing elective surgery: A prospective cohort study. *Indian J Anaesth* 2018; 62:538-44
  321. Mathew S, Chaudhuri S, Arun Kumar HD, Joseph TT: Airway management in Escobar syndrome: A formidable challenge. *Indian J Anaesth* 2013; 57:603-5
  322. Maurtua MA, Fernando M, Finnegan PS, Mehta B, Wu J, Foss J, Perilla M, Zura A, Doyle DJ: Use of the CTrach laryngeal mask airway in adult patients: A retrospective review of 126 cases. *J Clin Anesth* 2012; 24:370-2
  323. Maxey-Jones CL, Palmerton A, Farmer JR, Bateman BT: Difficult airway management caused by local anesthetic allergy during emergent cesarean delivery: A case report. *A A Case Rep* 2017; 9:84-6
  324. Mazzinari G, Rovira L, Henao L, Ortega J, Casasempere A, Fernandez Y, Acosta M, Belaouchi M, Esparza-Minana JM: Effect of dynamic versus stylet-guided intubation on first-attempt success in difficult airways undergoing glidescope laryngoscopy: A randomized controlled trial. *Anesth Analg* 2019; 128:1264-71
  325. McCaffer CJ, Douglas C, Wickham MH, Picozzi GL: Acute upper airway obstruction and emergency front of neck access in an achondroplastic patient. *BMJ Case Rep* 2015; 2015
  326. McCrirrick A, Pracilio JA: Awake intubation: A new technique. *Anaesthesia* 1991; 46:661-3
  327. McCrory CR, Moriarty DC: Laryngeal mask airway positioning is related to Mallampati grading in adults. *Anesth Analg* 1995; 81:1001-4
  328. McGoldrick KE, Donlon JV: Sublingual hematoma following difficult laryngoscopy. *Anesth Analg* 1979; 58:343-4
  329. McManus S, Jones L, Anstey C, Senthuran S: An assessment of the tolerability of the Cook staged extubation wire in patients with known or suspected difficult airways extubated in intensive care. *Anaesthesia* 2018; 73:587-93
  330. McQuibban GA: LMA-FasTrach. *Can J Anaesth* 1998; 45:95-6
  331. Meco BC, Alanoglu Z, Yilmaz AA, Basaran C, Alkis N, Demirer S, Cuhruk H: Does ultrasonographic volume of the thyroid gland correlate with difficult intubation? An observational study. *Braz J Anesthesiol* 2015; 65:230-4
  332. Mehta T, Jayaprakash J, Shah V: Diagnostic value of different screening tests in isolation or combination for predicting difficult intubation: A prospective study. *Indian J Anaesth* 2014; 58:756-7
  333. Messeter KH, Pettersson KI: Endotracheal intubation with the fibre-optic bronchoscope. *Anaesthesia* 1980; 35:294-8
  334. Michailidou M, O'Keeffe T, Mosier JM, Friese RS, Joseph B, Rhee P, Sakles JC: A comparison of video laryngoscopy to direct laryngoscopy for the emergency intubation of trauma patients. *World J Surg* 2015; 39:782-8
  335. Michálek P, Donaldson W, McAleavey F, Abraham A, Mathers RJ, Telford C: The i-gel supraglottic airway as a conduit for fibreoptic tracheal intubation - A randomized comparison with the single-use intubating laryngeal mask airway and Ctrach laryngeal mask in patients with predicted difficult laryngoscopy. *Prague Med Rep* 2016; 117:164-75
  336. Michalek P, Hodgkinson P, Donaldson W: Fiberoptic intubation through an I-gel supraglottic airway in two patients with predicted difficult airway and intellectual disability. *Anesth Analg* 2008; 106:1501-4
  337. Min JJ, Kim G, Kim E, Lee JH: The diagnostic validity of clinical airway assessments for predicting difficult laryngoscopy using a grey zone approach. *J Int Med Res* 2016; 44:893-904
  338. Miner JR, Rubin J, Clark J, Reardon RF: Retrograde intubation with an extraglottic device in place. *J Emerg Med* 2015; 49:864-7

339. Mishkel L, Wang JF, Gutierrez F, Ballard JB: Nasotracheal intubation by fiberoptic laryngoscope. *South Med J* 1981; 74:1407-9
340. Misquith JCR, Ribeiro KNSA: Anaesthetic management of a patient with a mobile pedunculated oropharyngeal mass. *J Clin Diagn Res* 2018; 12:UD03-05
341. Moda N, Kumar N: Combined use of video laryngoscopy and fiberoptic for airway management in a patient with fixed cervical spine. *Asian J Pharma Clin Res* 2018; 11:1-3
342. Modir H, Moshiri E, Malekianzadeh B, Noori G, Mohammadbeigi A: Endotracheal intubation in patients with difficult airway: Using laryngeal mask airway with bougie versus video laryngoscopy. *Med Gas Resch* 2017; 7:150-155
343. Mohammadi SS, Saliminia A, Nejatifard N, Azma R: Usefulness of ultrasound view of larynx in pre-anesthetic airway assessment: A comparison with cormack-lehane classification during direct laryngoscopy. *Anesth Pain Med* 2016; 6:e39566
344. Mohan K, Rupa LM, Krishna Murthy SG, Greesham PG, Bhavana U: Anaesthesia for TMJ ankylosis with the use of TIVA, followed by endotracheal intubation. *J Clin Diagn Res* 2012; 6:1765-7
345. Moon E, Jeong H, Chung J, Yi J: Central venous catheter malposition in the azygos vein and difficult endotracheal intubation in severe ankylosing spondylitis: A case report. *Int J Clin Exp Med* 2015; 8:21755-9
346. Moon TS, Fox PE, Somasundaram A, Minhajuddin A, Gonzales MX, Pak TJ, Ogunnaike B: The influence of morbid obesity on difficult intubation and difficult mask ventilation. *J Anesth* 2019; 33:96-102
347. Moriyama K, Mitsuda M, Kurita M, Ozaki M, Moriyama K, Yorozu T: When can we give general anesthesia to an infant with anticipated difficult airway management caused by facial vascular malformation? *JA Clin Rep* 2017; 3:12
348. Morrison S, Aerts S, Van Rompaey D, Vanderveken O: Failed awake intubation for critical airway obstruction rescued with the Ventrain device and an Arndt exchange catheter: A case report. *A A Pract* 2019; 13:23-6
349. Mort TC, Braffett BH: Conventional versus video laryngoscopy for tracheal tube exchange: glottic visualization, success rates, complications, and rescue alternatives in the high-risk difficult airway patient. *Anesth Analg* 2015; 121:440-8
350. Mort TC: Continuous airway access for the difficult extubation: The efficacy of the airway exchange catheter. *Anesth Analg* 2007; 105:1357-62
351. Mosier JM, Stolz U, Chiu S, Sakles JC: Difficult airway management in the emergency department: GlideScope videolaryngoscopy compared to direct laryngoscopy. *J Emerg Med* 2012; 42:629-34
352. Mostafa M, Saeed M, Hasanin A, Badawy S, Khaled D: Accuracy of thyromental height test for predicting difficult intubation in elderly. *J Anesth* 2020; 34:217-23
353. Mukaiharu K, Godai K, Yamada T, Hasegawa-Moriyama M, Kanmura Y: Successful airway management using a MultiViewScope handle with a stylet scope in a patient with Schwartz-Jampel syndrome. *JA Clin Rep* 2016; 2:36
354. Nagai K, Sakuramoto C, Goto F: Unilateral hypoglossal nerve paralysis following the use of the laryngeal mask airway. *Anaesthesia* 1994; 49:603-4
355. Nagamine Y, Kurahashi K: The use of three-dimensional computed tomography images for anticipated difficult intubation airway evaluation of a patient with Treacher Collins syndrome. *Anesth Analg* 2007; 105:626-8
356. Naithani U, Gupta G, Keerti, Gupta M, Meena K, Sharma CP, Bajaj P: Predicting difficult intubation in surgical patients scheduled for general anaesthesia: A prospective study of 435 patients. *J Evol Med Dent Sci* 2013; 2:2270-86
357. Nájera-Losada DC, Pérez-Moreno JC, Sanabria-Carretero P, Castro-Parga L E: Airtraq in difficult paediatric airway: Report of three cases. *Colombian J Anesthesiol* 2018; 46:168-172

358. Nakazawa K, Ikeda D, Ishikawa S, Makita K: A case of difficult airway due to lingual tonsillar hypertrophy in a patient with Down's syndrome. *Anesth Analg* 2003; 97:704-5
359. Narendra PL, Vishal NS, Jenkins B: Ludwig's angina: need for including airways and larynx in ultrasound evaluation. *BMJ Case Rep* 2014; 2014:09
360. Narkhede HH, Patel RD, Narkhede HR: A prospective observational study of predictors of difficult intubation in Indian patients. *J Anaesthesiol Clin Pharmacol* 2019; 35:119-23
361. Nasa P, Singh A, Juneja D, Garg N, Singh O, Javeri Y: Emergency percutaneous tracheostomy in two cancer patients with difficult airway: An alternative to cricothyroidotomy? *South Asian J Canc* 2012; 1:90-2
362. Nasa VK, Kamath SS: Risk factors assessment of the difficult intubation using Intubation Difficulty Scale (IDS). *J Clin Diagn Res* 2014; 8:GC01-3
363. Nasreen F, Khalid A: An infant with beals-hecht syndrome: An airway challenge for the anaesthesiologist. *Sri Lankan J Anaesthesiol* 2020; 28:150-2
364. Nasreen F, Prakash M, Khalid A, Hushain D: Airway management with Airtraq in a neonate with Epignathus. A case report. *Indian J Anaesth* 2020; 64:S245-S247
365. Nassar M, Zanaty OM, Ibrahim M: Bonfils fiberscope vs GlideScope for awake intubation in morbidly obese patients with expected difficult airways. *J Clin Anesth* 2016; 32:101-5
366. Neligan PJ, Porter S, Max B, Malhotra G, Greenblatt EP, Ochroch EA: Obstructive sleep apnea is not a risk factor for difficult intubation in morbidly obese patients. *Anesth Analg* 2009; 109:1182-1186
367. Neuman GG, Weingarten AE, Abramowitz RM, Kushins LG, Abramson AL, Ladner W: The anesthetic management of the patient with an anterior mediastinal mass. *Anesthesiology* 1984; 60:144-7
368. Ng I, Hill AL, Williams DL, Lee K, Segal R: Randomized controlled trial comparing the McGrath videolaryngoscope with the C-MAC videolaryngoscope in intubating adult patients with potential difficult airways. *Br J Anaesth* 2012; 109:439-43
369. Nguyen NH, Morvant EM, Mayhew JF: Anesthetic management for patients with arthrogryposis multiplex congenita and severe micrognathia: case reports. *J Clin Anesth* 2000; 12:227-30
370. Ni H, Guan C, He G, Bao Y, Shi D, Zhu Y: Ultrasound measurement of laryngeal structures in the parasagittal plane for the prediction of difficult laryngoscopies in Chinese adults. *BMC Anesthesiol* 2020; 20
371. Nishimori M, Matsumoto M, Nakagawa H, Ichiishi N: Unanticipated difficult airway due to undiagnosed oropharyngeal stenosis: A case report. *JA Clin Rep* 2016; 2:10
372. Nishinarita R, Mihara T, Nakamura N, Miyamoto Y, Ka K: Anesthetic management of pediatric patients with Emanuel syndrome. *J Anesth* 2015; 29:450-452
373. Noppens RR, Geimer S, Eisel N, David M, Piepho T: Endotracheal intubation using the C-MAC(R) video laryngoscope or the Macintosh laryngoscope: a prospective, comparative study in the ICU. *Crit Care* 2012; 16:R103
374. Noppens RR, Mobus S, Heid F, Schmidtman I, Werner C, Piepho T: Evaluation of the McGrath Series 5 videolaryngoscope after failed direct laryngoscopy. *Anaesthesia* 2010; 65:716-20
375. Normand KC, Vargas LA, Burnett T, Sridhar S, Cai C, Zhang X, Markham TH, Guzman-Reyes S, Hagberg CA: Use of the McGRATH™ MAC: To view or not to view? *Trends Anaesth Crit Care* 2018; 19:25-33
376. Northrip DR, Bohman BK, Tsueda K: Total airway occlusion and superior vena cava syndrome in a child with an anterior mediastinal tumor. *Anesth Analg* 1986; 65:1079-82
377. Nurullah M, Alam MS, Hossen M, Shahnawaz M: Prediction of difficult airway by thyromental height test-a comparison with modified mallampati test. *Bangladesh J Med Sci* 2018; 17:455-61
378. Oates JD, Macleod AD, Oates PD, Pearsall FJ, Howie JC, Murray GD: Comparison of two methods for predicting difficult intubation. *Br J Anaesth* 1991; 66:305-9

379. Oe Y, Godai K, Masuda M, Kanmura Y: Difficult airway associated with bifid glottis and coexistent subglottic stenosis in a patient with Pallister-Hall syndrome: a case report. *JA Clin Rep* 2018; 4:20
380. Ohsima N, Amaya F, Yamakita S, Nakayama Y, Kato H, Muranishi Y, Numajiri T, Sawa T: Difficult tracheal intubation and post-extubation airway stenosis in an 11-month-old patient with unrecognized subglottic stenosis: a case report. *JA Clin Rep* 2017; 3:10
381. Okumura Y, Okuda M, Sato Boku A, Tachi N, Hashimoto M, Yamada T, Yamada M: Usefulness of Airway Scope for intubation of infants with cleft lip and palate-comparison with Macintosh laryngoscope: a randomized controlled trial. *BMC Anesthesiol* 2019; 19
382. Ono K, Goto T, Nakai D, Ueki S, Takenaka S, Moriya T: Incidence and predictors of difficult nasotracheal intubation with airway scope. *J Anesth* 2014; 28:650-4
383. Osses H, Poblete M, Asenjo F: Laryngeal mask for difficult intubation in children. *Paediatr Anaesth* 1999; 9:399-401
384. Ovassapian A, Doka JC, Romsa DE: Acromegaly--use of fiberoptic laryngoscopy to avoid tracheostomy. *Anesthesiology* 1981; 54:429-30
385. Ovassapian A, Doka JC, Romsa DE: Failure to withdraw flexible fiberoptic laryngoscope after nasotracheal intubation. *Anesthesiology* 1985; 63:124-125
386. Ovassapian A, Doka JC, Romsa DE: Fiberoptic nasotracheal intubation - incidence and causes of failure. *Anesth Analg* 1983; 62:692-695
387. Ozdilek A, Beyoglu CA, Erbabacan SE, Ekici B, Altindas F, Vehid S, Koksall GM: Correlation of neck circumference with difficult mask ventilation and difficult laryngoscopy in morbidly obese patients: An observational study. *Obes Surg* 2018; 28:2860-7
388. Ozkan AS, Akbas S, Yalin MR, Ozdemir E, Koylu Z: Successful difficult airway management of a child with Coffin-siris syndrome. *Clin Case Rep* 2017; 5:1312-4
389. Packiasabapathy S, Chandiran R, Batra RK, Agarwala S: Difficult airway in Mowat-Wilson syndrome. *J Clin Anesth* 2016; 34:151-3
390. Pai Bh P, Shariat AN: Revisiting a case of difficult airway with a rigid laryngoscope. *BMJ Case Rep* 2019; 12:15
391. Palmer JH, Ball DR: Awake tracheal intubation with the intubating laryngeal mask in a patient with diffuse idiopathic skeletal hyperostosis. *Anaesthesia* 2000; 55:70-4
392. Pande A, Ramachandran R, Rewari V: Bougie-associated bronchial injury complicated by a nephropleural fistula after percutaneous nephrolithotomy: A tale of two complications. *BMJ Case Rep* 2018; 2018:17
393. Pandey R, Garg R, Kumar A, Darlong V, Punj J, Singh SA: Case report: Airway management of a patient with popping pedunculated subglottic laryngeal polyp. *Acta Anaesthesiol Belg* 2009; 60:251-3
394. Pang L, Feng YH, Ma HC, Dong S: Fiberoptic bronchoscopy-assisted endotracheal intubation in a patient with a large tracheal tumor. *Int Surg* 2015; 100:589-92
395. Pappu A, Sharma B, Jain R, Dua N, Sood J: A randomised comparative study of 'videoendoscope' with the Truview EVO2, C-MAC D blade videolaryngoscope and the Macintosh laryngoscope. *Indian J Anaesth* 2020; 64:S186-92
396. Parameswari A, Govind M, Vakamudi M: Correlation between preoperative ultrasonographic airway assessment and laryngoscopic view in adult patients: A prospective study. *J Anaesthesiol Clin Pharmacol* 2017; 33:353-8
397. Park CD, Lee HK, Yim JY, Kang IH: Anesthetic management for a patient with severe mento-sternal contracture: difficult airway and scarce venous access -A case report. *Korean J Anesthesiol* 2013; 64:61-4
398. Park R, Peyton JM, Fiadjoe JE, Hunyady AI, Kimball T, Zurakowski D, Kovatsis PG, Bosenberg A, Hopkins P, Glover C, Olutoye O, Szmuk P, Olomu P, Jagannathan N, Burjek N, Watkins S, Reynolds P, Haydar B, Matuszczak M, Jain R, Khalil S, Polaner D, Zieg J, Szolnoki J, Sathyamoorthy M, Taicher B, Bhattacharya S, Raman V, Bhalla T, Stricker P, Lockman J, Galvez J, Rehman M, Nishisaki A, Von Ungern-Sternberg B,

- Sommerfield D, Soneru C, Chiao F, Richtsfeld M, Belani K, Sarmiento L, Mireles S, Rosas GB: The efficacy of GlideScope® videolaryngoscopy compared with direct laryngoscopy in children who are difficult to intubate: An analysis from the paediatric difficult intubation registry. *Br J Anaesth* 2017; 119:984-992
399. Park YH, Kim SH, Lee SJ, Yang J, Kim H: Fiberoptic intubation in patient who have had unilateral radical maxillectomy – A case report. *Rawal Med J* 2020; 45:990-3
  400. Parmet JL, Colonna-Romano P, Horrow JC, Miller F, Gonzales J, Rosenberg H: The laryngeal mask airway reliably provides rescue ventilation in cases of unanticipated difficult tracheal intubation along with difficult mask ventilation. *Anesth Analg* 1998; 87:661-5
  401. Parr MJ, Gregory M, Baskett PJ: The intubating laryngeal mask. Use in failed and difficult intubation. *Anaesthesia* 1998; 53:343-8
  402. Patel A, Venn PJ, Barham CJ: Fibreoptic intubation through a laryngeal mask airway in an infant with Robin sequence. *Eur J Anaesthesiol* 1998; 15:237-9
  403. Patel B, Khandekar R, Diwan R, Shah A: Validation of modified mallampati test with addition of thyromental distance and sternomental distance to predict difficult endotracheal intubation in adults. *Indian J Anaesth* 2014; 58:171-5
  404. Pavoni V, Froio V, Nella A, Simonelli M, Gianesello L, Horton A, Malino L, Micaglio M: Tracheal intubation with Aura-i and aScope-2: How to minimize apnea time in an unpredicted difficult airway. *Case Rep Anesthesiol* 2015; 2015:453547
  405. Pellis T, Leykin Y, Albano G, Zannier G, Di Capua G, Marzano B, Gullo A: Perioperative management and monitoring of a super-obese patient. *Obes Surg* 2004; 14:1423-7
  406. Petrișor C, Trancă S, Szabo R, Simon R, Prie A, Bodolea C: Clinical versus ultrasound measurements of hyomental distance ratio for the prediction of difficult airway in patients with and without morbid obesity. *Diagnostics* 2020; 10
  407. Phillips M, Jagannathan N: Placement of a supraglottic airway to overcome airway obstruction when performing nasal fiberoptic intubation in infants with pierre robin sequence: A case series. *A A Prac* 2020
  408. Pieters BM, Theunissen M, van Zundert AA: Macintosh blade videolaryngoscopy combined with rigid bonfils intubation endoscope offers a suitable alternative for patients with difficult airways. *Anesth Analg* 2018; 126:988-994
  409. Pinosky ML, Hardin CL, Bach DE, Shuman K: The reinforced laryngeal mask airway (LMA) as an alternative airway device to manage the difficult airway. *Pediatr Dent* 1998; 20:422-4
  410. Pinto J, Cordeiro L, Pereira C, Gama R, Fernandes HL, Assuncao J: Predicting difficult laryngoscopy using ultrasound measurement of distance from skin to epiglottis. *J Crit Care* 2016; 33:26-31
  411. Piro AJ, Weiss DR, Hellman S: Mediastinal Hodgkin's disease: A possible danger for intubation anesthesia. Intubation danger in Hodgkin's disease. *Int J Radiat Oncol Biol Phys* 1976; 1:415-9
  412. Pius J, Ioanidis K, Noppens RR: Use of the novel C-MAC video stylet in a case of predicted difficult intubation: A case report. *A A Pract* 2019
  413. Pollard BA, El-Beheiry H: Pott's disease with unstable cervical spine, retropharyngeal cold abscess and progressive airway obstruction. *Can J Anaesth* 1999; 46:772-5
  414. Portereiko JV, Perez MM, Hojman H, Frankel HL, Rabinovici R: Acute upper airway obstruction by an over-inflated Combitube esophageal obturator balloon. *J Trauma* 2006; 60:426-7
  415. Pradhan D, Bhattacharyya P: Difficult airway management from Emergency Department till Intensive Care Unit. *Indian J Crit Care Med* 2015; 19:557-9
  416. Prakash S, Kumar A, Bhandari S, Mullick P, Singh R, Gogia AR: Difficult laryngoscopy and intubation in the Indian population: An assessment of anatomical and clinical risk factors. *Indian J Anaesth* 2013; 57:569-75

417. Prakash S, Mullick P, Bhandari S, Kumar A, Gogia AR, Singh R: Sternomental distance and sternomental displacement as predictors of difficult laryngoscopy and intubation in adult patients. *Saudi J Anaesth* 2017; 11:273-8
418. Preis C, Czerny C, Preis I, Zimpfer M: Variations in ILMA external diameters: another cause of device failure. *Can J Anaesth* 2000; 47:886-9
419. Preis CA, Hartmann T, Zimpfer M: Laryngeal mask airway facilitates awake fiberoptic intubation in a patient with severe oropharyngeal bleeding. *Anesth Analg* 1998; 87:728-9
420. Rajendra Kulkarni K, Namazi I, Madnaik S: Failed fiberoptic intubation and surgical tracheostomy in a case of Down's syndrome with goiter. *Sri Lankan J Anaesthesiol* 2013; 21:83-5
421. Ramamani M, Ponnaiah M, Bhaskar S, Rai E: An uncommon cause of unanticipated difficult airway. *Paediatr Anaesth* 2009; 19:643-5
422. Ramkisson A, Hodgson RE: Nasal intubation of a difficult airway following supraglottic airway rescue facilitated by video laryngoscopy and a flexible intubation scope. *South Afr J Anaesth Analg* 2019; 25:42-5
423. Rana S, Verma V, Bhandari S, Sharma S, Koundal V, Chaudhary SK: Point-of-care ultrasound in the airway assessment: A correlation of ultrasonography-guided parameters to the Cormack-Lehane Classification. *Saudi J Anaesth* 2018; 12:292-6
424. Rao CS, Ranganath T, Rao SPB, Sujani K: Comparison of upper lip bite test with Modified Mallampati test and thyromental distance for predicting difficulty in endotracheal intubation- A prospective study. *J Evol Med Dent Sci* 2017; 6:1413-6
425. Rao KVN, Dhatchinamoorthi D, Nandhakumar A, Selvarajan N, Akula HR, Thiruvengkatarajan V: Validity of thyromental height test as a predictor of difficult laryngoscopy: A prospective evaluation comparing modified Mallampati score, interincisor gap, thyromental distance, neck circumference, and neck extension. *Indian J Anaesth* 2018; 62:603-8
426. Rao L, Jumana H, Gururajrao M, Venkatesh KH: Successful management of difficult airway in children with the use of adult fiberoptic bronchoscope. *Indian J Anaesth* 2015; 59:50-1
427. Rao TL, Mathru M, Gorski DW, Salem MR: Experience with a new intubation guide for difficult tracheal intubation. *Crit Care Med* 1982; 10:882-3
428. Rasch DK, Browder F, Barr M, Greer D: Anaesthesia for Treacher Collins and Pierre Robin syndromes: A report of three cases. *Can Anaesth Soc J* 1986; 33:364-70
429. Raval CB, Khan S: Airway management in submandibular abscess patient with awake fiberoptic intubation--a case report. *Middle East J Anaesthesiol* 2012; 21:647-51
430. Ravalia A, Goddard JM: The laryngeal mask and difficult tracheal intubation. *Anaesthesia* 1990; 45:168
431. Rawal P, Shrestha SM: The evaluation of thyromental height test as a single, accurate predictor of difficult laryngoscopy. *J Nepal Health Res Counc* 2020; 18:271-6
432. Redden RL, Biery KA, Campbell RL: Arterial oxygen desaturation during awake endotracheal intubation. *Anesth Prog* 1990; 37:201-4
433. Reddy PB, Punetha P, Chalam KS: Ultrasonography - A viable tool for airway assessment. *Indian J Anaesth* 2016; 60:807-13
434. Reena, Kumar A, Singh SK, Agrawal V: Unsuspected subglottic web in a child managed for severe respiratory obstruction. *Saudi J Anaesth* 2017; 11:99-101
435. Reena, Vikram A: Maxillary tumor in a child: An expected case of difficult airway. *Saudi J Anaesth* 2016; 10:233-5
436. Rhee KY, Lee JR, Kim J, Park S, Kwon WK, Han S: A comparison of lighted stylet (Surch-Lite) and direct laryngoscopic intubation in patients with high Mallampati scores. *Anesth Analg* 2009; 108:1215-9
437. Riad W, Ansari T, Shetty N: Does neck circumference help to predict difficult intubation in obstetric patients? A prospective observational study. *Saudi J Anaesth* 2018; 12:77-81

438. Riad W, Vaez MN, Raveendran R, Tam AD, Quereshy FA, Chung F, Wong DT: Neck circumference as a predictor of difficult intubation and difficult mask ventilation in morbidly obese patients: A prospective observational study. *Eur J Anaesthesiol* 2016; 33:244-9
439. Richa F: Intubating laryngeal mask airway combined to fiberoptic intubation in subglottic stenosis. *BMJ Case Rep* 2013
440. Roa NL, Moss KS: Treacher-Collins syndrome with sleep apnea: anesthetic considerations. *Anesthesiology* 1984; 60:71-3
441. Rocke DA, Murray WB, Rout CC, Gouws E: Relative risk analysis of factors associated with difficult intubation in obstetric anesthesia. *Anesthesiology* 1992; 77:67-73
442. Rogers SN, Benumof JL: New and easy techniques for fiberoptic endoscopy-aided tracheal intubation. *Anesthesiology* 1983; 59:569-72
443. Roodneshin F: Sevoflurane as the single anesthetic agent for management of anticipated pediatric difficult airway. *Tanaffos* 2012; 11:69-72
444. Rosenblatt W, Ianus AI, Sukhupragarn W, Fickenscher A, Sasaki C: Preoperative endoscopic airway examination (PEAE) provides superior airway information and may reduce the use of unnecessary awake intubation. *Anesth Analg* 2011; 112:602-7
445. Rosenstock CV, Thogersen B, Afshari A, Christensen AL, Eriksen C, Gatke MR: Awake fiberoptic or awake video laryngoscopic tracheal intubation in patients with anticipated difficult airway management: A randomized clinical trial. *Anesthesiology* 2012; 116:1210-6
446. Saasouh W, Laffey K, Turan A, Avitsian R, Zura A, You J, Zimmerman NM, Szarpak L, Sessler DI, Ruetzler K: Degree of obesity is not associated with more than one intubation attempt: A large centre experience. *Br J Anaesth* 2018; 120:1110-6
447. Sabapathy S, Kulkarni P, Hanumanthappa N, Sarangi TK: Caught in the subglottic web- unanticipated difficult intubation. *Biomed Res (Aligarh)* 2018; 29:3174-6
448. Saini S, Hooda S, Nandini S, Sekhri C: Difficult airway management in a maxillofacial and cervical abnormality with intubating laryngeal mask airway. *J Oral Maxillofac Surg* 2004; 62:510-3
449. Salhotra R, Sharma C, Tyagi A, Kumar S, Sethi A, Bhatt S: An unanticipated difficult airway in Lesch-Nyhan syndrome. *J Anaesthesiol Clin Pharmacol* 2012; 28:239-41
450. Salzarulo HH, Taylor LA: Diabetic "stiff joint syndrome" as a cause of difficult endotracheal intubation. *Anesthesiology* 1986; 64:366-8
451. Samra T, Banerjee N: Anaesthesia for emergency ventriculo-peritoneal shunt in an adolescent with Noonan's Syndrome. *Indian J Anaesth* 2014; 58:452-5
452. Sanchez-Morillo J, Estruch-Perez MJ, Hernandez-Cadiz MJ, Tamarit-Conejeros JM, Gomez-Diago L, Richart-Aznar M: Indirect laryngoscopy with rigid 70-degree laryngoscope as a predictor of difficult direct laryngoscopy. *Acta Otorrinolaringol Esp* 2012; 63:272-9
453. Sardar A, Khanna P, Singh A, Sharma A: Long-standing meningomyelocele can be a predictor of difficult airway and postoperative hypoventilation: challenge to the anaesthesiologist. *BMJ Case Rep* 2016; 2016:16
454. Saricicek V, Mizrak A, Gul R, Goksu S, Cesur M: GlideScope video laryngoscopy use tracheal intubation in patients with ankylosing spondylitis: a series of four cases and literature review. *J Clin Monit Comput* 2014; 28:169-72
455. Sarkar S, Jafra A, Mathew P: Emergency airway management in Pierre Robin Sequence, our nightmare experiences. *Trends Anaesth Crit Care* 2020
456. Sarma J: Unexpected difficult intubation in a patient with prominent mandibular tori. *Open Anesthesiol J* 2015; 8
457. Saruki N, Saito S, Sato J, Takahashi T, Tozawa R: Difficult airway management with the combination of a fiberoptic stylet and McCoy laryngoscope. *Can J Anaesth* 2001; 48:212
458. Savva D: Prediction of difficult tracheal intubation. *Br J Anaesth* 1994; 73:149-53
459. Sawyer T, Foglia EE, Ades A, Moussa A, Napolitano N, Glass K, Johnston L, Jung P, Singh N, Quek BH, Barry J, Zenge J, DeMeo SD, Brei B, Krick J, Kim JH, Nadkarni V, Nishisaki A, National Emergency Airway Registry for Neonates: Incidence, impact and



- indicators of difficult intubations in the neonatal intensive care unit: a report from the National Emergency Airway Registry for Neonates. *Arch Dis Child Fetal Neonatal Ed* 2019; 104:F461-6
460. Saxena KN, Bansal P: Endotracheal intubation under local anesthesia and sedation in an infant with difficult airway. *J Anaesthesiol Clin Pharmacol* 2012; 28:358-60
  461. Schmitt HJ, Mang H: Head and neck elevation beyond the sniffing position improves laryngeal view in cases of difficult direct laryngoscopy. *J Clin Anesth* 2002; 14:335-8
  462. Schuschnig C, Walzl B, Erlacher W, Reddy B, Stoik W, Kapral S: Intubating laryngeal mask and rapid sequence induction in patients with cervical spine injury. *Anaesthesia* 1999; 54:793-7
  463. Selim M, Mowafi H, Al-Ghamdi A, Adu-Gyamfi Y: Intubation via LMA in pediatric patients with difficult airways. *Can J Anaesth* 1999; 46:891-3
  464. Selvi O, Kahraman T, Senturk O, Tulgar S, Serifsoy E, Ozer Z: Evaluation of the reliability of preoperative descriptive airway assessment tests in prediction of the Cormack-Lehane score: A prospective randomized clinical study. *J Clin Anesth* 2017; 36:21-6
  465. Sen R, Mallepally AR, Sakrikar G, Marathe N, Rathod T: Comparison of TruView and King Vision video laryngoscopes in subaxial cervical spine injury: A randomized controlled trial. *Surg Neurol Int* 2020; 11
  466. Seo SH, Lee JG, Yu SB, Kim DS, Ryu SJ, Kim KH: Predictors of difficult intubation defined by the intubation difficulty scale (IDS): predictive value of 7 airway assessment factors. *Korean J Anesthesiol* 2012; 63:491-7
  467. Serdiuk AA, Bosek V: An adult patient with Klippel-Feil syndrome presenting for repeat operation: a cautionary tale of the GlideScope. *J Clin Anesth* 2012; 24:238-41
  468. Serocki G, Bein B, Scholz J, Döriges V: Management of the predicted difficult airway: A comparison of conventional blade laryngoscopy with video-assisted blade laryngoscopy and the GlideScope. *Eur J Anaesthesiol* 2010; 27:24-30
  469. Serocki G, Neumann T, Scharf E, Dorges V, Cavus E: Indirect videolaryngoscopy with C-MAC D-Blade and GlideScope: a randomized, controlled comparison in patients with suspected difficult airways. *Minerva Anesthesiol* 2013; 79:121-9
  470. Sethi S, Arora V: Use of glidescope and external manipulation in airway management of an unusual retropharyngeal lipoma. *J Anaesthesiol Clin Pharmacol* 2010; 26:557-8
  471. Sethi SK, Jain N, Khare A, Patodi V: Anaesthetic management in a case of large plunging ranula with difficult airway: A case report. *Egypt J Anaesth* 2017; 33:209-12
  472. Shah AA, Rafique K, Islam M: Can difficult intubation be accurately predicted using upper lip bite test? *J Postgrad Med Instit* 2014; 28:282-7
  473. Shah AC, Ng WCK, Sinnott S, Cravero JP: Population analysis of predictors of difficult intubation with direct laryngoscopy in pediatric patients with and without thyroid disease. *J Anesth* 2018; 32:54-61
  474. Shah PJ, Dubey KP, Yadav JP: Predictive value of upper lip bite test and ratio of height to thyromental distance compared to other multivariate airway assessment tests for difficult laryngoscopy in apparently normal patients. *J Anaesthesiol Clin Pharmacol* 2013; 29:191-5
  475. Shah PN, Sundaram V: Incidence and predictors of difficult mask ventilation and intubation. *J Anaesthesiol Clin Pharmacol* 2012; 28:451-5
  476. Shamim F, Yahya M, Ikram M: Awake fiberoptic intubation in a patient with known difficult airway due to huge thyroid goiter. *Anaesth Pain Intensive Care* 2017; 21:94-7
  477. Shaw JC, Welch EA, Harrison BJ, Michael S: Complete airway obstruction during awake fiberoptic intubation. *Anaesthesia* 1997; 52:582-585
  478. Sheeran P, Maguire T, Browne G: Mechanical failure of the McCoy laryngoscope during difficult intubation. *Anaesthesia* 2000; 55:184-5

479. Sheff SR, May MC, Carlisle SE, Kallies KJ, Mathiason MA, Kothari SN: Predictors of a difficult intubation in the bariatric patient: Does preoperative body mass index matter? *Surg Obes Relat Dis* 2013; 9:344-9
480. Shetty SR, V.T S: Validation of clinical versus ultrasound parameters in assessment of airway. *Trends Anaesth Crit Care* 2020; 35:21-7
481. Shigematsu T, Miyazawa N, Kobayashi M, Yorozu T, Toyoda Y, Morisaki H: Nasal intubation with Bullard laryngoscope: a useful approach for difficult airways. *Anesth Analg* 1994; 79:132-135
482. Shih CC, Wang JC, Chen SJ, Hsu YP: Focused ultrasound assists in diagnosis and management of difficult airway in Ludwig's angina. *J Med Ultrasound* 2019; 27:61-62
483. Shindo Y, Toda S, Kido K, Masaki E: Massive ameloblastoma: A case report of difficult fiberoptic intubation. *Ann Med Surg* 2018; 32:6-9
484. Shivakumar PS, Sarvesh NK, Patil V, Sagar GC, Sirigeri S, Monnaiah KP, Gundappa RS: Ratio of patient's height to thyromental distance (RHTMD) compared to thyromental distance for prediction of difficult intubation. *J Evol Med Dent Sci* 2016; 5:4462-7
485. Shollik NA, Ibrahim SM, Ismael A, Agnoletti V, Piraccini E, Corso RM: Use of the Bonfils intubation fiberscope in patients with limited mouth opening. *Case Rep Anesthesiol* 2012; 2012:297306
486. Shraavanalakshmi D, Bidkar PU, Narmadalakshmi K, Lata S, Mishra SK, Adinarayanan S: Comparison of intubation success and glottic visualization using King Vision and C-MAC videolaryngoscopes in patients with cervical spine injuries with cervical immobilization: A randomized clinical trial. *Surg Neurol Int* 2017; 8:19
487. Shukla P, Yadav SS, Jain V, Sharma G, Chourasia HK, Choudhary B: A comparative evaluation of Airtraq and Macintosh laryngoscopes for difficult intubation. *J Evolution Med Dent Sci* 2017; 6:5098-5102
488. Shukry M, Hanson RD, Koveleskie JR, Ramadhyani U: Management of the difficult pediatric airway with Shikani optical stylet. *Paediatr Anaesth* 2005; 15:342-5
489. Shung J, Avidan MS, Ing R, Klein DC, Pott L: Awake intubation of the difficult airway with the intubating laryngeal mask airway. *Anaesthesia* 1998; 53:645-9
490. Shyam R, Chaudhary AK, Sachan P, Singh PK, Singh GP, Bhatia VK, Chandra G, Singh D: Evaluation of Fastrach laryngeal mask airway as an alternative to fiberoptic bronchoscope to manage difficult airway: a comparative study. *J Clin Diagn Res* 2017; 11:UC09-12
491. Siddiqui S, Seet E, Chan WY: The use of laryngeal mask airway Supreme™ in rescue airway situation in the critical care unit. *Singapore Med J* 2014; 55:e205-6
492. Sidhu VS, Whitehead EM, Ainsworth QP, Smith M, Calder I: A technique of awake fibreoptic intubation. Experience in patients with cervical spine disease. *Anaesthesia* 1993; 48:910-3
493. Sime J, Bailitz J, Moskoff J: The bougie: An inexpensive lifesaving airway device. *J Emerg Med* 2012; 43:e393-5
494. Singh J, Yadav MK, Marahatta SB, Lal Shrestha B: Randomized crossover comparison of the laryngeal mask airway classic with i-gel laryngeal mask airway in the management of difficult airway in post burn neck contracture patients. *Indian J Anaesth* 2012; 56:348-52
495. Singh N, Rao PB, Samal RL: TruView video laryngoscope for lateral position intubation in a patient with giant presacral neurofibroma. *J Emerg Med* 2019; 57:380-2
496. Sinha R, Rewari V, Varma P, Kumar A: Successful use of C-Mac video laryngoscope in a child with large parapharyngeal mass. *Paediatr Anaesth* 2014; 24:531-3
497. Sizlan A, Ozhan MO, Cekmen N, Suzer MA, Orhan ME, Comak I, Sahin S, Sahin I: Nasogastric tube aided fiberoptic intubation through laryngeal mask airway (case report). *Acta Anaesthesiol Belg* 2009; 60:185-8
498. Sjogren P, Pedersen T: Anaesthetic problems in Hurler-Scheie syndrome. Report of two cases. *Acta Anaesthesiol Scand* 1986; 30:484-6

499. Smith BB, Barbara DW, Hyder JA, Smith MM: Anesthetic considerations for patients with Bardet-Biedl syndrome: a case series and review of the literature. *Paediatr Anaesth* 2016; 26:429-37
500. Smith M, Crockyard A, Calder I, Isert P, Nicol ME: Oxygen saturation and cardiovascular changes during fiberoptic intubation under general anaesthesia. *Anaesthesia* 1992; 47:158-161
501. Sohn L, Sawardekar A, Jagannathan N: Airway management options in a prone achondroplastic dwarf with a difficult airway after unintentional tracheal extubation during a wake-up test for spinal fusion: to flip or not to flip? *Can J Anaesth* 2014; 61:741-4
502. Song JA, Bae HB, Choi JI, Kang J, Jeong S: Difficult intubation and anesthetic management in an adult patient with undiagnosed congenital tracheal stenosis: a case report. *J Int Med Res* 2020; 48
503. Sowers N, Kovacs G: Use of a flexible intubating scope in combination with a channeled video laryngoscope for managing a difficult airway in the emergency department. *J Emerg Med* 2016; 50:315-9
504. Srinivasan C, Kuppaswamy B: Comparison of validity of airway assessment tests for predicting difficult intubation. *Indian Anaesth Forum* 2017; 18:63-8
505. Srinivasan G, Sivakumar RK, Bidkar P, Sharma D: Paediatric King Vision((R)) videolaryngoscope in a case of infantile oral mass: A useful alternative to fiberoptic bronchoscope as a first choice in paediatric difficult airway. *Indian J Anaesth* 2019; 63:325-7
506. Srivastava D, Dhiraaj S: Airway management of a difficult airway due to prolonged enlarged goiter using loco-sedative technique. *Saudi J Anaesth* 2013; 7:86-9
507. Srivilaithon W, Muengtaweepongsa S, Sittichanbuncha Y, Patumanond J: Predicting difficult intubation in emergency department by intubation assessment score. *J Clin Med Res* 2018; 10:247-53
508. St. Mont G, Biesler I, Pförtner R, Mohr C, Groeben H: Easy and difficult nasal intubation - A randomised comparison of Macintosh vs Airtraq® laryngoscopes. *Anaesthesia* 2012; 67:132-138
509. Stevic M, Bokun Z, Milojevic I, Budic I, Jovanovic B, Krstic Z, Simic D: Management of anesthesia in a child with a large neck rhabdoid tumor. *Med Princ Pract* 2016; 25:290-2
510. Stone DJ, Stirt JA, Kaplan MJ, McLean WC: A complication of lightwand-guided nasotracheal intubation. *Anesthesiology* 1984; 61:780-1
511. Strutt JR, Thompson NR, Stotesbery JL, Horvath B: Emergency endotracheal intubation with a rigid stylet of an infant with severe subglottic stenosis. *J Emerg Med* 2020; 58:e157-60
512. Subedi A, Tripathi M, Bhattarai B, Pokharel K, Dhital D: Successful intubation with McCoy laryngoscope in a patient with ankylosing spondylitis. *J Nepal Health Res Counc* 2014; 12:70-2
513. Subhash, Dhama VK, Manik YK, Tiwari T, Singh G: Use of lightwand for nasotracheal intubation in adult patients with limited mouth opening undergoing elective surgery. *Anaesth Pain Intensive Care* 2015; 19:468-72
514. Sugita T, Arisaka H: AirWay Scope™ for difficult ventilation in a patient with epiglottic cyst. *Anesth Prog* 2018; 65:204-5
515. Sung JK, Kim HG, Kim JE, Jang MS, Kang JM: Endotracheal tube intubation with the aid of a laryngeal mask airway, a fiberoptic bronchoscope, and a tube exchanger in a difficult airway patient: a case report. *Korean J Anesthesiol* 2014; 66:237-9
516. Sutton CD, Carvalho B: Supraglottic airway rescue after failed fiberoptic intubation in a patient with osteogenesis imperfecta: A case report. *A A Pract* 2019; 13:7-9
517. Tafesse D, Ataro G: Predictors of difficult tracheal intubation on adult elective patients in a teaching hospital. *Open Anesthesiol J* 2016; 10:34-9

518. Tai FMT, Ho YL, Cheng BCP, Mui KSY, Chan MTV, Wong OF: The use of the Airtraq (R) in awake tracheal intubation: a case series. *Hong Kong J Emerg Med* 2015; 22:248-255
519. Takaishi K, Kawahito S, Tomioka S, Eguchi S, Kitahata H: Cuffed oropharyngeal airway for difficult airway management. *Anesth Prog* 2014; 61:107-10
520. Takeshita S, Ueda H, Goto T, Muto D, Kakita H, Oshima K, Tainaka T, Ono T, Kazaoka Y, Yamada Y: Case report of Pierre Robin sequence with severe upper airway obstruction who was rescued by fiberoptic nasotracheal intubation. *BMC Anesthesiol* 2017; 17
521. Tamire T, Demelash H, Admasu W: Predictive values of preoperative tests for difficult laryngoscopy and intubation in adult patients at Tikur Anbessa specialized hospital. *Anesthesiol Res Pract* 2019; 2019:1790413
522. Tantia P, Malik S, Jamil S, Samal R: Intubation with Airtraq™ laryngoscope in a morbidly obese patient. *Saudi J Anaesth* 2011; 5: 85–86
523. Tantri AR, Firdaus R, Salomo ST: Predictors of Difficult intubation among Malay patients in Indonesia. *Anesth Pain Med* 2016; 6:e34848
524. Tassonyi E, Lehmann C, Gunning K, Coquoz E, Montandon D: Fiberoptically guided intubation in children with gangrenous stomatitis (noma). *Anesthesiology* 1990; 73:348-9
525. Tekeli AE, Eker E, Bartın MK, Öner MÖ: Anesthesia management in laparoscopic sleeve gastrectomy cases. *East J Med* 2019; 24:335-9
526. Theiler L, Kleine-Brueggeney M, Urwyler N, Graf T, Luyet C, Greif R: Randomized clinical trial of the i-gel and Magill tracheal tube or single-use ILMA and ILMA tracheal tube for blind intubation in anaesthetized patients with a predicted difficult airway. *Br J Anaesth* 2011; 107:243-50
527. Thompson C, Moga R, Crosby ET: Failed videolaryngoscope intubation in a patient with diffuse idiopathic skeletal hyperostosis and spinal cord injury. *Can J Anaesth* 2010; 57:679-82
528. Thompson NCP: Concurrent use of videolaryngoscope and fiberoptic bronchoscope in a child with neurofibromatosis to facilitate endotracheal intubation. *J Nat Med Assoc* 2020
529. Thomsen JLD, Norskov AK, Rosenstock CV: Supraglottic airway devices in difficult airway management: A retrospective cohort study of 658,104 general anaesthetics registered in the Danish anaesthesia database. *Anaesthesia* 2019; 74:151-7
530. Thomson KD, Ordman AJ, Parkhouse N, Morgan BD: Use of the Brain laryngeal mask airway in anticipation of difficult tracheal intubation. *Br J Plast Surg* 1989; 42:478-80
531. Toshniwal G, McKelvey GM, Wang H: STOP-Bang and prediction of difficult airway in obese patients. *J Clin Anesth* 2014; 26:360-7
532. Tremblay MH, Williams S, Robitaille A, Drolet P: Poor visualization during direct laryngoscopy and high upper lip bite test score are predictors of difficult intubation with the GlideScope videolaryngoscope. *Anesth Analg* 2008; 106:1495-500
533. Tse JC, Rimm EB, Hussain A: Predicting difficult endotracheal intubation in surgical patients scheduled for general anesthesia: A prospective blind study. *Anesth Analg* 1995; 81:254-8
534. Tsuchiya M, Terai H, Mizutani K, Funai Y, Tanaka K, Yamada T, Mori T, Nishikawa K: General anesthesia management for adult mucopolysaccharidosis patients undergoing major spine surgery. *Med Princ Pract* 2019; 28:581-5
535. Tsukamoto M, Hitosugi T, Yokoyama T: Flexible laryngeal mask airway management for dental treatment cases associated with difficult intubation. *J Dent Anesth Pain Med* 2017; 17:61-4
536. Tungaria H, Raiger LK, Paliwal R, Saxena SS, Bairwa BK: Palatonasal fistula repair - A case of unanticipated difficult intubation. *J Clin Diagn Res* 2016; 10:UD01-02
537. Tutuncu AC, Erbabacan E, Teksoz S, Ekici B, Koksall G, Altintas F, Kaya G, Ozcan M: The assessment of risk factors for difficult intubation in thyroid patients. *World J Surg* 2018; 42:1748-53

538. Uakritdathikarn T: Lightwand-assisted nasotracheal intubation in awake ankylosing spondylitis. *J Med Assoc Thai* 2006; 89:1976-80
539. Ul Haq MI, Shamim F, Lal S, Shafiq F: Airway management in a patient with severe ankylosing spondylitis causing bamboo spine: Use of Aintree intubation catheter. *J Coll Physicians Surg Pak* 2015; 25:900-2
540. Ul Haq MI, Ullah H: Comparison of Mallampati test with lower jaw protrusion maneuver in predicting difficult laryngoscopy and intubation. *J Anaesthesiol Clin Pharmacol* 2013; 29:313-7
541. Uribe AA, Zvara DA, Puente EG, Otey AJ, Zhang J, Bergese SD: BMI as a predictor for potential difficult tracheal intubation in males. *Front Med* 2015; 2:38
542. Ushiroda J, Inoue S, Egawa J, Kawano Y, Kawaguchi M, Furuya H: Life-threatening airway obstruction due to upper airway edema and marked neck swelling after labor and delivery. *Braz J Anesthesiol* 2013; 63:508-10
543. Usui T, Saito S, Goto F: Arytenoid dislocation while using a McCoy laryngoscope. *Anesth Analg* 2001; 92:1347-8
544. Van Zundert T, Wong D, Marcus M, Brimacombe JR: Prospective evaluation of the LMA-Supreme™ as an airway intubation conduit in patients with a predicted difficult airway. *Eur J Anaesthesiol* 2012; 29:240
545. Van Zundert TCRV, Wong DT, Van Zundert AAJ: The LMA-Supreme™ as an intubation conduit in patients with known difficult airways: A prospective evaluation study. *Acta Anaesthesiol Scand* 2013; 57:77-81
546. Verchere S, Khalil B, Maddukuri V, Hagberg CA: Use of the DCI video laryngoscope system in a pediatric patient with amniotic band syndrome and craniofacial abnormalities affecting the airway. *J Clin Anesth* 2012; 24:151-4
547. Vest D, Lee D, Newcome K, Stamper H: A retrospective review of difficult intubations: is obstructive sleep apnea a predictor? *Clin Nurse Spec* 2013; 27:128-31
548. Viderman D, Nurpeisov A, Balabayev O, Urunbayev Y, de Almeida G, Bilotta F: Hydatid cyst in the cervical spinal cord complicated by potentially life-threatening difficult airway: A case report. *Braz J Anesthesiol* 2020
549. Vinayagam S, Dhanger S, Tilak P, Gnanasekar R: C-MAC® video laryngoscope with D-BLADE™ and Frova introducer for awake intubation in a patient with parapharyngeal mass. *Saudi J Anaesth* 2016; 10:471-3
550. Vincent RD, Jr., Wimberly MP, Brockwell RC, Magnuson JS: Soft palate perforation during orotracheal intubation facilitated by the GlideScope videolaryngoscope. *J Clin Anesth* 2007; 19:619-21
551. Wahba SS, Tammam TF, Saeed AM: Comparative study of awake endotracheal intubation with Glidescope video laryngoscope versus flexible fiber optic bronchoscope in patients with traumatic cervical spine injury. *Egypt J Anaesth* 2012; 28:257-60
552. Wahlen BM, Al-Thani H, El-Menyar A: Ventrain: from theory to practice. Bridging until re-tracheostomy. *BMJ Case Rep* 2017; 2017:16
553. Wajekar AS, Chellam S, Toal PV: Prediction of ease of laryngoscopy and intubation-role of upper lip bite test, modified mallampati classification, and thyromental distance in various combination. *J Fam Med Prim Care* 2015; 4:101-5
554. Wakeling HG, Ody A, Ball A: Large goitre causing difficult intubation and failure to intubate using the intubating laryngeal mask airway: lessons for next time. *Br J Anaesth* 1998; 81:979-81
555. Walker RW: The laryngeal mask airway in the difficult paediatric airway: an assessment of positioning and use in fiberoptic intubation. *Paediatr Anaesth* 2000; 10:53-8
556. Walls RD, Timmis DP, Finucane BT: Difficult intubation associated with calcified stylohyoid ligament. *Anaesth Intens Care* 1990; 18:110-2

557. Wang B, Zheng C, Yao W, Guo L, Peng H, Yang F, Wang M, Jin X: Predictors of difficult airway in a Chinese surgical population: the gender effect. *Minerva Anestesiologica* 2019; 85:478-86
558. Wang CMZ, Pang KP, Guan TS, Pang KA, Pang EB, Cherilynn TYN, Chan YH, Rotenberg BW: Predictors of difficulty in intubation in patients with obstructive sleep apnoea. *Med J Malaysia* 2019; 74:133-7
559. Watson NC, Hokanson M, Maltby JR, Todesco JM: The intubating laryngeal mask airway in failed fibreoptic intubation. *Can J Anaesth* 1999; 46:376-8
560. Watt S, Kalpan J, Kolli V: Case report of the use of videolaryngoscopy in thyroid goiter masses: An airway challenge. *Int J Surg Case Rep* 2016; 27:119-21
561. Wei W, Qiu HR, Wang HX, Xue FS: Anesthesia and airway managements for emergency removal of esophageal foreign body in a trisomy 21 patient with mental retardation and predicted difficult airway: A case report. *Medicine* 2020; 99:e23710
562. Weinberg S, Kravath R, Phillips L, Mendez H, Wolf GL: Episodic complete airway obstruction in children with undiagnosed obstructive sleep apnea. *Anesthesiology* 1984; 60:356-8
563. Williamson JA, Webb RK, Szekely S, Gillies ER, Dreosti AV: The Australian Incident Monitoring Study. Difficult intubation: An analysis of 2000 incident reports. *Anaesth Intensive Care* 1993; 21:602-7
564. Wilson ME, Spiegelhalter D, Robertson JA, Lesser P: Predicting difficult intubation. *Br J Anaesth* 1988; 61:211-216
565. Winterhalter M, Kirchhoff K, Groschel W, Lullwitz E, Heermann R, Hoy L, Heine J, Hagberg C, Piepenbrock S: The laryngeal tube for difficult airway management: a prospective investigation in patients with pharyngeal and laryngeal tumours. *Eur J Anaesthesiol* 2005; 22:678-82
566. Wojtczak JA: Submandibular sonography: assessment of hyomental distances and ratio, tongue size, and floor of the mouth musculature using portable sonography. *J Ultrasound Med* 2012; 31:523-8
567. Wolf LH, Gravenstein D: Capnography during fiberoptic bronchoscopy to verify tracheal intubation. *Anesth Analg* 1997; 85:701-3
568. Wong DT, Woo JA, Arora G: Lighted stylet-guided intubation via the intubating laryngeal airway in a patient with Hallermann-Streiff syndrome. *Can J Anaesth* 2009; 56:147-150
569. Wong P, Iqbal R, Light KP, Williams E, Hayward J: Head and neck surgery in a tertiary centre: Predictors of difficult airway and anaesthetic management. *Proceed Singapore Healthcare* 2016; 25:19-26
570. Wong SH, Hung CT: Prevalence and prediction of difficult intubation in Chinese women. *Anaesth Intensive Care* 1999; 27:49-52
571. Workeneh SA, Gebregzi AH, Denu ZA: Magnitude and predisposing factors of difficult airway during induction of general anaesthesia. *Anesthesiol Res Pract* 2017; 2017:5836397
572. Wu CN, Ma WH, Wei JQ, Wei HF, Cen QY, Cai QX, Cao Y: Laryngoscope and a new tracheal tube assist lightwand intubation in difficult airways due to unstable cervical spine. *PLoS One* 2015; 10:e0120231
573. Xu M, Li X, Wang J, Guo X: Application of a new combined model including radiological indicators to predict difficult airway in patients undergoing surgery for cervical spondylosis. *Chin Med J (Engl)* 2014; 127:4043-8
574. Xu M, Liu Y, Yang J, Liu H, Ding C: Ossification of the cervical anterior longitudinal ligament is an underdiagnosed cause of difficult airway: A case report and review of the literature. *BMC Anesthesiology* 2020; 20
575. Xue FS, Yang QY, Liao X, He N, Liu HP: Lightwand guided intubation in paediatric patients with a known difficult airway: a report of four cases. *Anaesthesia* 2008; 63:520-5

576. Yabuki S, Iwaoka S, Murakami M, Miura H: Reliability of the thyromental height test for prediction of difficult visualisation of the larynx: A prospective external validation. *Indian J Anaesth* 2019; 63:270-6
577. Yadav N, Rudingwa P, Mishra S, Pannerselvam S: Ultrasound measurement of anterior neck soft tissue and tongue thickness to predict difficult laryngoscopy - An observational analytical study. *Indian J Anaesth* 2019; 63:629-34
578. Yadav SS, Gupta S, Choudhary B: Smaller size laryngeal mask airway and gum elastic bougie combination is a failsafe technique for tracheal intubation in a child with temporomandibular joint ankylosis. *J Evol Med Dent Sci* 2014; 3:12225-9
579. Yakushiji H, Goto T, Shirasaka W, Hagiwara Y, Watase H, Okamoto H, Hasegawa K, Japanese Emergency Medicine Network i: Associations of obesity with tracheal intubation success on first attempt and adverse events in the emergency department: An analysis of the multicenter prospective observational study in Japan. *PLoS One* 2018; 13:e0195938
580. Yamagata K, Kawamura A, Kasai S, Akazawa M, Takeda M, Tachibana K: Anesthetic management of a child with Kagami-Ogata syndrome complicated with marked tracheal deviation: a case report. *JA Clin Rep* 2018; 4:62
581. Yaman F, Arslan B, Yuvañç E, Büyükkoçak U: Unexpected difficult airway with hypogonadotropic hypogonadism. *Int Med Case Rep J* 2014; 7:75-7
582. Yang D, Tong SY, Jin JH, Tang GZ, Sui JH, Wei LX, Deng XM: Shikani optical stylet-guided intubation via the intubating laryngeal airway in patients with scar contracture of the face and neck. *Chin Med Sci J* 2013; 28:195-200
583. Yao W, Wang B: Can tongue thickness measured by ultrasonography predict difficult tracheal intubation? *Br J Anaesth* 2017; 118:601-9
584. Yegian CC, Volz LM, Galgon RE: Use of an Airway exchange catheter-assisted extubation with continuous end-tidal carbon dioxide monitoring in a pediatric patient with a known difficult airway: A case report. *A A Prac* 2018; 11:233-5
585. Yıldırım İ, İnal MT, Memiş D, Turan FN: Determining the efficiency of different preoperative difficult intubation tests on patients undergoing caesarean section. *Balkan Med J* 2017; 34:436-43
586. Yilmaz C, Karasu D, Dilektasli E, Taha A, Ozgunay SE, Korfali G: An evaluation of ultrasound measurements of anterior neck soft tissue and other predictors of difficult laryngoscopy in morbidly obese patients. *Bariatr Surg Pract Patient Care* 2018; 13:18-24
587. Yoshikawa M, Shinomura T, Kishimoto K, Uga H: Anesthetic management of an adult patient with hyaline fibromatosis syndrome undergoing laparoscopic colectomy: A case report. *A A Practice* 2020; 14:87-9
588. Yoshimatsu Y, Morita R, Suginaka M, Furukawa K, Nakamura N, Yamairi K, Maruyama N, Kaji M, Kamimori T, Fujiwara H: Difficult intubation due to unknown congenital tracheal stenosis in the adult: A case report and literature review. *J Thorac Dis* 2018; 10:E93-7
589. Yu T, Wang B, Jin XJ, Wu RR, Wu H, He JJ, Yao WD, Li YH: Predicting difficult airways: 3-3-2 rule or 3-3 rule? *Ir J Med Sci* 2015; 184:677-83
590. Yun HJ, So E, Karm MH, Kim HJ, Seo KS: Orotracheal intubation in a patient with difficult airway by using fiberoptic nasotracheal intubation: A case report. *J Dent Anesth Pain Med* 2018; 18:125-8
591. Yunoki K, Miyawaki I, Yamazaki K, Mima H: Extracorporeal membrane oxygenation-assisted airway management for difficult airways. *J Cardiothorac Vasc Anesth* 2018; 32:2721-5
592. Zaballos M, Ginel MD, Portas M, Barrio M, López AM: Awake insertion of a laryngeal mask Airway-Proseal™ as alternative to awake fiberoptic intubation in management of anticipated difficult airway in ambulatory surgery. *Braz J Anesthesiol* 2016; 66:539-42
593. Zbeidy R, Torres Buendia N, Souki FG: Anaesthetic management of a parturient with spondylothoracic dysostosis. *BMJ Case Rep* 2020; 13:23

594. Zhang X, Li W: Misguidance of peroral rigid laryngoscopy in assessment of difficult airway: two comparable cases in microlaryngeal surgery. *BMJ Case Reports* 2013; 22:22
595. Zhou ZB, Yang XY, Zhou X, Wen SH, Xiao Y, Feng X: Anesthetic manipulation in extreme airway stenosis: a case report. *J Med Case Rep* 2014; 8:292
596. Zhu H, Liu J, Suo L, Zhou C, Sun Y, Jiang H: A randomized controlled comparison of non-channeled king vision, McGrath MAC video laryngoscope and Macintosh direct laryngoscope for nasotracheal intubation in patients with predicted difficult intubations. *BMC Anesthesiol* 2019; 19
597. Ziyaeifard M, Azarfarin R, Ferasatkish R, Dashti M: Management of difficult airway with laryngeal mask in a child with mucopolysaccharidosis and mitral regurgitation: A case report. *Res Cardiovascul Med* 2014; 3:e17456