

Supplemental Digital Content 6: Polyp detection by sedation type in full and restricted sample

Supplemental Table 6a: Full sample prevalence of polyp outcomes by sedation¹

	Propofol		Moderate Sedation	
	N	% (95% CI)	N	% (95% CI)
Any neoplastic lesion	16936	58.1 (57.6-58.7)	12765	51.8 (51.2-52.5)
Any adenoma	11335	39.5 (38.9-40.0)	9109	37.3 (36.7-38.0)
Any serrated polyp	9728	34.0 (33.4-34.5)	5912	24.5 (24.0-25.1)

¹ Sample includes all exams meeting inclusion criteria, regardless of endoscopist and facility average propofol use

Supplemental Table 6b: Restricted sample prevalence of polyp outcomes by sedation¹

	Propofol		Moderate Sedation	
	N	% (95% CI)	N	% (95% CI)
Any neoplastic lesion	2597	56.0 (54.6-57.5)	7784	54.6 (53.7-55.4)
Any adenoma	1841	40.2 (38.8-41.6)	5693	40.4 (39.6-41.2)
Any serrated polyp	1371	30.3 (28.9-31.6)	3589	25.7 (25.0-26.5)

¹ Sample restricted to exams performed by endoscopists with 5%-95% propofol use at facilities with 5%-95% propofol use