

**Supplemental Digital Content 7. Number of patients with systolic blood pressure below 110 mmHg during 24 h post-treatment in both groups.**

Number of patients with systolic blood pressure below 110 mmHg, n (%)	Intervention group	Control group	<i>P</i> value
1 h	6/161 (3.7)	4/166 (2.4)	0.711
2 h	13/152 (8.6)	8/163 (4.9)	0.195
3 h	14/151 (9.3)	9/160 (5.6)	0.219
4 h	17/151 (11.3)	11/160 (6.9)	0.177
5 h	12/151 (7.9)	12/159 (7.5)	0.895
6 h	14/152 (9.2)	10/157 (6.4)	0.351
7 h	10/149 (6.7)	13/153 (8.5)	0.559
8 h	9/149 (6.0)	19/153 (12.4)	0.056
9 h	14/148 (9.5)	18/154 (11.7)	0.529
10 h	8/149 (5.4)	14/154 (9.1)	0.212
11 h	10/145 (6.9)	12/146 (8.2)	0.670
12 h	12/149 (8.1)	10/154 (6.5)	0.601
13 h	11/148 (7.4)	7/150 (4.7)	0.316
14 h	14/149 (9.4)	9/150 (6.0)	0.271
15 h	15/146 (10.3)	6/150 (4.0)	0.036
16 h	14/145 (9.7)	6/150 (4.0)	0.053
17 h	12/141 (8.5)	8/149 (5.4)	0.291
18 h	7/141 (5.0)	6/149 (4.0)	0.700
19 h	12/138 (8.7)	6/146 (4.1)	0.113
20 h	13/136 (9.6)	8/142 (5.6)	0.216
21 h	6/133 (4.5)	8/140 (5.7)	0.652
22 h	15/134 (11.2)	5/138 (3.6)	0.017
23 h	12/130 (9.2)	3/137 (2.2)	0.013
24 h	8/132 (6.1)	10/135 (7.4)	0.661