

Appendix 1. Outcome Assessment Method Among Women Having Medication Abortion and Receiving Mifepristone at a Pharmacy (n=260)

	n	%
Participants who had follow-up with study clinic (n=235)		
Ultrasound	202	77.7%
Serial serum hCG testing	14	5.4%
Clinical history and home urine pregnancy test (telephone visit)	11	4.2%
Clinical history alone (telephone visit)	8	3.1%
Participants who did not follow-up with study clinic and indicated abortion completion in survey responses (n=25)		
Pregnancy symptoms went away*	16	6.2%
Normal period returned*	6	2.3%
Negative urine pregnancy test*	8	3.1%
Ultrasound at another facility	1	0.4%
Reported the clinic said abortion was complete	2	0.8%
Not reported	1	0.4%

*Some participants reported more than one response.

Appendix 2. Effectiveness of Medication Abortion by Weeks of Gestational Age

Gestational age at initial clinic visit (d)	Abortion outcome (n=260)					
	Complete abortion with medications only (n=243)		Incomplete abortion, had vacuum aspiration (n=17)		Ongoing pregnancy (n=0)	
	n	%	n	%	n	%
<=49	166	94.3%	10	5.7%	0	0.0%
50-56	40	93.0%	3	7.0%	0	0.0%
57-63	28	87.5%	4	12.5%	0	0.0%
64-70	9	100.0%	0	0.0%	0	0.0%

Grossman D, Baba CF, Kaller S, Biggs A, Raifman S, Gurazada T, et al. Medication abortion with pharmacist dispensing of mifepristone. *Obstet Gynecol* 2021;137.

The authors provided this information as a supplement to their article.

©2021 American College of Obstetricians and Gynecologists.