

Appendix 1. Baseline Demographics: ITT Population Compared With the 12-Month Completer Population

Demographic	Statistics	ITT population	Completed 12 month follow up	p-value	Completed 12 month follow up	ITT population	p-value
		Control n=165	Control n=151		Intervention n=135	Intervention n=169	
Age	Mean ±SD	51.3±12.7	51.6±12.7	0.89†	52.2 ±13.0	51.4 ± 13.3	0.6†
Race, n (%)	Asian	8 (2.4)	7 (4.6)	0.96‡	2 (1.5)	5 (1.5)	0.84‡
	Black	12 (3.9)	12 (8.0)		17 (12.6)	18 (5.4)	
	Middle Eastern/North African	0 (0.0)	0 (0.0)		1 (0.7)	0 (0.0)	
	Multi	2 (0.6)	2 (1.3)		5 (3.7)	8 (2.4)	
	White	138 (41.3)	127 (84.1)		105 (77.8)	130 (39.0)	
	Other*	4 (1.2)	3 (2.0)		4 (3.0)	6 (1.8)	
	Unknown*	0 (0.0)	0 (0.0)		1 (0.7)	2 (0.6)	
Ethnicity, n (%)	Hispanic/Latino	151 (45.2)	138 (91.4)	0.95‡	14 (10.4)	20 (6.0)	0.86‡
	Not Hispanic/Latino	14 (4.2)	13(8.6)		119 (88.2)	147 (44.0)	
	Declined to answer	0 (0.00%)	0 (0.0)		2 (1.5)	2 (0.6)	
BMI (kg/m ²)	Mean ±SD	31.7 ±7.5	31.9 ±7.5	0.78†	31.6 ±7.3	31.64 (7.4)	0.96†
Parity	Median (IQR)	2(1-4)	2 (1-4)		3 (2-3)	3(2-3)	
Mode of delivery, n (%)	Vaginal	80 (29.0)	71 (47.0)	0.9‡	73 (54.1)	88 (31.9)	0.9‡
	Forceps/Vacuum	34 (12.3)	33 (21.9)		29 (21.5)	32 (11.6)	
	Cesarean Section	20 (7.3)	20 (13.3)		16 (11.9)	22 (8.0%)	
Baseline UDI-6 Score	Mean ±SD	54.6±18.8	55.1±18.9	0.81†	52.2±20.0	53.7±19.9	0.56†

Abbreviations: IQR = Interquartile range; BMI = Body Mass Index

*Other/Unknown were prespecified categories from which participants could choose

†Student's t-test, ‡ ANOVA=One-way analysis of variance

Weinstein MM, Dunivan GC, Guaderrama NM, Richter HE. Digital therapeutic device for urinary incontinence: a longitudinal analysis at 6 and 12 Months. *Obstet Gynecol* 2023;141.

The authors provided this information as a supplement to their article.

©The Authors.