

Supplemental Table3 – Donor- and transplant-specific characteristics stratified by groups

Characteristics	<i>De Novo</i> n=12,295	Bridge-To-Transplant n=5,584	p-value
Donor-specific			
Age, years	30 [22, 42]	30 [22, 39]	0.029
Male sex	8,138 (66.2)	4,334 (77.6)	<0.001
Donor race			<0.001
Caucasian	7,796 (63.4)	3,696 (66.2)	
African American	1,943 (15.8)	951 (17.0)	
Hispanic	2,143 (17.4)	795 (14.2)	
Other	413 (3.4)	142 (2.5)	
Body mass index, kg/m ²	25.90 [22.91, 29.85]	26.78 [23.79, 30.74]	<0.001
Blood group			<0.001
A	4,603 (37.4)	1,874 (33.6)	
AB	385 (3.1)	74 (1.3)	
B	1,465 (11.9)	572 (10.2)	
O	5,842 (47.5)	3,064 (54.9)	
Tobacco use			0.026
Yes	1,634 (13.3)	669 (12.0)	
Missing	135 (1.1)	74 (1.3)	
Diabetes			0.449
Yes	437 (3.6)	179 (3.2)	
Missing	54 (0.4)	22 (0.4)	
Hypertension			0.719
Yes	1,892 (15.4)	833 (14.9)	
Missing	70 (0.6)	32 (0.6)	
Cytomegalovirus			0.003
Positive	7,629 (62.0)	3,317 (59.4)	
Missing	54 (0.4)	26 (0.5)	
BUN/creatinine ratio >= 30			0.407
Yes	1044 (8.5)	470 (8.4)	
Missing	24 (0.2)	6 (0.1)	
Transplant-specific			
Recipient/donor BMI ratio	1.00 [0.86, 1.16]	1.05 [0.90, 1.20]	<0.001
Race mismatch	5,994 (48.8)	2,698 (48.3)	0.601
ABO compatible	2,073 (16.9)	647 (11.6)	<0.001
Allocation type			<0.001
Local	7,379 (60.0)	3,598 (64.4)	
Regional	2,011 (16.4)	653 (11.7)	
National	2,888 (23.5)	1,327 (23.8)	
Foreign	17 (0.1)	6 (0.1)	
Distance, kilometers	149.67 [20.92, 481.19]	133.58 [16.09,	<0.001

Ischemic time, hours	3.13 [2.42, 3.78]	414.00] 3.20 [2.42, 3.85]	0.019
----------------------	-------------------	------------------------------	-------

Values presented as median [25th, 75th percentile] or count (percentage). All variables are complete unless stated otherwise. *BMI*, body mass index.