

**Supplemental Table6** – Incremental factors for conditional, risk-adjusted mortality at 1-year<sup>a</sup>

	Coefficients	HR	95% CI	p-value
Group				
<i>De novo</i>	Ref.	Ref.	Ref.	Ref.
<b>Bridge-to-transplant</b>	<b>0.228</b>	<b>1.256</b>	<b>(1.044-1.511)</b>	<b>0.016</b>
Recipient age, years	0.001	1.001	(0.992-1.009)	0.897
Diagnosis				
Ischemic	Ref.	Ref.	Ref.	Ref.
Dilated, non-ischemic	-0.042	0.959	(0.794-1.157)	0.661
Congenital	0.343	1.409	(0.863-2.300)	0.170
Restricted	0.592	1.808	(1.258-2.600)	0.001
Other	-0.042	0.959	(0.663-1.385)	0.822
Mechanical ventilation	0.527	1.694	(0.840-3.413)	0.141
Dialysis	0.567	1.762	(1.227-2.530)	0.002
Donor age, years	0.015	1.015	(1.008-1.023)	<0.001
Ischemic time, hours	0.017	1.017	(0.917-1.128)	0.746
Glomerular filtration rate, mL/min/1.73m <sup>2</sup>	-0.005	0.995	(0.991-0.998)	0.004
Total waitlist time, days	0.000	1.000	(1.000-1.000)	0.023
Race mismatch	0.197	1.218	(1.033-1.437)	0.019
Recipient body mass index, kg/m <sup>2</sup>	0.001	1.001	(0.980-1.022)	0.926
Distance, kilometers	0.000	1.000	(1.000-1.000)	0.847
Recipient-donor BMI ratio	0.089	1.093	(0.705-1.694)	0.692
Primary payment source				
Private	Ref.	Ref.	Ref.	Ref.
Medicare	0.211	1.236	(1.027-1.487)	0.025
Medicaid	0.382	1.466	(1.133-1.897)	0.004
Other public	0.145	1.156	(0.686-1.949)	0.586
Other	-0.549	0.578	(0.143-2.326)	0.440

<sup>a</sup>Conditioned on survival to 90 days after transplant.

*BMI*, body mass index; *CI*, confidence interval; *HR*, hazards ratio.