

Jonathan P. Troost¹, Howard Trachtman², Patrick H. Nachman³, Matthias Kretzler⁴, Cathie Spino⁵, Radko Komers⁶, Sarah Tuller⁶, Kalyani Perumal⁷, Susan F. Massengill⁸, Elaine S. Kamil⁹, Gia Oh¹⁰, David T. Selewski¹, Patrick Gipson¹, Debbie S. Gipson¹

1. Division of Nephrology, Department of Pediatrics and Communicable Diseases, C.S. Mott Children's Hospital, University of Michigan, Ann Arbor, MI
2. Division of Nephrology, Department of Pediatrics, New York University Langone Medical Center, New York, NY
3. Department of Pathology and Laboratory Medicine, Department of Medicine, and Kidney Center, School of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina
4. Division of Nephrology, Department of Internal Medicine, University of Michigan, Ann Arbor, MI
5. Department of Biostatistics, School of Public Health, University of Michigan, Ann Arbor, MI
6. Retrophin Inc., San Diego, CA
7. Division of Nephrology, Department of Internal Medicine, John H. Stroger, Jr. Hospital, Chicago, IL
8. Division of Pediatric Nephrology, Levine Children's Hospital, Charlotte, NC
9. Division of Pediatric Nephrology, Department of Pediatrics, Cedars-Sinai Medical Center, David Geffen School of Medicine at The University of California, Los Angeles, Los Angeles, CA
10. Division of Nephrology, Department of Pediatrics, Stanford University School of Medicine, Stanford, CA