

## Supplemental Material Table of Contents

*Supplemental Table 1.* Cox proportional hazards model in discontinuation population.

*Supplemental Figure 1.* Flow chart of inclusion of patients in study

*Supplemental Table 1.* Cox proportional hazards model in discontinuation population.

<b>Variable</b>	<b>Hazard ratio<sup>^</sup></b>	<b>95% confidence interval</b>	<b>P value</b>
Systolic blood pressure change, mmHg	0.99	0.95 – 1.04	0.76
Log-transformed albuminuria change, %	2.15	1.10 – 4.22	0.02
Female, n	1.00	0.41 – 2.40	0.99
Age, years	0.92	0.76 – 1.08	0.27
eGFR, ml/min/1.73m <sup>2</sup> †	0.92	0.86 – 0.98	0.009
Albuminuria, 100 mg/g†	1.11	1.04 – 1.18	0.001
Systolic blood pressure, mmHg†	1.00	0.97 – 1.05	0.63

<sup>^</sup>composite endpoint: a sustained 50% reduction in eGFR or progression to end-stage kidney disease (eGFR < 10 ml/min/1.73m<sup>2</sup> or start of kidney replacement therapy)

†measurement at last visit before discontinuation

Supplemental Figure 1. Flow chart of inclusion of patients in study

