Supplemental Figure 1. Bivariate relationship of markers of mineral metabolism in patients with ESRD requiring dialysis with interaction by race. (A) relationship between 1,25$(OH)_2$D and 25(OH)D; (B) iPTH and 25(OH)D; (C) iPTH and 1,25$(OH)_2$D; (D) 1,25$(OH)_2$D and FGF-23; (E) iPTH and FGF-23


Race
Black
White
Log 1,25(OH)2D (pg/mL)
0.6
0.8
1.0
1.2
1.4
1.6
1.8
2.0
Interaction P−value: 0.8251
1.0
1.5
2.0
2.5
Log 25(OH)D (ng/mL)


Race
Black
White
Log iPTH (pg/mL)
3.0
2.5
2.0
1.5
1.0
Interaction P−value: 0.3744
1.0
1.5
2.0
2.5
Log 25(OH)D (ng/mL)


Race
Black
White
Log iPTH (pg/mL)
3.0
2.5
2.0
1.5
1.0
Interaction P−value: 0.7433
0.6
0.8
1.0
1.2
1.4
1.6
1.8
2.0
Log 1,25(OH)2D (pg/mL)


Race
Black
White
Interaction P−value: 0.0917
Log 1,25(OH)2D (pg/mL)
0.6
0.8
1.0
1.2
1.4
1.6
1.8
2.0
2.0
2.5
3.0
3.5
4.0
4.5
Log FGF−23 (RU/mL)


Race
Black
White
Interaction P−value: 0.3750
Log iPTH (pg/mL)
1.0
1.5
2.0
2.5
3.0
2.0
2.5
3.0
3.5
4.0
4.5
Log FGF−23 (RU/mL)