

## Sudden Cardiac Arrest (SCA) Prevention Treatment Algorithms

- Inpatient to Outpatient Transition
- ACE Inhibitors and/or ARBs
- Beta Blockers
- Aldosterone Antagonists
- Implantable Cardioverter Defibrillator Therapy (ICD Inpatient)
- Cardiac Resynchronization Therapy (CRT Inpatient)
- Anticoagulation Therapy in Patients with Atrial Fibrillation (Outpatient)
- Management of Volume Overload (Outpatient)
- Implantable Cardioverter Defibrillator (ICD Outpatient)
- Cardiac Resynchronization Therapy (CRT Outpatient)
- Device Therapy

## Sudden Cardiac Arrest (SCA) Prevention Pathways and Tools Objectives

- **Facilitate** optimal care for post-MI and HF patients at risk for SCA
- **Educate** healthcare providers and patients about SCA and treatment options and increase awareness and patient access to diagnostics and lifesaving therapies
- **Promote** evidence-based, guideline-recommended medical and device therapy and increase guideline awareness and adoption among healthcare providers
- **Assist** hospitals and practices in closing treatment gaps by providing practical information, disease management, and communication tools to identify and treat patients at risk for SCA


### SCA Prevention Medical Advisory Team:

Gregg Fonarow, MD  
Nancy Albert, PhD, RN  
David Cannom, MD  
William Lewis, MD  
Julie Shea, MS, RNCS  
Mary Norine Walsh, MD

This clinical tool is not intended to replace individual medical judgment or individual patient needs.

Sponsored by Medtronic, Inc.  
April 2007  
Revised September 2008

## Angiotensin-Converting Enzyme Inhibitor (ACEI)/ Angiotensin Receptor Blocker (ARB) Inpatient/Outpatient Treatment Algorithm



**Note:** Careful consideration of patient characteristics and choice of drugs is warranted.

\* For hospitalized patients previously treated with ACEI/ARB at the time of admission, therapy should be continued in the absence of contraindications.

	ACEIs							ARBs <sup>†</sup>	
	Captopril	Enalapril	Lisinopril	Ramipril	Trandolapril	Quinapril	Fosinopril	Candesartan	Valsartan
<b>Initial dose</b>	6.25 mg tid	2.5 mg bid	2.5-5 mg qd	2.5 mg qd	1 mg qd	10 mg qd	10 mg qd	4-8 mg qd	40 mg bid
<b>Titration steps</b> (typically double dose at each step)	12.5 mg tid 25 mg tid	5 mg bid	5 mg qd 10 mg qd	5 mg qd	2 mg qd	20 mg qd 40 mg qd	20 mg qd 40 mg qd	16 mg qd	80 mg bid
<b>Target dose</b>	50 mg tid	10 mg bid	20 mg qd	10 mg qd	4 mg qd	80 mg qd	80 mg qd	32 mg qd	160 mg bid

<sup>†</sup> ARBs are most frequently used in place of ACEIs when side effects limit ACEI use (intolerable cough may occur in as many as 20% of patients receiving ACEIs).

Routine combined use of an ACEI, ARB, and aldosterone antagonist is not recommended for patients with current or prior symptoms of HF and reduced LVEF.

### Reference Sources


- Fonarow GC, Yancy CW, Albert NM, et al. Improving the use of evidence-based heart failure therapies in the outpatient setting: the IMPROVE HF Registry Rationale, Design, and Implementation. *Circulation*. May 30, 2006;113(21):e789 (Abstract #P15).
- Garg R, Yusuf S. Overview of randomized trials of angiotensin-converting enzyme inhibitors on mortality and morbidity in patients with heart failure. Collaborative Group on ACE Inhibitor Trials [published correction appears in *JAMA*. 1995;274:462]. *JAMA*. May 10, 1995;273(18):1450-1456.
- Hunt SA, Abraham WT, Chin MH, et al. ACC/AHA 2005 Guideline Update for the Diagnosis and Management of Chronic Heart Failure in the Adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure). *Circulation*. September 20, 2005;112(12):e154-235.
- Pfeffer MA, McMurray JJ, Velazquez EJ, et al, for the Valsartan in Acute Myocardial Infarction Trial Investigators. Valsartan, captopril, or both in myocardial infarction complicated by heart failure, left ventricular dysfunction, or both [published correction appears in *N Engl J Med*. 2004;350:203]. *N Engl J Med*. November 13, 2003;349(20):1893-1906.
- Young JB, Dunlap ME, Pfeffer MA, et al. Mortality and morbidity reduction with Candesartan in patients with chronic heart failure and left ventricular systolic dysfunction: results of the CHARM low-left ventricular ejection fraction trials. *Circulation*. October 26, 2004; 110(17):2618-2626.

Adapted, with permission, by the SCA Prevention Medical Advisory Team, from the IMPROVE HF toolkit.  
This is a general algorithm to assist in the management of patients. This clinical tool is not intended to replace individual medical judgment or individual patient needs.

Refer to the manufacturers' prescribing information and/or instructions for use for the indications, contraindications, warnings, and precautions associated with the medications and devices referenced in these materials.

Sponsored by Medtronic, Inc.  
April 2007

## Beta Blocker Inpatient/Outpatient Treatment Algorithm



## References

- <sup>1</sup> Beta blocker therapy initiation is recommended for patients with a recent decompensation of HF after optimization of volume status and successful discontinuation of IV diuretics and vasoactive agents, including inotropic support. Whenever possible, beta blockade should be initiated in the hospital setting at a low dose in stable patients prior to discharge.
- <sup>2</sup> Patients hospitalized with decompensated HF already treated with beta blocker therapy prior to hospitalization should continue on beta blocker therapy as long as they do not have any contraindications, are not in cardiogenic shock, and do not show signs of systemic hypoperfusion (altered mental status, narrow pulse pressure, cold or clammy skin, rising BUN/serum Cr). A temporary reduction of dose in this setting may be considered. Abrupt discontinuation should be avoided, if possible. If discontinued or reduced, beta blockers should be reinstated or the dose should be gradually increased before the patient is discharged.
- <sup>3</sup> ACC/AHA 2005 guidelines recommend using only those beta blockers proven to be effective in heart failure (carvedilol, sustained-release metoprolol succinate, and bisoprolol) at the doses studied in large clinical trials. If patient is currently on a beta blocker other than those listed, consider switching.
- <sup>4</sup> Applies to both outpatients and those hospitalized for heart failure.
- <sup>5</sup> Beta blocker titration steps are generally at 2-week intervals. BP and HR should be carefully monitored. If SBP is < 80 mmHg or HR is < 55 bpm, assess the dose and recheck patient carefully for signs of hypoperfusion. Recheck status as needed.
- <sup>6</sup> Patients who cannot achieve target dose of the beta blocker should be maintained on the highest tolerated dose.
- <sup>7</sup> For patients weighing > 85 kg, carvedilol 50 mg bid may be used.


## Reference Sources

- Fonarow GC, Yancy CW, Albert NM, et al. Improving the use of evidence-based heart failure therapies in the outpatient setting: the IMPROVE HF Registry Rationale, Design, and Implementation. *Circulation*. May 30, 2006;113(21):e789 (Abstract #P15).
- HFSA 2006 Comprehensive Heart Failure Practice Guideline. *J Card Fail*. February 2006;12(1):e1-2.
- Hjalmarson A, Goldstein S, Fagerberg B, et al. Effects of controlled-release metoprolol on total mortality, hospitalizations, and well-being in patients with heart failure: the Metoprolol CR/XL Randomized Intervention Trial in congestive heart failure (MERIT-HF) MERIT-HF Study Group. *JAMA*. March 8, 2000;283(10):1295-1302.
- Hunt SA, Abraham WT, Chin MH, et al. ACC/AHA 2005 Guideline Update for the Diagnosis and Management of Chronic Heart Failure in the Adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure). *Circulation*. September 20, 2005;112(12):e154-235.
- Packer M, Coats AJ, Fowler MB, et al, for the Carvedilol Prospective Randomized Cumulative Survival Study Group. Effect of carvedilol on survival in severe chronic heart failure. *N Engl J Med*. May 31, 2001;344(22):1651-1658.

Adapted, with permission, by the SCA Prevention Medical Advisory Team, from the IMPROVE HF toolkit.  
This is a general algorithm to assist in the management of patients. This clinical tool is not intended to replace individual medical judgment or individual patient needs.  
Refer to the manufacturers' prescribing information and/or instructions for use for the indications, contraindications, warnings, and precautions associated with the medications and devices referenced in these materials.

Sponsored by Medtronic, Inc.  
April 2007

## Aldosterone Antagonist Inpatient/Outpatient Treatment Algorithm



Routine combined use of an ACEI, ARB, and aldosterone antagonist is not recommended for patients with current or prior symptoms of heart failure and reduced LVEF.

\* Switch to eplerenone if signs of gynecomastia.

† Start at 6.25 mg qd in patients at increased risk for hyperkalemia.

- Impaired renal function is a risk factor for hyperkalemia during treatment with aldosterone antagonists. The risk of hyperkalemia increases progressively when serum Cr > 1.6 mg/dL.
- Hyperkalemia risk is increased with concomitant use of higher doses of ACEIs or ARBs
- NSAIDs and COX-2 inhibitors should be avoided
- Potassium supplements should be discontinued or reduced
- Dehydration, by diarrhea or other causes, should be addressed emergently

Ongoing clinical trials are evaluating the benefit of aldosterone antagonists in patients with mild heart failure.

### Reference Sources

- Fonarow GC, Yancy CW, Albert NM, et al. Improving the use of evidence-based heart failure therapies in the outpatient setting: the IMPROVE HF Registry Rationale, Design, and Implementation. *Circulation*. May 30, 2006;113(21):e789 (Abstract #P15).
- Hunt SA, Abraham WT, Chin MH, et al. ACC/AHA 2005 Guideline Update for the Diagnosis and Management of Chronic Heart Failure in the Adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure). *Circulation*. September 20, 2005;112(12):e154-235.
- Pitt B, Remme W, Zannad F, et al, for the Eplerenone Post-Acute Myocardial Infarction Heart Failure Efficacy and Survival Study Investigators. Eplerenone, a selective aldosterone blocker, in patients with left ventricular dysfunction after myocardial infarction [published correction appears in *N Engl J Med*. 2003;348:2271]. *N Engl J Med*. April 3, 2003;348(14):1309-1321.
- Pitt B, Zannad F, Remme WJ, et al. The effect of spironolactone on morbidity and mortality in patients with severe heart failure. Randomized Aldactone Evaluation Study Investigators. *N Engl J Med*. September 2, 1999;341(10):709-717.
- Swedberg K, Cleland J, Dargie H, et al. Guidelines for the diagnosis and treatment of chronic heart failure: executive summary (update 2005): The Task Force for the Diagnosis and Treatment of Chronic Heart Failure of the European Society of Cardiology. *Eur Heart J*. June 2005;26(11):1115-1140.


Adapted, with permission, by the SCA Prevention Medical Advisory Team, from the IMPROVE HF registry toolkit.

This is a general algorithm to assist in the management of patients. This clinical tool is not intended to replace individual medical judgment or individual patient needs.

Refer to the manufacturers' prescribing information and/or instructions for use for the indications, contraindications, warnings, and precautions associated with the medications and devices referenced in these materials.

Sponsored by Medtronic, Inc.  
April 2007

## Implantable Cardioverter Defibrillator (ICD) Therapy Inpatient Algorithm



\* Class I recommendation.

<sup>†</sup> Functional status as documented prior to current hospitalization while on chronic optimal medical therapy.

<sup>††</sup> CMS coverage for primary prevention ICD implants: patients with nonischemic dilated cardiomyopathy (NIDCM) > 9 months, NYHA Class II and III heart failure, and measured LVEF ≤ 35%. Patients with NIDCM > 3 months and < 9 months, NYHA Class II or III heart failure, and measured LVEF ≤ 35% at this time are only covered by Medicare if these patients are enrolled in an FDA-approved category B IDE clinical trial, a trial under the CMS clinical trial policy, or the American College of Cardiology National Cardiovascular Data Registry (ACC-NCDR).



## Reference Sources

- Bardy GH, Lee KL, Mark DB, et al, for the Sudden Cardiac Death in Heart Failure Trial (SCD-HeFT) Investigators. Amiodarone or an implantable cardioverter-defibrillator for congestive heart failure [published correction appears in *N Engl J Med.* 2005;352:2146]. *N Engl J Med.* January 20, 2005;352(3):225-237.
- Buxton AE, Lee KL, Fisher JD, Josephson ME, Prystowsky EN, Hafley G. A randomized study of the prevention of sudden death in patients with coronary artery disease. Multicenter Unsustained Tachycardia Trial Investigators [published correction appears in *N Engl J Med.* 2000;342:1300]. *N Engl J Med.* December 16, 1999;341(25):1882-1890.
- Epstein AE, Dimarco JP, Ellenbogen KA, et al. ACC/AHA/HRS 2008 Guidelines for device-based therapy of cardiac rhythm abnormalities. *Heart Rhythm.* June 2008;5(6):e1-62.
- Fonarow GC, Yancy CW, Albert NM, et al. Improving the use of evidence-based heart failure therapies in the outpatient setting: the IMPROVE HF Registry Rationale, Design, and Implementation. *Circulation.* May 30, 2006;113(21):e789 (Abstract #P15).
- Gregoratos G, Abrams J, Epstein AE, et al. ACC/AHA/NASPE 2002 guideline update for implantation of cardiac pacemakers and antiarrhythmia devices: summary article: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (ACC/AHA/NASPE Committee to Update the 1998 Pacemaker Guidelines). *Circulation.* October 15, 2002; 106(16):2145-2161.
- Hohnloser SH, Kuck KH, Dorian P, et al, for the DINAMIT Investigators. Prophylactic use of an implantable cardioverter-defibrillator after acute myocardial infarction. *N Engl J Med.* December 9, 2004;351(24):2481-2488.
- Hunt SA, Abraham WT, Chin MH, et al. ACC/AHA 2005 Guideline Update for the Diagnosis and Management of Chronic Heart Failure in the Adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure). *Circulation.* September 20, 2005;112(12):e154-235.
- Moss AJ, Hall WJ, Cannom DS, et al. Improved survival with an implanted defibrillator in patients with coronary disease at high risk for ventricular arrhythmia. Multicenter Automatic Defibrillator Implantation Trial Investigators. *N Engl J Med.* December 26, 1996; 335(26):1933-1940.
- Moss AJ, Zareba W, Hall WJ, et al, for the Multicenter Automatic Defibrillator Implantation Trial II Investigators. Prophylactic implantation of a defibrillator in patients with myocardial infarction and reduced ejection fraction. *N Engl J Med.* March 21, 2002;346(12):877-883.

Adapted, with permission, by the SCA Prevention Medical Advisory Team, from the IMPROVE HF registry toolkit.

This is a general algorithm to assist in the management of patients. This clinical tool is not intended to replace individual medical judgment or individual patient needs.


Refer to the manufacturers' prescribing information and/or instructions for use for the indications, contraindications, warnings, and precautions associated with the medications and devices referenced in these materials.

Sponsored by Medtronic, Inc.

April 2007

Revised September 2008

## Cardiac Resynchronization Therapy (CRT) Inpatient Algorithm



**Note:** Ongoing clinical trials are evaluating the benefit of CRT in NYHA Class II patients.

\* Functional status as documented prior to current hospitalization while on chronic optimal medical therapy.

† CRT is a Class I guideline recommendation for patients with normal sinus rhythm (NSR). CRT is a Class IIa recommendation for patients with atrial fibrillation. Preliminary data suggest a possible functional improvement in these patients.

†† Inclusion of defibrillator to be based on ICD algorithm and physician discretion.

## Reference Sources

- Abraham WT. Rationale and design of a randomized clinical trial to assess the safety and efficacy of cardiac resynchronization therapy in patients with advanced heart failure: the Multicenter InSync Randomized Clinical Evaluation (MIRACLE). *J Card Fail*. December 2000; 6(4):369-380.
- Bristow MR, Feldman AM, Saxon LA. Heart failure management using implantable devices for ventricular resynchronization: Comparison of Medical Therapy, Pacing, and Defibrillation in Chronic Heart Failure (COMPANION) Trial. COMPANION Steering Committee and COMPANION Clinical Investigators. *J Card Fail*. September 2000;6(3):276-285.
- Cleland JG, Daubert JC, Erdmann E, et al, for the Cardiac Resynchronization-Heart Failure (CARE-HF) Study Investigators. The effect of cardiac resynchronization on morbidity and mortality in heart failure. *N Engl J Med*. April 14, 2005;352(15):1539-1549.
- Epstein AE, Dimarco JP, Ellenbogen KA, et al. ACC/AHA/HRS 2008 Guidelines for device-based therapy of cardiac rhythm abnormalities. *Heart Rhythm*. June 2008;5(6):e1-62.
- Fonarow GC, Yancy CW, Albert NM, et al. Improving the use of evidence-based heart failure therapies in the outpatient setting: the IMPROVE HF Registry Rationale, Design, and Implementation. *Circulation*. May 30, 2006;113(21):e789 (Abstract #P15).
- Hunt SA, Abraham WT, Chin MH, et al. ACC/AHA 2005 Guideline Update for the Diagnosis and Management of Chronic Heart Failure in the Adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure). *Circulation*. September 20, 2005;112(12):e154-235.
- Strickberger SA, Conti J, Daoud EG, et al. Patient selection for cardiac resynchronization therapy: from the Council on Clinical Cardiology Subcommittee on Electrocardiography and Arrhythmias and the Quality of Care and Outcomes Research Interdisciplinary Working Group, in collaboration with the Heart Rhythm Society. *Circulation*. April 26, 2005;111(16):2146-2150.
- Young JB, Abraham WT, Smith AL, et al, for the Multicenter InSync ICD Randomized Clinical Evaluation (MIRACLE ICD) Trial Investigators. Combined cardiac resynchronization and implantable cardioversion defibrillation in advanced chronic heart failure: the MIRACLE ICD Trial. *JAMA*. May 28, 2003;289(20):2685-2694.

Adapted, with permission, by the SCA Prevention Medical Advisory Team, from the IMPROVE HF registry toolkit.

This is a general algorithm to assist in the management of patients. This clinical tool is not intended to replace individual medical judgment or individual patient needs.


Refer to the manufacturers' prescribing information and/or instructions for use for the indications, contraindications, warnings, and precautions associated with the medications and devices referenced in these materials.

Sponsored by Medtronic, Inc.

April 2007

Revised September 2008

## Anticoagulation Therapy in Atrial Fibrillation Outpatient Algorithm



\*Atrial flutter should be similarly treated.

• Contraindications to warfarin include:

- Allergy
- Pregnancy
- Risk of bleeding (such as active peptic ulcer disease); hemorrhagic stroke; other hemorrhage; hepatic failure; bleeding disorder; metastatic cancer; recent or planned surgery or biopsy procedure; other physician-documented bleeding risk
- High risk of fall documented by physician
- Psychosocial concerns (such as active psychosis; terminal illness/comfort care only; alcoholism or drug abuse)
- Other potential contraindication (such as seizure disorder; malignant hypertension; intracranial aneurysm, repaired or unrepaired)

• Aspirin should be used in patients with absolute contraindications to oral anticoagulation

### Reference Sources


- Bonow RO, Bennett S, Casey DE Jr, et al. ACC/AHA clinical performance measures for adults with chronic heart failure: a report of the American College of Cardiology/American Heart Association Task Force on Performance Measures (Writing Committee to Develop Heart Failure Clinical Performance Measures) endorsed by the Heart Failure Society of America. *J Am Coll Cardiol*. September 20, 2005;46(6):1144-1178.
- Fonarow GC. Quality indicators for the management of heart failure in vulnerable elders. *Ann Intern Med*. October 16, 2001;135(8 Pt 2):694-702.
- Fonarow GC, Yancy CW, Albert NM, et al. Improving the use of evidence-based heart failure therapies in the outpatient setting: the IMPROVE HF Registry Rationale, Design, and Implementation. *Circulation*. May 30, 2006;113(21):e789 (Abstract #P15).
- Fuster V, Rydén LE, Cannom DS, et al. ACC/AHA/ESC 2006 guidelines for the management of patients with atrial fibrillation – executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the European Society of Cardiology Committee for Practice Guidelines (Writing Committee to Revise the 2001 Guidelines for the Management of Patients With Atrial Fibrillation) [published correction appears in *J Am Coll Cardiol*. 2007;50:562]. *J Am Coll Cardiol*. August 15, 2006;48(4):854-906.
- Singer DE, Albers GW, Dalen JE, Go AS, Halperin JL, Manning WJ. Antithrombotic therapy in atrial fibrillation: the Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy. *Chest*. September 2004;126(3 Suppl):429S-456S.

Adapted, with permission, by the SCA Prevention Medical Advisory Team, from the IMPROVE HF registry toolkit.  
This is a general algorithm to assist in the management of patients. This clinical tool is not intended to replace individual medical judgment or individual patient needs.

Refer to the manufacturers' prescribing information and/or instructions for use for the indications, contraindications, warnings, and precautions associated with the medications and devices referenced in these materials.

Sponsored by Medtronic, Inc.  
April 2007

## Management of Volume Overload Outpatient Algorithm



### Diuretic Maintenance Dosing

Weight returned to baseline (identifiable cause for weight increase, e.g., nonadherence)	Resume original dose
Weight returned to baseline, but patient failed original dose previously, or no known cause for weight gain	Continue at current increased dose
Weight returned to baseline, but required two or more diuretic titrations	Resume dose prior to last increase (i.e., down one titration level)
Symptoms improved, but weight has not returned to baseline	Continue at current increased dose
Persistent symptoms with no change in weight	Continue next titration level
Persistent or worsening symptoms, and/or increase in weight, and/or history of frequent hospitalizations for volume overload	Consider adding metolazone, IV diuretic, or hospitalization. PO metolazone may be added in resistant cases for no more than 3 days, then reassess. <sup>4</sup>

<sup>1</sup> Fluid restriction (< 2 L/day) is recommended in patients with moderate hyponatremia (serum sodium < 130 mEq/L) and should be considered to assist in treatment of fluid overload in other patients.

<sup>2</sup> Consider also loosening the degree of dietary sodium restriction, then reassess.

<sup>3</sup> Initial dose of loop diuretic at physician discretion. Careful observation for the development of side effects, including electrolyte abnormalities, symptomatic hypotension, and renal dysfunction, is recommended in patients treated with diuretics. Patients should undergo routine laboratory studies and clinical examination as dictated by their clinical response.

<sup>4</sup> Addition of chlorothiazides or metolazone, once or twice daily, to loop diuretics should be considered in patients with persistent fluid retention despite high-dose loop diuretic therapy. But chronic daily use, especially of metolazone, should be avoided if possible because of the potential for electrolyte shifts and volume depletion. These drugs may be used periodically (every other day or weekly) to optimize fluid management. Volume status and electrolytes must be monitored closely when multiple diuretics are used. Consider administering metolazone 30-60 minutes before administration of loop diuretic.

#### Reference Sources


- Cody RJ, Covit AB, Schaer GL, Laragh JH, Sealey JE, Feldschuh J. Sodium and water balance in chronic congestive heart failure. *J Clin Invest.* May 1986;77(5):1441-1452.
- Fonarow GC, Yancy CW, Albert NM, et al. Improving the use of evidence-based heart failure therapies in the outpatient setting: the IMPROVE HF Registry Rationale, Design, and Implementation. *Circulation.* May 30, 2006;113(21):e789 (Abstract #P15).
- HFSA 2006 Comprehensive Heart Failure Practice Guideline. *J Card Fail.* February 2006;12(1):e1-2.
- Hunt SA, Abraham WT, Chin MH, et al. ACC/AHA 2005 Guideline Update for the Diagnosis and Management of Chronic Heart Failure in the Adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure). *Circulation.* September 20, 2005; 112(12):e154-235.

Adapted, with permission, by the SCA Prevention Medical Advisory Team, from the IMPROVE HF registry toolkit.  
This is a general algorithm to assist in the management of patients. This clinical tool is not intended to replace individual medical judgment or individual patient needs.

Refer to the manufacturers' prescribing information and/or instructions for use for the indications, contraindications, warnings, and precautions associated with the medications and devices referenced in these materials.

Sponsored by Medtronic, Inc.  
April 2007

## Implantable Cardioverter Defibrillator (ICD) Therapy Outpatient Algorithm



\* Class I recommendation.

† CMS coverage for primary prevention ICD implants: patients with nonischemic dilated cardiomyopathy (NIDCM) > 9 months, NYHA Class II and III heart failure, and measured LVEF ≤ 35%. Patients with NIDCM > 3 months and < 9 months, NYHA Class II or III heart failure, and measured LVEF ≤ 35% at this time are only covered by Medicare if these patients are enrolled in an FDA-approved category B IDE clinical trial, a trial under the CMS clinical trial policy, or the American College of Cardiology National Cardiovascular Data Registry (ACC-NCDR).



### Reference Sources

- Bardy GH, Lee KL, Mark DB, et al, for the Sudden Cardiac Death in Heart Failure Trial (SCD-HeFT) Investigators. Amiodarone or an implantable cardioverter-defibrillator for congestive heart failure [published correction appears in *N Engl J Med*. 2005;352:2146]. *N Engl J Med*. January 20, 2005;352(3):225-237.
- Buxton AE, Lee KL, Fisher JD, Josephson ME, Prystowsky EN, Hafley G. A randomized study of the prevention of sudden death in patients with coronary artery disease. Multicenter Unsustained Tachycardia Trial Investigators [published correction appears in *N Engl J Med*. 2000;342:1300]. *N Engl J Med*. December 16, 1999;341(25):1882-1890.
- Epstein AE, Dimarco JP, Ellenbogen KA, et al. ACC/AHA/HRS 2008 Guidelines for device-based therapy of cardiac rhythm abnormalities. *Heart Rhythm*. June 2008;5(6):e1-62.
- Fonarow GC, Yancy CW, Albert NM, et al. Improving the use of evidence-based heart failure therapies in the outpatient setting: the IMPROVE HF Registry Rationale, Design, and Implementation. *Circulation*. May 30, 2006;113(21):e789 (Abstract #P15).
- Gregoratos G, Abrams J, Epstein AE, et al. ACC/AHA/NASPE 2002 guideline update for implantation of cardiac pacemakers and antiarrhythmia devices: summary article: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (ACC/AHA/NASPE Committee to Update the 1998 Pacemaker Guidelines). *Circulation*. October 15, 2002; 106(16):2145-2161.
- Hohnloser SH, Kuck KH, Dorian P, et al, for the DINAMIT Investigators. Prophylactic use of an implantable cardioverter-defibrillator after acute myocardial infarction. *N Engl J Med*. December 9, 2004;351(24):2481-2488.
- Hunt SA, Abraham WT, Chin MH, et al. ACC/AHA 2005 Guideline Update for the Diagnosis and Management of Chronic Heart Failure in the Adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure). *Circulation*. September 20, 2005; 112(12):e154-235.
- Moss AJ, Hall WJ, Cannom DS, et al. Improved survival with an implanted defibrillator in patients with coronary disease at high risk for ventricular arrhythmia. Multicenter Automatic Defibrillator Implantation Trial Investigators. *N Engl J Med*. December 26, 1996; 335(26):1933-1940.
- Moss AJ, Zareba W, Hall WJ, et al, for the Multicenter Automatic Defibrillator Implantation Trial II Investigators. Prophylactic implantation of a defibrillator in patients with myocardial infarction and reduced ejection fraction. *N Engl J Med*. March 21, 2002;346(12):877-883.


Adapted, with permission, by the SCA Prevention Medical Advisory Team, from the IMPROVE HF registry toolkit.

This is a general algorithm to assist in the management of patients. This clinical tool is not intended to replace individual medical judgment or individual patient needs.

Refer to the manufacturers' prescribing information and/or instructions for use for the indications, contraindications, warnings, and precautions associated with the medications and devices referenced in these materials.

Sponsored by Medtronic, Inc.  
April 2007  
Revised September 2008

## Cardiac Resynchronization Therapy (CRT) Outpatient Algorithm



**Note:** Ongoing clinical trials are evaluating the benefit of CRT in NYHA Class II patients.

\* CRT is a Class I guideline recommendation for patients with normal sinus rhythm (NSR). CRT is a Class IIa recommendation for patients with atrial fibrillation. Preliminary data suggest a possible functional improvement in these patients.

† Inclusion of defibrillator to be based on ICD algorithm and physician discretion.

## Reference Sources

- Abraham WT. Rationale and design of a randomized clinical trial to assess the safety and efficacy of cardiac resynchronization therapy in patients with advanced heart failure: the Multicenter InSync Randomized Clinical Evaluation (MIRACLE). *J Card Fail*. December 2000; 6(4):369-380.
- Bristow MR, Feldman AM, Saxon LA. Heart failure management using implantable devices for ventricular resynchronization: Comparison of Medical Therapy, Pacing, and Defibrillation in Chronic Heart Failure (COMPANION) Trial. COMPANION Steering Committee and COMPANION Clinical Investigators. *J Card Fail*. September 2000;6(3):276-285.
- Cleland JG, Daubert JC, Erdmann E, et al, for the Cardiac Resynchronization-Heart Failure (CARE-HF) Study Investigators. The effect of cardiac resynchronization on morbidity and mortality in heart failure. *N Engl J Med*. April 14, 2005;352(15):1539-1549.
- Epstein AE, Dimarco JP, Ellenbogen KA, et al. ACC/AHA/HRS 2008 Guidelines for device-based therapy of cardiac rhythm abnormalities. *Heart Rhythm*. June 2008;5(6):e1-62.
- Fonarow GC, Yancy CW, Albert NM, et al. Improving the use of evidence-based heart failure therapies in the outpatient setting: the IMPROVE HF Registry Rationale, Design, and Implementation. *Circulation*. May 30, 2006;113(21):e789 (Abstract #P15).
- Hunt SA, Abraham WT, Chin MH, et al. ACC/AHA 2005 Guideline Update for the Diagnosis and Management of Chronic Heart Failure in the Adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure). *Circulation*. September 20, 2005;112(12):e154-235.
- Strickberger SA, Conti J, Daoud EG, et al. Patient selection for cardiac resynchronization therapy: from the Council on Clinical Cardiology Subcommittee on Electrocardiography and Arrhythmias and the Quality of Care and Outcomes Research Interdisciplinary Working Group, in collaboration with the Heart Rhythm Society. *Circulation*. April 26, 2005;111(16):2146-2150.
- Young JB, Abraham WT, Smith AL, et al, for the Multicenter InSync ICD Randomized Clinical Evaluation (MIRACLE ICD) Trial Investigators. Combined cardiac resynchronization and implantable cardioversion defibrillation in advanced chronic heart failure: the MIRACLE ICD Trial. *JAMA*. May 28, 2003;289(20):2685-2694.

Adapted, with permission, by the SCA Prevention Medical Advisory Team, from the IMPROVE HF registry toolkit.

This is a general algorithm to assist in the management of patients. This clinical tool is not intended to replace individual medical judgment or individual patient needs.


Refer to the manufacturers' prescribing information and/or instructions for use for the indications, contraindications, warnings, and precautions associated with the medications and devices referenced in these materials.

Sponsored by Medtronic, Inc.

April 2007

Revised September 2008

## Device Therapy Algorithm



ICD Therapy is not indicated for patients who do not have a reasonable expectation of survival with an acceptable functional status for at least one year, even if they meet ICD implantation criteria specified in the algorithm. ICD therapy is not indicated for NYHA Class IV patients with drug-refractory congestive heart failure who are not candidates for cardiac transplantation or CRT-D.

<sup>†</sup> Functional status as documented prior to current hospitalization while on chronic optimal medical therapy.

## References

<sup>1</sup> CMS coverage for primary prevention ICD implants: patients with nonischemic dilated cardiomyopathy (NIDCM) > 9 months, NYHA Class II and III heart failure, and measured LVEF  $\leq$  35%. Patients with NIDCM > 3 months and < 9 months, NYHA Class II or III heart failure, and measured LVEF  $\leq$  35% at this time are only covered by Medicare if these patients are enrolled in an FDA-approved category B IDE clinical trial, a trial under the CMS clinical trial policy, or the American College of Cardiology National Cardiovascular Data Registry (ACC-NCDR).

## Reference Source

- Bardy GH, Lee KL, Mark DB, et al, for the Sudden Cardiac Death in Heart Failure Trial (SCD-HeFT) Investigators. Amiodarone or an implantable cardioverter-defibrillator for congestive heart failure [published correction appears in *N Engl J Med*. 2005;352:2146]. *N Engl J Med*. January 20, 2005;352(3):225-237.
- Buxton AE, Lee KL, Fisher JD, Josephson ME, Prystowsky EN, Hafley G. A randomized study of the prevention of sudden death in patients with coronary artery disease. Multicenter Unsustained Tachycardia Trial Investigators [published correction appears in *N Engl J Med*. 2000;342:1300]. *N Engl J Med*. December 16, 1999;341(25):1882-1890.
- Epstein AE, Dimarco JP, Ellenbogen KA, et al. ACC/AHA/HRS 2008 Guidelines for device-based therapy of cardiac rhythm abnormalities. *Heart Rhythm*. June 2008;5(6):e1-62.
- Fonarow GC, Yancy CW, Albert NM, et al. Improving the use of evidence-based heart failure therapies in the outpatient setting: the IMPROVE HF Registry Rationale, Design, and Implementation. *Circulation*. May 30, 2006;113(21):e789 (Abstract #P15).
- Gregoratos G, Abrams J, Epstein AE, et al. ACC/AHA/NASPE 2002 guideline update for implantation of cardiac pacemakers and antiarrhythmia devices: summary article: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (ACC/AHA/NASPE Committee to Update the 1998 Pacemaker Guidelines). *Circulation*. October 15, 2002; 106(16):2145-2161.
- Hohnloser SH, Kuck KH, Dorian P, et al, for the DINAMIT Investigators. Prophylactic use of an implantable cardioverter-defibrillator after acute myocardial infarction. *N Engl J Med*. December 9, 2004;351(24):2481-2488.
- Hunt SA, Abraham WT, Chin MH, et al. ACC/AHA 2005 Guideline Update for the Diagnosis and Management of Chronic Heart Failure in the Adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure). *Circulation*. September 20, 2005; 112(12):e154-235.
- Moss AJ, Hall WJ, Cannom DS, et al. Improved survival with an implanted defibrillator in patients with coronary disease at high risk for ventricular arrhythmia. Multicenter Automatic Defibrillator Implantation Trial Investigators. *N Engl J Med*. December 26, 1996; 335(26):1933-1940.
- Moss AJ, Zareba W, Hall WJ, et al, for the Multicenter Automatic Defibrillator Implantation Trial II Investigators. Prophylactic implantation of a defibrillator in patients with myocardial infarction and reduced ejection fraction. *N Engl J Med*. March 21, 2002;346(12):877-883.


Adapted, with permission, by the SCA Prevention Medical Advisory Team, from the IMPROVE HF registry toolkit.

This is a general algorithm to assist in the management of patients. This clinical tool is not intended to replace individual medical judgment or individual patient needs.

Refer to the manufacturers' prescribing information and/or instructions for use for the indications, contraindications, warnings, and precautions associated with the medications and devices referenced in these materials.

Sponsored by Medtronic, Inc.  
April 2007  
Revised September 2008

## Inpatient to Outpatient Transition Algorithm for Medical and Device Therapy



## References

- <sup>1</sup> Patients with secondary indications for device therapy should undergo evaluation and device placement prior to hospital discharge.
- <sup>2</sup> Please see individual medication and device algorithms for details.
- <sup>3</sup> 3-6 months of optimal medical therapy is suggested before reassessment of LVEF and functional status.
- <sup>4</sup> CMS coverage for primary prevention ICD implants: patients with nonischemic dilated cardiomyopathy (NIDCM) > 9 months, NYHA Class II and III heart failure, and measured LVEF  $\leq$  35%. Patients with NIDCM > 3 months and < 9 months, NYHA Class II or III heart failure, and measured LVEF  $\leq$  35% at this time are only covered by Medicare if these patients are enrolled in an FDA-approved category B IDE clinical trial, a trial under the CMS clinical trial policy, or the American College of Cardiology National Cardiovascular Data Registry (ACC-NCDR).

## Reference Source

- Fonarow GC, Yancy CW, Albert NM, et al. Improving the use of evidence-based heart failure therapies in the outpatient setting: the IMPROVE HF Registry Rationale, Design, and Implementation. *Circulation*. May 30, 2006;113(21):e789 (Abstract #P15).

Adapted, with permission, by the SCA Prevention Medical Advisory Team, from the IMPROVE HF registry toolkit.

This is a general algorithm to assist in the management of patients. This clinical tool is not intended to replace individual medical judgment or individual patient needs.

Refer to the manufacturers' prescribing information and/or instructions for use for the indications, contraindications, warnings, and precautions associated with the medications and devices referenced in these materials.

Sponsored by Medtronic, Inc.  
April 2007



## Guideline Recommendations for Heart Failure Device Therapy

Summary of recommendations for the use of ICD and/or CRT from the *2005 ACC/AHA Guideline Update for the Diagnosis and Management of Chronic Heart Failure in the Adult*<sup>1</sup> and the *2008 ACC/AHA/HRS Guidelines for Device-Based Therapy for Cardiac Rhythm Abnormalities*<sup>2</sup>

Classification of Recommendations	
<b>Class I</b>	Conditions for which there is evidence and/or general agreement that a given procedure or treatment is beneficial, useful, and effective
<b>Class II</b>	Conditions for which there is conflicting evidence and/or a divergence of opinion about the usefulness/efficacy of a procedure or treatment
<b>Class IIa</b>	Weight of evidence/opinion is in favor of usefulness/efficacy
<b>Class IIb</b>	Usefulness/efficacy is less well established by evidence/opinion
<b>Class III</b>	Conditions for which there is evidence and/or general agreement that a procedure/treatment is not useful/effective and in some cases may be harmful

Level of Evidence	
Level of Evidence A	Data derived from multiple randomized clinical trials or meta-analyses
Level of Evidence B	Data derived from a single randomized trial or nonrandomized studies
Level of Evidence C	Only consensus opinion of experts, case studies, or standard of care

Guideline Recommendations Patients with Current or Prior Symptoms of HF (Stage C) Patients with Reduced LVEF Recommendations for ICD Therapy	
<b>Class I</b> Level of Evidence A	An ICD is recommended as secondary prevention to prolong survival in patients with current or prior symptoms of HF and reduced LVEF who have a history of cardiac arrest, ventricular fibrillation, or hemodynamically destabilizing ventricular tachycardia
<b>Class I</b> Level of Evidence A	ICD therapy is recommended for primary prevention to reduce total mortality by a reduction in sudden cardiac death in patients with ischemic heart disease who are at least 40 days post-MI, have an LVEF $\leq 30\%$ with NYHA Class I symptoms or an LVEF $\leq 35\%$ with NYHA Class II/III symptoms while undergoing chronic medical therapy and have a reasonable expectation of survival with a good functional status for more than 1 year



<b>Class I</b> Level of Evidence B	ICD therapy is recommended for primary prevention to reduce total mortality by a reduction in sudden cardiac death in patients with nonischemic cardiomyopathy who have an LVEF $\leq$ 30%, with NYHA Class II or III symptoms while undergoing chronic medical therapy and have a reasonable expectation of survival with a good functional status for more than 1 year
<b>Class IIb</b> Level of Evidence C	ICD therapy may be considered in patients with nonischemic heart disease who have an LVEF $\leq$ 35% with NYHA Class I symptoms

**Patients with Current or Prior Symptoms of HF (Stage C)  
Patients with Reduced LVEF  
Recommendations for CRT Therapy\***

<b>Class I</b> Level of Evidence A	Patients with an LVEF $\leq$ 35%, sinus rhythm, and NYHA Class III or ambulatory Class IV symptoms despite recommended, optimal medical therapy and who have cardiac dyssynchrony, which is currently defined as a QRS duration $>$ 120 ms, should receive cardiac resynchronization therapy unless contraindicated
<b>Class IIa</b> Level of Evidence B	For patients with an LVEF $\leq$ 35%, a QRS duration of $\geq$ 120 ms, and AF, cardiac resynchronization therapy, with or without ICD therapy, is reasonable for the treatment of NYHA Class III or ambulatory Class IV symptoms on recommended optimal medical therapy
<b>Class IIa</b> Level of Evidence C	For patients with an LVEF $\leq$ 35%, with NYHA Class III or ambulatory Class IV symptoms, who are receiving recommended optimal medical therapy and who have frequent dependence on ventricular pacing, cardiac resynchronization therapy is reasonable

\*Inclusion of an ICD to be based on ICD recommendations and physician discretion.

**Reference**

- <sup>1</sup> Hunt SA, Abraham WT, Chin MH, et al. ACC/AHA 2005 Guideline Update for the Diagnosis and Management of Chronic Heart Failure in the Adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure). *Circulation*. September 20, 2005;112(12):e154-235.
- <sup>2</sup> Epstein AE, Dimarco JP, Ellenbogen KA, et al. ACC/AHA/HRS 2008 Guidelines for device-based therapy of cardiac rhythm abnormalities. *Heart Rhythm*. June 2008;5(6):e1-62.

Developed by the SCA Prevention Medical Advisory Team based on the ACC/AHA 2005 Heart Failure Guidelines.

This is a general algorithm to assist in the management of patients. This clinical tool is not intended to replace individual medical judgment or individual patient needs.

Refer to the manufacturers' prescribing information and/or instructions for use for the indications, contraindications, warnings, and precautions associated with the medications and devices referenced in these materials.

Sponsored by Medtronic, Inc.  
April 2007  
Revised September 2008

UC200705457a EN