

Supplemental Table 1. Influencing factors of chronological order of occurrence of the sequelae

Variables	Factors	χ^2	<i>P</i>	95% CI	
Time	(12 month <i>vs.</i> 9 month)	3.449	0.063	-6.285	0.170
	(6 month <i>vs.</i> 9 month)	0.270	0.603	-3.992	2.319
Age	(Youth <i>vs.</i> Old age)	2.215	0.137	-2.159	0.295
	(Middle-age <i>vs.</i> Old-age)	0.380	0.538	-1.412	0.736
Sex	(Female <i>vs.</i> Male)	2.064	0.151	-2.108	0.325
Clinical classification of COVID-19	(Non-severe type <i>vs.</i> Severe and critically type)	0.301	0.583	-1.347	0.757
	(Too light <i>vs.</i> Obesity)	0	0.999	-2.203	2.204
BMI	(Normal <i>vs.</i> Obesity)	0.499	0.480	-0.943	2.007
	(Excessive weight <i>vs.</i> Obesity)	0.933	0.334	-0.746	2.197
Smoking history	(No <i>vs.</i> Yes)	2.438	0.118	-0.313	2.761
History of alcohol intake	(No <i>vs.</i> Yes)	0.031	0.859	-1.250	1.498
Hypertension	(No <i>vs.</i> Yes)	0.539	0.463	-0.710	1.561
Diabetes	(No <i>vs.</i> Yes)	0.666	0.414	-2.034	0.838
Hyperlipemia	(No <i>vs.</i> Yes)	1.102	0.294	-3.029	0.916
Hyperuricemia	(No <i>vs.</i> Yes)	0.163	0.686	-2.551	1.679

BMI: Body mass index; CI: Confidence Interval.