

	# patients (%)	Odds Ratio (95% CI)	aOR† (95% CI)
Lineage			
Euro-American	14/28 (50.0)	Reference	Reference
<u>Indo-Oceanic</u>	<u>13/17 (76.5)</u>	<u>3.3 (0.9 - 13.8)*</u>	<u>3.4 (0.8 - 13.7)*</u>
<u>East-Asian</u>	<u>1/9 (11.1)</u>	<u>0.1 (0.01 - 0.8)*</u>	<u>0.1 (0.01 - 1.2)*</u>
East-African Indian	9/23 (39.1)	0.6 (0.2 - 2.0)	0.6 (0.2 - 2.0)
Age, years			
<2	3/9 (33.3)	0.5 (0.1 - 2.6)	0.5 (0.1 - 2.6)
2-4	2/5 (40.0)	0.7 (0.1 - 5.1)	0.8 (0.1 - 6.1)
5-9	3/5 (60.0)	1.6 (0.2 - 14.0)	1.8 (0.2 - 14.0)
10-15	12/25 (48.0)	Reference	Reference
>15	17/33 (51.5)	1.2 (0.4 - 3.3)	1.3 (0.4 - 4.1)
Gender			
Male	21/42 (50.0)	1.2 (0.5 - 2.9)	1.3 (0.5 - 3.7)
Female	16/35 (45.7)	Reference	Reference
Ethnic Community			
<u>Indian subcontinent</u>	<u>12/40 (30.0)</u>	<u>0.3 (0.1 - 0.9)**</u>	
Africa	10/16 (62.5)	Reference	
Americas	2/3 (66.7)	1.2 (0.1 - 16.2)	
East Asia/SE Asia	11/16 (68.8)	1.3 (0.31 - 5.7)	
Middle East/Central Asia	2/2 (100.0)	9.4x10 ⁶ (0 - ∞)	
Place of Birth			
Canada	10/31 (32.3)	Reference	
<u>Foreign born</u>	<u>27/46 (58.7)</u>	<u>3.0 (1.2 - 8.0)**</u>	
Time since immigration‡			
<2	7/17 (41.2)	0.7 (0.2 - 3.1)	
2-5	8/11 (72.7)	2.7 (0.5 - 17.2)	
6-10	6/12 (50.0)	Reference	
>10	4/4 (100)	4.3x10 ⁷ (0 - ∞)	
Unknown	2/2 (100)	NA	
Born in Canada	10/31 (32.3)	NA	
Drug sensitivity			
Pan-sensitive	31/68 (45.6)	Reference	
Drug-resistant	6/9 (66.7)	2.4 (0.6 - 12.0)	

*p<0.1; **p <0.05

† Adjusted for age and gender

‡ Time since immigration unavailable for 2 individuals, excluded from analysis

Table. The distribution of extrathoracic TB disease (isolated or mixed) based on demographic and clinical characteristics. Unadjusted odds of extrathoracic disease were calculated based on the indicated reference population. 95% confidence interval is shown in parentheses. Adjusted odds ratios shown where applicable.