

	Odds Ratio (95% CI)	aOR (95% CI)
Ethnic Community†		
<u>Indian subcontinent</u>	<u>0.2 (0.08 - 0.5)</u>	<u>0.2 (0.06 - 0.5)***</u>
Africa	2.1 (0.7 - 6.5)	2.4 (0.7 - 7.7)
Americas	2.2 (0.2 - 25.7)	3.3 (0.3 - 41.8)
<u>East Asia/SE Asia</u>	<u>2.9 (0.9 - 9.6)</u>	<u>2.8 (0.8 - 9.4)*</u>
Lineage		
Euro-American	1.1 (0.5-2.9)	1.2 (0.4 - 3.1)
<u>Indo-Oceanic</u>	<u>4.9 (1.4 - 16.7)</u>	<u>5.4 (1.5 -19.2)**</u>
<u>East-Asian</u>	<u>0.1 (0.01- 0.9)</u>	<u>0.1 (0.01- 0.9)**</u>
East-African Indian	0.6 (0.2 - 1.6)	0.6 (0.2 - 1.6)

*p<0.1; **p <0.05; ***p<0.01

† Middle East/Central Asia was excluded from the analysis due to the small number of patients

Table. The distribution of extrathoracic disease (isolated or mixed) based on ethnic community and case isolate lineage. Unadjusted odds odds of extrathoracic disease were calculated based on a dichotomized population. 95% confidence interval is shown in parentheses. Odds ratios adjusted for age and sex are shown for lineage and adjusted for place of birth (Canada or foreign) are shown for ethnic community.