

Supplemental Digital Content 3: Summary of baseline characteristics

| | All (N=1031) | INF (N=202, 19.6%) | IMM (N=83, 8.1%) | UNS (N=746, 72.4%) | P value ¹ |
|----------------------------|-------------------------|--------------------------|------------------------|--------------------------|----------------------|
| Male | 567 (55.0) | 119 (58.9) | 44 (53.0) | 404 (54.2) | 0.4503 |
| Age (years) | 4.0 (1.0 - 10.0) | 2.0 (0.0 - 7.0) | 5.0 (2.0 - 8.0) | 5.0 (1.0 - 10.0) | <0.0001 |
| Age category | | | | | |
| <1 y | 195 (18.9) | 68 (33.7) | 3 (3.6) | 124 (16.6) | <0.0001 |
| 1-4y | 330 (32.0) | 67 (33.2) | 35 (42.2) | 228 (30.6) | |
| 5-9y | 239 (23.2) | 30 (14.9) | 31 (37.4) | 178 (23.9) | |
| 10-14y | 206 (20.0) | 26 (12.9) | 10 (12.1) | 170 (22.8) | |
| ≥ 15y | 61 (5.9) | 11 (5.4) | 4 (4.8) | 46 (6.2) | |
| Invasive ventilation | 808/1024 (78.9) | 155/202 (76.7) | 62/83 (74.7) | 591 /739 (80.0) | 0.3751 |
| DOIV | 3.0 (2.0 - 5.0) (N=806) | 3.0 (2.0 - 5.0) | 4.5 (2.0 -10.0) | 2.0 (2.0 - 5.0) | <0.0001 |
| Vasoactive treatment (VT) | 216/983 (22.0) | 46 /200 (23.0) | 14/79 (17.7) | 156/704 (22.2) | 0.6157 |
| Renal support | 50/890 (5.6) | 5/194 (2.6) | 12/76 (15.8) | 33/620 (5.3) | 0.0001 |
| PIM 2 score | 3.3 (1.9 - 4.8) | 3.4 (1.5 - 5.7) | 2.9 (1.5 - 3.9) | 3.4 (2.3 - 4.8) | 0.0181 |
| PIM2 score category | | | | | |
| <1% | 96 (9.3) | 26 (12.9) | 9 (10.8) | 61 (8.2) | 0.0038 |
| 1 to <5% | 699 (67.8) | 115 (56.9) | 64 (77.1) | 520 (69.7) | |
| 5 to <15% | 180 (17.5) | 42 (20.8) | 9 (10.8) | 129 (17.3) | |
| 15 to <30% | 19 (1.8) | 8 (4.0) | 0 (0) | 11 (1.5) | |
| 30%+ | 37 (3.6) | 11 (5.4) | 1 (1.2) | 25 (3.4) | |
| Length of PICU stay (days) | 2.3 (1.1- 4.9) | 2.5 (1.2 - 4.9) | 3.5 (1.8 - 9.0) | 2.1 (1.1 - 4.7) | 0.0013 |

¹ Kruskal Wallis / Chi Square Test

| | All (N=1031) | INF (N=202, 19.6%) | IMM (N=83, 8.1%) | UNS (N=746, 72.4%) | P value ¹ |
|------------------------------|--------------|--------------------------|------------------------|--------------------------|----------------------|
| Length of PICU stay category | | | | | |
| <1d | 218 (21.1) | 41 (20.3) | 11 (13.3) | 166 (22.3) | 0.0009 |
| 1 to <3d | 395 (38.3) | 77 (38.1) | 25 (30.1) | 293 (39.3) | |
| 3 to <7d | 262 (25.4) | 61 (30.2) | 21 (25.3) | 180 (24.1) | |
| 7 to <14d | 97 (9.4) | 18 (8.9) | 14 (16.9) | 65 (8.7) | |
| 14 to <28d | 35 (3.4) | 5 (2.5) | 8 (9.6) | 22 (2.9) | |
| 28d+ | 24 (2.3) | 0 (0) | 4 (4.8) | 20 (2.7) | |
| Mortality | 87 (8.4) | 22 (10.9) | 4(4.8) | 61 (8.2) | 0.2181 |

Key: Data shown are median (interquartile range, IQR) for continuous variables or number (%)

for categorical variables. Data presented are for the entire cohort unless where a denominator is specified. Where presented, each denominator represents the number of children who received a particular intervention and had available data. INF=infective encephalitis, IMM=immune mediated encephalitis; UNS= unspecified encephalitis; DOIV=duration of invasive ventilation; PIM2= Pediatric Index of Mortality score, second version.

¹ Kruskal Wallis / Chi Square Test