## **SUPPLEMENTAL DIGITAL CONTENT 1.** High- and medium-risk criteria for sepsis in children per National Institute for Health and Care excellence guidelines:

Risk Level	Category	Criteria
High Risk	Behavior	<ul><li>No response to social cues</li><li>Appears ill or unwell</li><li>Unable to wake or stay awake</li></ul>
	Respiratory	<ul> <li>- Grunting</li> <li>- Apnea</li> <li>- Severe chest indrawing</li> <li>- Respiratory rate: ≥60 breaths/min (any age)</li> </ul>
	Circulation and Skin	<ul> <li>Cyanosis of skin, lips, or tongue</li> <li>Mottled, ashen, or pale appearance</li> <li>Cold extremities</li> <li>Heart rate: &lt;60 bpm (bradycardia, any age)</li> <li>Capillary refill time: &gt;3 seconds</li> </ul>
	Other	- Hypotension (age-specific thresholds)
Medium Risk	Behavior	- Not responding normally to social cues - Wakes only with prolonged stimulation
	Respiratory	<ul> <li>Nasal flaring</li> <li>Oxygen saturation &lt;92% in air or increased oxygen requirement</li> <li>Respiratory rate: &gt;50 breaths/min (2–12 months), &gt;40 breaths/min (1–5 years)</li> </ul>
	Circulation and Skin	<ul> <li>Pale or flushed skin</li> <li>Reduced urine output</li> <li>Capillary refill time: 2–3 seconds</li> <li>Heart rate: &gt;160 bpm (under 1 year), &gt;150 bpm (1–2 years), &gt;140 bpm (2–5 years)</li> </ul>
	Other	<ul> <li>- Temperature: ≥38°C or ≤36°C (in children under 3 months)</li> <li>- Signs of infection (e.g., rash, redness)</li> </ul>