

Mental Health Issues and Psychological Factors in Athletes: Detection, Management, Effect on Performance, and Prevention: American Medical Society for Sports Medicine Position Statement

Cindy J. Chang, MD,*# Margot Putukian, MD, FACSM, FAMSSM,†# Giselle Aerni, MD,‡ Alex B. Diamond, DO, MPH,§ Eugene S. Hong, MD,¶ Yvette M. Ingram, PhD,|| Claudia L. Reardon, MD,** and Andrew T. Wolanin, PsyD††

Abstract: The American Medical Society for Sports Medicine convened a panel of experts to provide an evidence-based, best practices document to assist sports medicine physicians and other members of the athletic care network with the detection, treatment, and prevention of mental health issues in competitive athletes. This statement discusses how members of the sports medicine team, including team physicians, athletic trainers, and mental health providers, work together in providing comprehensive psychological care to athletes. It specifically addresses psychological factors in athletes including personality issues and the psychological response to injury and illness. The statement also examines the athletic culture and environmental factors that commonly impact mental health, including sexuality and gender issues, hazing, bullying, sexual misconduct, and transition from sport. Specific mental health disorders in athletes, such as eating disorders/disordered eating, depression and suicide, anxiety and stress, overtraining, sleep disorders, and attention-deficit/hyperactivity disorder, are reviewed with a focus on detection, management, the effect on performance, and prevention. This document uses the Strength of Recommendation Taxonomy (SORT) to grade level of evidence.

Key Words: mental health, athlete, sports psychology, depression, anxiety, eating concerns, gambling, hazing, psychological response to injury

(*Clin J Sport Med* 2020;00:1–27)

BACKGROUND AND PURPOSE

Although participation in athletics has many benefits, the very nature of competition can provoke, augment, or expose specific psychological issues in athletes. Certain personality traits can aid in athletic success, yet these same traits can also be associated with mental health (MH) disorders. The athletic culture may have an impact on performance and psychological health through its effect on existing personality traits and

MH disorders. This article will focus on the competitive athlete, from the youth and collegiate athlete to the Olympian and professional athlete, and how the athletic care network¹ and MH care providers can assist with the detection and treatment of psychological issues in this population.

Unique signs and symptoms in athletes, prevalence in the athlete population, and utilization of available screening tools will be reviewed. Specific Diagnostic and Statistical Manual of Mental Disorders (DSM-5) diagnostic criteria and the pathophysiology of MH disorders will not be discussed. The discussion of management may include psychosocial approaches and pharmacological treatments, emphasizing the selection of the most effective treatments with the fewest side effects of relevance for athletic performance. Lastly, this paper will present recommendations for prevention, including the identification and possible elimination of risk factors in the athlete population.

METHODS

The American Medical Society for Sports Medicine (AMSSM) Board of Directors appointed co-chairs (C.J.C. and M.P.) to assemble a writing group that was carefully selected to include a balanced panel of sports medicine physicians and other professionals experienced in managing MH issues in athletes, actively engaged in research, and with demonstrated leadership in the topic. Important members of the panel included an athletic trainer (AT), a clinical psychologist, and a sport psychiatrist. The co-chairs generated the outline and the

Submitted for publication November 25, 2019; Accepted December 3, 2019.

From the *Departments of Orthopaedics and Family & Community Medicine, University of California, San Francisco, San Francisco, California; †Department of Athletic Medicine, University Health Services, Princeton University, Princeton, New Jersey; ‡Department of Sports Medicine and Family Medicine, WellSpan Health, York, Pennsylvania; §Departments of Pediatrics and Orthopaedic Surgery, Vanderbilt University Medical Center, Nashville, Tennessee; ¶Departments of Orthopaedics and Family Medicine, Medical University of South Carolina, Charleston, South Carolina; ||Department of Health Science, Lock Haven University, Lock Haven, Pennsylvania; **Department of Psychiatry, University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin; and ††Wolanin Consulting and Assessment Inc, Bala Cynwyd, Pennsylvania.

The authors report no conflicts of interest.

#C. J. Chang and M. Putukian are Co-Chairs, and Co-Lead authors.

Corresponding Author: Margot Putukian, MD, FACSM, FAMSSM, Department of Athletic Medicine, University Health Services, Princeton University, Princeton, NJ 08544 (putukian@princeton.edu).

Copyright © 2020 AMERICAN MEDICAL SOCIETY FOR SPORTS MEDICINE. All rights reserved. The AMERICAN MEDICAL SOCIETY FOR SPORTS MEDICINE has granted Wolters Kluwer Health, Inc. permission for the reproduction of this article. <http://dx.doi.org/10.1097/JSM.0000000000000817>

writing group subsequently conducted an in-depth literature review using PubMed, SportDiscus, and the Cochrane database for each topic. The writing group engaged in conference calls and written communications to discuss the evidence and compile the manuscript. The panel used the Strength of Recommendation Taxonomy (SORT) to grade level of evidence⁷ (Table 1).

This AMSSM position statement is novel in several respects in its contribution to the topic, including the following:

- addressing topics not fully explored in previous publications about MH issues in athletes, including key personality issues, demographic and cultural variables, and environmental conditions;
- discussing the interaction and impact of these variables both positively and negatively on competitive athletes, and how to monitor the athletic environment that may precipitate or exacerbate MH issues; and
- defining the level of evidence and the knowledge gaps in this rapidly expanding field.

HOW TEAMS WORK

In many sport settings, the team physician, working with ATs and other members of the athletic care network, provides medical care to athletes; this should include management of health and wellness issues, including MH disorders.² The team physician often coordinates multiple aspects of the athletes' overall care and may oversee MH treatment, including psychiatric medication management.³ The MH care network^{4,5} includes clinical or counseling psychologists, neuropsychologists, psychiatrists, licensed clinical social workers, MH nurses, licensed MH counselors, and primary care physicians with core competencies (ie, additional expertise) to treat MH disorders. It is critical that members of the MH care network and athletic care network have clear, established guidelines of how to best communicate to optimize the treatment of the athlete.

It is important that team physicians attempt to normalize health-seeking behaviors and understand that there may be barriers in seeking care that athletes uniquely experience. Providing screening for depression, anxiety, eating disorders (EDs), and substance use, as part of preparticipation sports physical examinations, can help in normalizing discussions regarding these important MH disorders. It is also important that team physicians include protocols to allow for both routine MH referrals as well as MH emergencies, as part of the emergency MH action plan.³

Strength of Recommendation	Basis for Recommendation
A	Consistent, good-quality patient-oriented evidence
B	Inconsistent or limited-quality patient-oriented evidence
C	Consensus, disease-oriented evidence, usual practice, expert opinion, or case series for studies of diagnosis, treatment, prevention, or screening

PERSONALITY ISSUES AND ATHLETIC CULTURE

There is an interactive relationship between athlete personality characteristics and the athletic culture that may have both positive and negative effects on the individual athlete as well as the team or sport environment. This relationship between internal and external variables is fluid and may mirror changes in culture as a whole and changing developmental patterns of individuals in their time of athletic engagement. Internal variables may include traits such as perfectionism, pessimism, or introversion, while external variables may include factors such as coaching culture, team performance, or socioeconomic status.

For example, the internal trait of perfectionism may contribute to anxiety about physical appearance in the athletic environment,⁶ while the external coaching environment may also influence athlete anxiety,⁷ or the internal trait of pessimism, in addition to the external factor of low socioeconomic status, may result in adjustment and performance issues. An athlete's personality traits may also influence the athletic culture. For example, athletes who engage in aggressive or high-risk behaviors may see a performance benefit in sport, while simultaneously possibly being prone for substance use disorders, hazing or other behavioral issues outside of sport because they are high-risk takers.

The following sections will provide insight into some of the key personality and environmental issues that impact athletes and their MH. Consequently, it is important for team physicians to be attuned to personality styles that may have a higher risk of dysfunctional behaviors and athletic environments that may trigger or exacerbate MH disorders in the athletes under their care.

Personality Issues

Detection in Athletes

There has been much interest in whether or not there is an "athletic personality" that is particularly conducive to success or failure in sport. Personality in general is defined as the stable, consistent aspects of one's thinking, emotions, and behaviors. Findings regarding athletic personality have been inconsistent.⁸ There is no single personality profile that reliably appears in athletes or predicts athletic success or failure. However, many studies suggest that athletes generally show more positive personality characteristics than do nonathletes.^{9,10} Specifically, athletes show greater extraversion and conscientiousness and less hastiness and anger.^{10,11} Athletes also generally show higher sensation-seeking tendencies than do nonathletes.⁹ Contact sport athletes and male athletes express higher tendencies toward sensation-seeking than noncontact sport athletes and female athletes, respectively.¹²

Research looking at individual versus team-sport athletes also demonstrates some personality differences. Individual sport athletes demonstrate more conscientiousness and autonomy.¹³ Team-sport athletes score higher on agreeableness and a tendency to attend to and depend on others for personal satisfaction.¹³

Perfectionism is defined as an achievement-related personality trait that includes the setting and pursuit of excessively high standards of performance together with overly critical self-evaluations.^{14,15} Some distinguish *positive* perfectionism,

with the underlying motivation being to obtain a favorable outcome, from *negative* perfectionism, which seeks to avoid adverse consequences.¹⁶ Both types may be risk factors for the development of EDs in athletes.⁸

Athlete identity is the degree to which an individual views themselves within the athletic role and looks to others for confirmation of that role.¹⁷ High athletic identity has been associated with both positive outcomes, such as better athletic performance, and negative outcomes, such as overtraining and the use of performance-enhancing drugs.¹⁷ Strong tendencies for athletes to evaluate themselves exclusively according to their athletic performance may be associated with depression.¹⁸

Personality testing of individual athletes is not typically undertaken in the clinical setting, as unstructured clinical interview is typically the preferred method for the assessment of personality traits and disorders.^{19,20} However, a variety of validated instruments are available for objective assessment or research.²¹ One such instrument is the Minnesota Multiphasic Personality Inventory-2 (MMPI-2).²²

Management

By definition, personality traits and disorders are long-standing issues. If deemed problematic for athletes, they may be addressed through psychotherapy.²³ Comorbid psychiatric disorders should be treated with medication, psychotherapy, or a combination.²³ Unique challenges related to personality issues have been described when undertaking psychotherapy with high-level athletes in particular.²³ Because this population may be accustomed to being in charge, with others around them giving them special attention and being solicitous of their opinions due to their celebrity status, a “situational narcissism” can develop that can contribute to entitlement regarding scheduling of appointments and payment of fees for services.²³ It is important for health care providers to be aware that establishing a relationship that does not follow usual treatment parameters may lead to boundary violations.

Effect on Performance

Athletes who compete at professional levels have higher self-esteem and sensation-seeking tendencies than those who do not compete at that level.⁹ In addition, lower levels of self-esteem and sensation-seeking in athletes have been associated with greater risk for depression, anxiety, social anxiety, and negative physical symptoms, all of which may impede performance.⁹ Extremes of athletic identity, either too much or too little, may limit performance.¹⁸ Externally driven perfectionism, that is, perfectionism encouraged by parents, coaches, teammates, or others, is likely more problematic for performance than is internally driven perfectionism. Externally driven perfectionism is associated with negative mood, anxiety, anger, and hostility, all of which can decrease performance.²⁴ If perfectionism leads to EDs, there can also be negative performance effects.

Sexuality and Gender Issues

Detection in Athletes

Athletes may be adversely affected by discrimination and other environmental factors in sport related to sexuality and

gender. For example, there may be discrimination against sexual minorities, defined as sexual groups other than heterosexuals [including members of the lesbian, gay, bisexual, transgender, and questioning/queer (LGBTQ) populations]. The masculine “culture of toughness” can also contribute to stigma that prevents athletes from seeking MH treatment.

The psychological issues experienced by sexual minority athletes may be associated with the anxiety of “coming out,” the fear of being ridiculed by teammates, pressures from others to be “normal,” or because they may have been a victim of or witnessed homophobic behaviors. The 2012 National Collegiate Athletic Association (NCAA) Campus Pride Report Score Card indicated 4.6% of student-athletes identified as either lesbian, gay, bisexual, or questioning compared with 6.2% of college students.²⁵ Sexual minority athletes have long been subjected to negative, homophobic attitudes of other athletes and fans. Thirty-two percent of male athletes and 14% of female athletes recall sexuality-related bullying when engaged in school athletics.²⁶ Lesbian, gay, bisexual, or transsexual identification is a risk factor for harassment and abuse in sport.²⁷ Gay male athletes have concerns that their teammates, and coaches will discover their sexual identity, and they fear being “outed.”²⁸ The term “lesbian” may be used inappropriately to stereotype female athletes.²⁹ Across genders, sexual minority student-athletes have reported experiences of negative MH harm in the last 12 months in significantly greater numbers than their heterosexual peers.³⁰

There seem to be more homophobic attitudes in male-dominated sports than in female sports.³¹ If they are aware of these negative attitudes before entering the athletic realm, some sexual minority male athletes will attempt to avoid sport all together.^{26,32,33} When analyzing the differences between male and female sexual minority student-athletes, researchers found that female athletes reported lower rates of negative MH outcomes than their male and nonathlete peers.³⁰ However, female athletes were more likely to report use of both illicit and prescription drugs when compared with heterosexual female student-athletes, and as a group, sexual minority student-athletes have reported a greater likelihood of use of illicit drugs than have heterosexual student-athletes. Student-athletes identifying as sexual minority have an increased risk for poor health, possibly due to avoidance of athletics and associated lower rates of physical activity and increase in drug use. This subsequently leads to poor MH and an increased risk of chronic medical diseases such as obesity and cardiac issues.³⁰

Athletes of any sex, gender identity, and sexual orientation suffer from stigma regarding mental illness, in part related to the masculine culture of toughness, and thus may have difficulty admitting to psychiatric symptoms and asking for and accepting treatment for those symptoms.³⁴ Athletes attempt to avoid displaying vulnerability on the playing field, making it hard to transition to being able to admit vulnerability and accept help when off the field. Moreover, athletes perceive that they risk playing time, starting roles, or endorsements if discovered seeking treatment for mental illness.³⁴

Management

Health care providers should ask athletes about any concerns related to gender and sexuality, and athletes should be educated about the role of sexuality- and gender-related

issues in sport and informed that they are not immune from mental MH disorders. The creation of a supportive environment that is welcoming to sexual minorities is key to the health of athletes and their teams. Sensitivity training for student-athletes and coaches,²⁶ debunking stereotypes, increasing exposure to LGBTQ community, creating open and educational discussions, and using correct terminology³⁵ are important.³⁶ Having gender-neutral locker rooms available to sexual minority athletes and uniforms that allow all participants to be comfortable are also important.²⁶ Policies and laws can have a role; for example, antidiscrimination policies in university settings can protect against homophobic actions.³⁷

Numerous organizations have developed educational programs to assist athletes/teams, coaches, officials, administrators, institutions, and other sport-related groups in overcoming the challenges of discrimination and stigma. The NCAA LGBTQ Inclusion Initiative³⁸ reports that one of its core values is a commitment to diversity, inclusion, and gender equity in collegiate sport and available on its website is a document outlining best practices and sample policies and resources for allies and sexual minority student-athlete groups.³⁹ GLSEN (formerly known as Gay, Lesbian, and Straight Education Network)^{40,41} provides educational resources for parents, schools, and teams aimed at creating and maintaining an athletic climate that is based on respect, safety, and equal access for all students regardless of sexual orientation or gender identity. Go! Athletes is an online social support network consisting of current and former sexual minority collegiate and high school athletes and coaches.⁴² Athlete Ally is an online forum designed to provide support for the sexual minority athlete community, with its mission to eliminate homophobia and transphobia in sports.⁴³

Effect on Performance

There are minimal data evaluating the effect on performance, though it is likely that negative attitudes related to sexuality and gender negatively impact sports performance. Female and sexual minority athletes may be deterred from participating in sports. One study demonstrated that 46% of female athletes felt their involvement in sports led others to conclude that they were lesbian,²⁹ and this may lead to hesitation to become involved in certain sports in particular. If they identify as LGBTQ, some athletes report that they cannot be open with their sexuality for fear of losing sponsorships.²⁹ Athletes may also feel the need to guard against any behavior that suggests close relationships with teammates of the same sex. With the masculine culture of toughness making it difficult for many athletes to reach out for help, these potentially distracting issues and associated potentially untreated MH disorders are likely to have a negative impact on physical performance.

Hazing

Detection in Athletes

The terms hazing and bullying are often used together; however, it is important to differentiate these 2 forms of interpersonal violence.⁴⁴ Hazing is any nonrepetitive, humiliating, or dangerous activity expected of a student (irrespective of their willingness to participate) to become part of the group.^{45,46} Bullying will be discussed in the next section of the paper.

Although cases of extreme hazing result in significant morbidity and mortality and receive considerable public attention, the true incidence of hazing that occurs in sports is believed to be common but is not well known. Greatest reporting rates have been seen in higher levels of competition, team sports, and contact sports.⁴⁷ Rates appear equal across genders, although boys tend to be subjected to more physical forms of hazing.⁴⁸ It remains largely unclear why certain personal demographics, sports, or levels of participation lead to more or fewer hazing-related activities. Populations at higher risk for any type of abuse involve elite athletes, children, LGBTQ groups, disabled athletes,⁴⁹ and those with a lower grade point average.⁵⁰ These characteristics may also place these individuals at a higher risk for hazing specifically. Team characteristics that lead to a higher risk of hazing include denial or failure to recognize the authority of the coaching staff, an unsupervised team area or locker room, and an imbalance of power shifted toward masculine authority.⁴⁹

Eighty percent of NCAA athletes experienced some form of hazing during college, while 42% of these same athletes also reported a history of being hazed in high school.⁵¹ A similar study almost a decade later revealed that 74% of student-athletes experienced at least one form of hazing while in college.⁴⁶ Given that there are over 380 000 student-athletes participating in NCAA sports,⁵² it is likely that 280 000 student-athletes were potential victims of hazing in college athletics alone.

In youth sports, the incidence of hazing is reported to increase with age, starting with 25% of high school athletes reporting their first incident occurring before the age of 13.⁵³ Rates of hazing in sport ranged from 5% to 17.4% in middle schools and 17.4% to 48% at the high school levels.^{48,54}

Although athletes may endorse participation in activities that meet criteria for hazing, hazing itself is grossly underreported. In fact, 60% to 95% of college athletes who were the victim of hazing stated that they would not report their incident.^{46,51} Some of the reasons espoused for the refusal to report included allegiance to fellow teammates, fear of retribution, uncertainty of trust with authority, normalization or positive perception of hazing behavior, and victim blaming, with the perception that participants choose to be involved with hazing activities.^{46,51,53,55-57}

Management

The management of hazing requires a global investment from athletes, coaches, administrators, and health care providers all playing key and specific roles as outlined in Table 2. Although adults must set the tone of zero tolerance toward hazing and promote an environment of respect and dignity, it is crucial that athletes take ownership of the process. A comprehensive approach that involves education, incorporation of programs with validated measures, and the establishment of clearly defined policies and penalties will allow the necessary shift from reacting to incidents to preventing them.

Early engagement of MH providers is critical to care for the victims of hazing. Engagement of the athletic care network, coaching staff, athletes, and administration is important to address factors that allowed hazing to happen. Creating support networks and open lines of communication are important parts of providing advice and help for athletes

TABLE 2. Key Roles in Hazing Awareness and Prevention

Administrators	Coaches	Student-Athletes	Team Captains
Dispel myths/set the record straight	Create environment of total respect and dignity for all	Understand what constitutes hazing	Encourage others to speak up without fear of retribution
Ensure comprehensive approach	Address topic/consequences early and consistently	Insist on open communication among teammates and coaches	Understand potential level of accountability
Provide appropriate resources/support	Involve the team in process; be supportive	Plan alternate activities that enhance bonding	Ensure new members feel supported
Establish protocols and procedures for the process (reporting, investigating, and adjudicating)	Help develop positive traditions that are significant and meaningful	Understand team acceptance and success should be based on strong work ethic, positive attitude, positive relationships	Help develop positive traditions that are significant and meaningful
Enforce clearly defined sanctions	Discuss expectations about character and values on/off field	Discuss among your team how to report potential incidents	Create conduct guidelines along with coaches
	Take it seriously	Make caring about each other a high priority	Accept role as leader; set proper tone with your actions and words
	Avoid creating division between veterans and new players	Seek advice from trusted adults/leaders	

Adapted with permission from Wilfert.⁹⁷ Copyright 2007 by the National Collegiate Athletic Association.

regarding issues of hazing. A NCAA survey demonstrated that men were most likely to turn first to teammates (27%), while women were most likely to turn first to their parents (29%).⁵⁸

One of the most significant roles for the team physician is to recognize a problem has occurred, followed by appropriate response, reporting, and referral. The team physician should encourage open disclosure by serving as an empathetic, nonjudgmental listener who creates an environment of support for the athlete, where they can feel safe to discuss events that have occurred.

All forms of hazing (physical, psychological, and/or substance-related) are maltreatment and include the potential for significant harm, which can be underestimated. Of those who experienced hazing, 71% reported awareness of negative consequences, including relationship difficulties and physical and psychological symptoms.⁵⁹ Psychological sequelae include anxiety, depression, EDs, suicidal thoughts and behaviors, loss of confidence and self-esteem, aggression toward self and/or others, delinquency, criminality, interpersonal conflicts mostly centered on trust issues, sexual difficulties, emotional instability, substance use, and changes in weight, energy, sleep, and concentration, all of which can lead to several additional health issues.^{59,60} The treatment of some of these selected issues will be discussed in detail in other sections of this article.

Effect on Performance

Cycles of hazing may be perpetuated by those who believe these activities support team building. However, it has been shown that as hazing increases, the less attraction, team bonding, and team closeness is felt by the athlete.⁶¹ Conversely, there were higher levels of team social cohesiveness if an athlete participated in appropriate team building activities that promote dignity, positive MH, and teamwork as opposed to victimization.⁶¹ Therefore, the dangerous consequences of hazing provide no added benefit to the concept of team unity and may likely have a paradoxical effect, theoretically resulting in a poorer individual and team performance.

Prevention

Properly addressing the needs of athletes requires the development of an extensive prevention plan regarding hazing. This plan should be centered on a zero-tolerance policy for all types and degrees of hazing and incorporate principles and practices that safeguard the health and wellness of the athlete. In one study, only 39% of college students were introduced to anti-hazing policies when joining teams or organizations, and just 15% attended hazing prevention workshops hosted by either adults or peers.⁴⁶ However, recent NCAA data show that over 70% of college athletes indicated that a coach or athletic department official had talked to them about hazing.⁵⁸

For meaningful change to be enacted, the culture of hazing within sport needs to be addressed. It should be understood that hazing is an inappropriate and misguided example of a traditional initiation. There should be no allowance of inappropriate language or “a little hazing,” which can lead to more severe forms of hazing. Situations that involve secrecy, alcohol and other substances, and power imbalance should be eliminated because these are ripe conditions for hazing incidents to occur. Finally, it is important to encourage spheres of influence (eg, media and professional teams) to avoid intentionally or unintentionally delivering messages that promote hazing.

Bullying

Detection in Athletes

Bullying in athletics is another under-researched topic. Defining the term can be difficult because actions and comments that one athlete perceives as bullying may be experienced by another athlete, who responds positively to negative feedback, as motivating. Bullying has been defined as “abuse and mistreatment of someone vulnerable by someone stronger, more powerful.”⁶² This negative behavior can be exhibited by teammates or coaches and includes yelling, physically trying to overpower an individual, or using body language to intimidate an athlete. It can also involve a coach

making an athlete feel that they are worthless, despised, inadequate, or valued only as a result of their athletic performance.⁶³

Bullying has been reported to be less prevalent in sport than in school and occurring at a relatively low frequency overall.⁶⁴ Participants who reported experiencing one or more acts of bullying on their team reported weaker connections with peers. Detecting bullying in athletics can be difficult because both the athlete and the coaching/team environment need to be observed. Swigonski et al⁶⁵ provided advice to parents to look for 4 defensive behaviors by the coach that may help determine whether their child's coach is a bully. The first behavior is moral justification of their actions by attempting to explain their poor behavior as part of the athletic culture. A second is the coach offering a "backhanded" apology when called out about their poor behavior. The third defensive behavior is an attempt to shift the focus of the conversation away from the bullying behavior to something more severe like physical contact; by doing so, they are explaining that the behavior is not as bad as it could be. The fourth is the coach trying to escalate the situation by attempting to "back the complainant into a corner," so that they will stop questioning the coach's behavior.

Cyberbullying is another form of bullying, with one study reporting that 38% of students knew someone who had been cyberbullied, whereas almost 9 percent reported cyberbullying someone else.⁶⁶ In the 2017 NCAA Study of Student Athlete Social Environments report, 12% of male and 4 percent of female student-athletes reported receiving negative or threatening messages from fans through social media.⁶⁷ Black student-athletes reported receiving negative or threatening messages at twice the rate of white student-athletes.⁶⁸ Negative social media engagement is not limited to college-age student-athletes; student-athletes in secondary schools are also victims.

Athletes may present with changes in sleeping and eating patterns, frequent mood swings, aggressive behavior, apprehension about attending team functions, or physical symptoms such as feeling ill before practices.⁶⁹ Depending on the severity of the bullying, athletes may present with unexplained injuries, loss of personal items, frequent headaches, or faking illness or injuries.⁵⁹ They may also contemplate and/or attempt suicide, exhibit poor psychosocial adjustment, experience difficulty with academics that once came much easier, or display rebellious behaviors such as skipping practices or misusing substances.

No screening tool specifically designed for assessing bullying in athletics exists; however, the Bullying and Ostracism Screening Experience and School Climate (BOSS-E) short form⁷⁰ is a 16-item self-report questionnaire that health care providers and administrators may consider for implementation. It is used to provide a quick indicator of school climate for boys and girls aged 10 to 14 years. A second screening tool available to assess bullying behaviors is the 3-part California Bully Victimization Scale instrument.⁷¹

Management

The management of bullying starts with athlete, parent, school/athletic administrator, athletic care network, and coach intervention. All stakeholders must be aware of both the psychological signs and the physical symptoms that can be exacerbated by bullying and exhibited by an athlete, to intervene on their behalf.

The athletic care network should work with affected athletes to help resolve the issue at both the individual and system levels. Possible resolutions include acting as an advocate for the athletes, referring them to counseling, offering educational programs to the athletics department, or reporting the behavior of the bully up the administrative chain while adhering to any confidentiality requirements. Creating an overall supportive environment where all stakeholders agree that bullying is unacceptable is the key to preventing repeat occurrences and providing a safe environment in which athletes can participate.

Effect on Performance

Because most research uses the terms bullying and hazing interchangeably, effect on performance is discussed under the hazing section above.

Prevention

Athletes need to be taught that bullying is not acceptable and should be reported to an uninvolved, trusted adult. Athletes should be encouraged to defend their teammates who are experiencing bullying or help remove them from the situation. Step UP! is an example of a prosocial behavior and bystander intervention program.⁷² The NCAA website has details and links for Green Dot, Mentors in Violence Prevention, and Bringing in the Bystander educational programs.⁷³

Sexual Misconduct

Detection in Athletes

Another form of misconduct and maltreatment of athletes is sexual misconduct. The exact incidence of sexual misconduct (including sexual harassment, sexual abuse, gender harassment, and homophobic behaviors) in sports is difficult to determine because of a lack of reporting, particularly by boys. Sexual misconduct has been shown to occur in all sports across all ages and levels of participation.⁷⁴ It is estimated that 2 percent to 48% of athletes will experience sexual abuse, and 19% to 92% will be subjected to some form of sexual harassment.^{27,51,54,75,76} Peer athletes are far more likely than coaches to perpetrate sexual harassment.²⁷ Male athletes are typically more often the offender, with 34% of women experiencing sexual harassment from men and 12% from women.⁷⁷ In one study, 13% of female athletes and 6% of male athletes had experienced sexual abuse in the sport setting.⁷⁸

Specifically regarding underage athletes, in one study, 3 percent of coaches admitted to having intimate relations with an athlete under the age of 18.⁷⁹ Two percent to 22% of children and teens are victims of sexual abuse through sport, with 98% of cases being perpetrated by those in positions of power such as coaches, teachers, and instructors, with victims more often being male.⁸⁰

The prevalence of sexual abuse seems to be higher in elite-level athletes where the risks of being sexually exploited by coaches and team support staff are higher.^{54,75} It is unclear whether this is due to intrinsic changes in the athlete-coach relationship or reflects greater exposure to high-risk situations.⁸⁰⁻⁸² In athletes who compete in individual sports, the period of "imminent achievement" is a distinctly

vulnerable period. In this developmental phase, athletes are considered pre-elite but have not yet acquired an elite-level status. There is often a heightened level of stress and dependence on coaches and training staff, which in turn may leave the athlete more vulnerable to predation. The pre-elite athlete is also more likely to tolerate inappropriate behaviors rather than compromise pending achievement.⁸³ Athletes who specialize at a younger age, particularly around puberty, have been found to be highly vulnerable to sexual abuse.⁴⁸

Populations at higher risk for abuse in youth sport reflect trends in the general population. Disabled athletes are at 2 to 3 times increased risk as compared to young athletes in general, and LGBTQ athletes are also at markedly higher risk for abuse.^{84,85} Sport type, amount of touching, or degree of clothing cover during participation do not appear to correlate with rates of abuse.^{86,87} Unsupervised situations (locker room, travel/away trips, and coach's car or home), times of isolation (individual training sessions), and group events that involve alcohol are particularly high-risk environments that increase the risk for sexual harassment and abuse.^{81,82,88,89}

Psychological abuse often serves as the "gateway" to other forms of nonaccidental violence,²⁷ and 75% of youth athletes reported emotional abuse during organized sport.⁹⁰ Sexual abuse can be prefaced by "grooming behavior" where a coach shows preferential treatment toward an athlete, and sometimes even the parents, in efforts to gain favor and trust.

The victimized athlete may present with various nonspecific and recurrent medical and somaticizing concerns including headaches, lethargy, sleep disturbances, bed wetting, acting out or engagement in risky behaviors, self-harm, weight fluctuations, and poorer general health satisfaction.⁷⁴ Previous sexual abuse results in an increased risk of developing a wide array of psychiatric conditions and substance abuse.⁷⁴

Management

The International Olympic Committee (IOC) has defined "safe sport" as an athletic environment that is respectful, equitable, and free from all forms of nonaccidental violence to athletes.²⁷ Sporting organizations should foster the evolution and growth of the "safe sport" movement to properly address the culture of maltreatment and violence in sport, including all forms of physical, sexual, and psychological abuse. Barriers to implementation include a lack of knowledge and resources as well as fear and administrative difficulties.⁹¹ Therefore, it may be challenging to transfer policies that have been established at the sports federation or national governing body level to the local, individual club level.^{92,93}

Recommendations have been directed primarily at administrative and athletic staff personnel, with a focus on education to prevent, recognize, and react to instances of abuse. In the United States, Title IX in 1972 included that "No person in the United States shall, on the basis of sex...be subjected to discrimination under any education program or activity receiving federal financial assistance."⁹⁴ In recent legal proceedings, it has been an important litigation tool in challenging the power structure that has perpetuated instances of sexual violence in the sporting arena in the secondary school and collegiate levels.⁸¹

Consequences of abuse may be physical, psychological, or both. The most important role for the team physician is the recognition of any potential abuse, with immediate and

appropriate response. The first step is to encourage open disclosure and avoid any suggestive, directing, or leading questions. It is essential to consider that the victim may be experiencing feelings of shame, guilt, and/or fear.^{54,95} Active listening and creating an environment of psychological and emotional support for the athlete wherein they feel safe to discuss events are crucial. Although certain emotions and negative feelings toward the perpetrator may be present, it is important to maintain a neutral tone.⁵⁴ It can be helpful to acknowledge the courage required to speak about abuse that may implicate teammates, coaches, or other seemingly respected members of the community. Finally, it is critical to reinforce to the athlete that the victimization and abuse is not their fault, and that abuse is not a normal, healthy, or helpful part of the team or athletic structure.^{96,97}

Be aware of the policies regarding the duty to report the sexual abuse to authorities depending on the legal statutes in the particular country/jurisdiction. Similarly, depending on privacy laws, providers should involve any and all medical, MH, administrative, and/or legal professionals in a timely manner. Failure to act appropriately may empower the perpetrator(s) and increase the psychological sequelae resulting from the abuse.^{98,99} Ensure accurate record completion and documentation. Despite the conflicting interests from multiple layers of stakeholders and unique pressures faced by members of the sports medicine team, there should be no lack of ethical clarity. The primary and only obligation is to the athlete's well-being.

Effect on Performance

Sexual abuse can have consequences that are psychological and physical in nature. Signs of abuse unique to athletics may also include the excessive taking of risk within their sport, loss of confidence, lack of concentration during participation, declining performance, skipping training sessions, self-injurious behaviors to avoid having to participate in sport, early dropout, excessive training to cope (which leads to burn out), and unexplained injuries that are clinically atypical or fail to resolve.^{78,100} Those who have been sexually abused in the context of sport often have a variety of physical ailments and psychosocial concerns such as low self-esteem, strained relationships with their parents and peers, and higher rates of EDs, all of which can interfere with their athletic abilities.⁷⁸

Prevention

Detailed in Table 3, prevention efforts need to account for the 3 main categories of risk regarding sexual harassment and abuse in sport-athlete, coach, and sport.^{82,89} Several well-established strategies in the protection of minors can also be applied to help prevent sexual abuse in sport. Four key components of the grooming process in a coach-athlete relationship are the targeting of the victim, building trust and friendship, developing control and loyalty, and building and securing secrecy.⁷⁴ Creating early recognition and intervention at each of these levels has the potential to prevent abuse before it occurs. By maintaining a culture with open lines of communication without secrecy and ensuring the presence of adults who will legitimize an athlete's concerns and act on their behalf, grooming can be thwarted.

Criminal background checks should be required for all adults who will be interacting with young athletes, but

TABLE 3. Risk Factors for Sexual Misconduct in Sport.^{82,89}

Athlete Variables	Coach Variables	Sport Variables
Female sex	Male sex	Weak employment controls
Younger age	Older age	No existence of parent and athlete contracts
Smaller physique	Larger physique	Weak or no codes of ethics
High performance status	Good qualifications	Increased opportunities for travel
Low self-esteem	High reputation	Limited opportunities for reporting concerns
Moderate to high medical problems	Unknown or ignored previous record of sexual abuse	
Weak relationship with parents	High trust from parents	
Low awareness of sexual abuse	Low commitment to codes of ethics	
Complete devotion to coach		

additional assessments such as written applications, personal interviews, and reference checks should be instituted as well. An athlete protection policy, including a written “statement of intent that demonstrates a commitment to create a safe and mutually respectful environment,” should be created for every sporting organization.¹⁰¹ Codes of conduct should be included that clearly establish expectations and boundaries of behavior for interactions with those less powerful, especially in terms of physical contact, dependency, and control.⁸² Interventions include increasing sexual abuse awareness and training efforts, establishing policies on rules of behavior and procedures for filing complaints that guarantee prompt attention, disciplinary measures, and follow-up, and providing resources and support for victims.^{74,82,89,101}

Parents should be included in any written or electronic communication between coaching staff and youth athletes. All adult-child interactions, including medical examinations, should be viewed by others at all times. There should also be established policies for adult athletes that patient chaperones should be provided anytime when requested, or necessary, especially during sensitive examinations or procedures, to protect and enhance the athlete’s comfort, safety, privacy, and dignity. Parents should be encouraged to ask organizations for copies of the existing safeguarding policies and procedures in place, including training, reporting, and monitoring. When travel is involved, athletes should use the buddy system, and detailed itineraries should be provided to parents before departure. There should be zero tolerance of behaviors in sport that would not be held as acceptable in other youth environments. The culture can be changed by challenging those who behave or speak inappropriately, and all allegations or suspicions of child abuse or other inappropriate behavior by staff and volunteers must be reported to local law enforcement. Although most studies have looked at sexual misconduct from coaching staff, others who participate in the athlete’s care, including medical personnel, can also be involved in sexual misconduct.

Transition From Sport

Detection in Athletes

Athletes at all levels of competition transition out of sport for numerous reasons, including (1) they have reached the end of their playing career, (2) they can no longer compete at the needed level, (3) they suffer a season- or career-ending or catastrophic injury, (4) the risk-reward ratio may no longer be worth it for them to play, (5) the game is no longer fun, or (6)

they develop other interests.¹⁰² This transition can be a problem for some athletes, whether or not they made the decision to finish competing. As with most events in life, “those who perceive that they are in control of the decisions are less likely to experience a difficult transition than those who perceive they are not.”¹⁰² A recent literature review demonstrated a correlation between voluntary retirement and a less difficult transition out of athletics.¹⁰³

Athletes from youth to elite levels transition from sport every day, and health care providers may want to assess the reasons their athlete is leaving a sport that has played a large role in forming their identity. The British Athletes Lifestyle Assessment Needs and Career and Education scale (BALANCE) and the Transition Coping Questionnaire are available to assist athletic health care providers in recognizing an athlete who may be struggling with transitioning out of sport.¹⁰⁴

Management

Management of the difficulties associated with transitioning from sports will vary depending on the athlete’s symptoms. Psychotherapy may be necessary if the athlete is diagnosed with anxiety or depression due to a loss of athletic identity and demonstrates a need for increased coping skills.¹⁰² Short workshops such as the NCAA Moving On!¹⁰⁵ and the Mind Body and Sport¹⁰⁶ program teach coping and stress management skills to help athletes with this transition. Athletes should be encouraged to begin lifelong activities or join other groups or recreational teams if they are missing or craving the social support they once received from participating in sports.^{102,107} Different treatment methods will work with different patients, and athletes should be encouraged to try multiple alternatives to find the most fulfilling treatment or substitution for them.

Although most athletes transition out of athletics without much difficulty, some will have a much harder time with this process. For injured athletes, the adjustment period from participation to retirement has been reported to range from 6 months to 1 year.^{103,107} Athletes who sustain career-ending injuries report lower life satisfaction for 5 to 10 years after retirement.¹⁰⁸ Mental health disorders, negative coping skills such as substance use, and criminal activity have all been reported after athletic retirement.^{102,109} These negative coping strategies may continue throughout retirement. Research shows greater adjustment difficulties, lack of retirement planning, and more frequent and severe psychological issues in athletes with higher levels of athletic identity.¹⁰³ The athletic care network should be aware that

a difficult transition may occur and when possible educate and observe individuals in the months after retirement.

It is worth noting that positivity can come from this transition. Many athletes see the value in getting involved in other activities; those who spent more time on themselves, their education, and their future had a more balanced life.¹⁰⁷ Conversely, athletes who focus too much on sport may have difficulty with helplessness and feeling lost upon retirement.¹¹⁰ These findings support the idea of a multidimensional identity as being a strong protective factor against the negative effects of retirement from sport. In addition, participants with a broader self-identity reported an easier transition after an injury than those participants who had a strong athletic identity.¹⁰⁷

Prevention

A preretirement plan to assist athletes with issues surrounding the transition out of athletic participation may be helpful.¹⁰³ Having a life development intervention plan in place may mitigate some of the negative fallout surrounding an early retirement.¹⁰⁴ There is evidence that suggests transitioning out of sport can be effectively managed at the professional and/or national levels.^{104,111} Participation in a psychoeducational retirement planning intervention focused on diversifying athletic identity, enhancing coping skills, building social support, and initiating the grief process earlier may create a more positive athletic transition process for the athlete.¹⁰³

THE PSYCHOLOGICAL RESPONSE TO INJURY AND ILLNESS

Relationship Between Injury, Performance, and Mental Health

The relationship between injury, performance, and MH in elite sport is complex and an area of interest.^{4,5,112–115} Elite sport brings specific stressors that potentially increase the likelihood of injury/illness, including MH disorders, and injury can potentially unmask and/or trigger MH disorders.^{116–122} Finally, there may be MH disorders that increase the likelihood of, or complicate recovery from, injury.^{116–119} There is minimal prospective research in this area with several, mostly unanswered, questions.

Detection in Athletes

Certain physical and MH disorders may be more common in athletes compared with their nonathlete peers, including performance anxiety, disordered eating (DE), menstrual dysfunction, and binge drinking.^{123–125} The psychological response to injury may trigger and/or unmask MH disorders including depression/suicide, anxiety, gambling, DE/EDs, and substance use/substance use disorders.^{126–138}

Certain psychological and sociocultural factors have been raised as potential risk factors for injury.^{120,121,139} Although there are limited prospective and/or conflicting results regarding some of these risk factors, stress consistently demonstrates a relationship with injury risk as well as with the ability to rehabilitate from injury and return to sport.^{116,139–142} A meta-analysis found that high stress response and history of negative stressors had the strongest association with injury rates.¹¹⁶ Perceived negative life event

stress from teammates and the coach was associated with an increased risk of acute and overuse injuries.¹⁴¹ Another study of team-sport athletes found an association between negative perceived recovery and the risk of both acute and overuse injuries.¹⁴² Emotional reactivity and stressful life events are also associated with poor on-field performance and injury.^{117,143,144} Athletes with high resiliency, self-efficacy, optimism, and a supportive social network respond more effectively to stress.^{118,145–148}

Response to Injury

A systematic review looking at psychosocial factors associated with outcome in sport injury rehabilitation in elite athletes concluded that 3 athlete responses were most important: cognitions, emotions, and behaviors.¹³⁹ The cognitive response, or how the athlete interprets the injury/illness, may determine an emotional response, which may then affect an athlete's behavioral response (eg motivation, goal setting, and compliance with treatment) as well as response to treatment. The cognitive and emotional responses to injury and illness can be considered either "normal" or "problematic."

Examples of problematic cognitive responses include concerns about reinjury ("I can't do this as it might increase my risk for injury"), doubts about competency ("I can't play this sport"), self-efficacy ("I won't be good enough to start anymore"), loss of identity ("Who am I if I'm not an athlete"), or concerns about the medical staff ("I don't think my medical staff is capable"). When an athlete is injured, one of the first responses is coping with and processing the medical information provided.

The emotional response to illness and injury includes sadness, isolation, lack of motivation, anger, irritation, frustration, changes in appetite and sleep, and disengagement. Injured athletes report a higher level of symptoms of depression and generalized anxiety disorder (GAD) compared with noninjured athletes.¹⁴⁹ The response of each person may differ, and the response may also vary within the same individual depending on the circumstances. Emotional responses to injury and illness can be normal. Problematic emotional responses are those that do not resolve, worsen over time, or in which the severity of symptoms seem excessive.⁴

Management

Resiliency and "high mental toughness" are associated with lower injury rates; those athletes with higher resiliency have a lower incidence of depression, anxiety, stress, and obsessive-compulsive symptoms.^{150,151} Psychological aspects of recovery and rehabilitation can influence an athlete's perception of their injury, life outside of sport, expectation for long-term outcome, and satisfaction with health care providers.^{119,152–156} Prominent individual factors may impact outcome, including pain perception, optimism/self-efficacy, and depression/stress.^{119,152–156}

A systematic review identified 3 psychological elements (self-determination theory—autonomy, competence, and relatedness) as the factors most important in positive rehabilitation and return to preinjury level of play.¹¹⁹ Examples of strategies that support positive return to sport experiences include (1) reducing reinjury anxieties using modeling techniques, (2) building confidence using functional tests and goal setting, (3) providing social support, (4) keeping athletes involved in their

sport, (5) reducing stressors related to premature return to sport, and (6) foster athlete autonomy.¹⁵⁵

Self-Medication in Response to Injury/Illness

Detection in Athletes

Participation in sports can include a greater risk of experiencing injury-related pain and the loss of one's core identity due to earlier than expected retirement.¹⁵⁷ An athlete may seek unhealthy methods of coping, including self-medication. Determining that an athlete has a problem with self-medication can be difficult, but evidence can include uncharacteristic behavioral changes such as arriving for practice too early or too late, missing training altogether, or frequent heightened conflict with teammates.

Although unable to distinguish purely social consumption from self-medication, a recent study by the NCAA sheds some light on the status of student-athlete substance use.¹⁵⁸ Overall, around 80% of student-athletes reported alcohol use in the past year, similar to rates for same aged nonathletes.¹⁵⁸ The percentage of student-athletes drinking excessively has dropped over the past decade for both male athletes (63%-44%) and female athletes (41%-33%).¹⁵⁸ It also showed that the use of recreational drugs such as tobacco, marijuana, and cocaine by student-athletes is generally less than their nonathlete college peers, with 22% of student-athletes reporting using marijuana compared with 33% of the general student body.¹⁵⁸ Cocaine use in high school sports peaked in the mid-1980s at 17%, but sharply fell to less than 2% in the 1990s.¹⁵⁹

Cannabis is viewed by many users as a way to experience anesthetic or relaxing effects and as such is one of the most commonly used agents by athletes to self-medicate.¹⁵⁹ Contrary to use patterns in the general population, among athletes, marijuana seems to have taken the place of tobacco as the second most widely used drug after alcohol.¹⁶⁰ Given the association between the 3 variables of marijuana use, depression, and body image-related stress in athletes, this pattern can be exacerbated after illness or injury.¹⁶¹⁻¹⁶³

Perceived attempts at self-medication have been demonstrated in a survey of collegiate athletes, where a significant relationship exists between reported alcohol abuse and self-reported symptoms of depression and general psychiatric symptoms.¹²⁷ Athletes who affirmed ratings in the "severe" range for depression and other psychiatric symptoms had a significantly higher rate of alcohol abuse than those who had low depression scores and low or mild symptom ratings.¹²⁷ Conversely, subjects reporting higher rates of alcohol abuse also had more psychiatric symptoms.¹²⁷ Nevertheless, only 0.6% of NCAA athletes reported using alcohol to deal with the stress of athletics.¹⁶⁴ A possible explanation for this disparity is that sport-related coping motives contribute to excessive substance use among athletes, without athletes recognizing or explicitly identifying coping-related intentions.¹⁶⁵ However, athletes who acknowledged alcohol use as a coping mechanism also experienced more negative consequences as a result of that drinking.¹⁶⁵

Opioid use disorder involving prescription opioids is recognized as one of the most important health problems today in medicine and public health.¹⁶⁶ The use and misuse of opioids crosses all levels of competition from teens to professionals. Approximately 23% of NCAA student-

athletes used prescription pain medication in the past year, with an additional 6% reporting use of these substances without an actual prescription.¹⁵⁸ Meanwhile, high school student-athletes who are involved in one competitive sport are at a greater risk of being prescribed, misusing, and being approached to divert opioid medications.^{167,168} Student-athletes who played more than 3 sports were 3 times more likely than nonsport participants to be approached to divert their opioid medications.¹⁶⁹ Diversion of controlled substances has been found to be associated with other forms of prescription drug misuse.¹⁷⁰ A study of opioid use among retired professional football players found that of the 52% who were prescribed opioids during their career, 71% misused their prescriptions; of these, 15% admitted to ongoing misuse, with pain, undiagnosed/unreported concussions, and heavy drinking all predicting current misuse.¹⁷¹ An additional concern is that 63% reported getting prescription opioids from nonmedical sources during their playing days.¹⁷¹

Male adolescents participating in organized sports are more likely to be prescribed, use, and misuse opioids compared with their nonsport peers.^{160,167} Although most sports in which adolescents participate during high school are not associated with the use of heroin or nonmedical use of prescription opioids (NUPO), certain high-contact sports such as wrestling and football show a greater risk for engagement in NUPO, with ice hockey players having an increased risk for concurrent heroin use and NUPO.¹⁷² It is believed that injury may be a key driver in this association; however, normative behaviors among some athletes or the stress associated with high-level competition may also play a role in this greater risk.¹⁷²

Many athletes do not recognize prescription opioids as a potential health risk and are known to self-medicate without consulting a physician first.¹⁷³ One third of NCAA athletes believe there is nothing wrong with using "painkillers" to cope with pain associated with competition.¹⁷⁴ However, it has been demonstrated that prescription opioid misuse is a strong risk factor for heroin use.¹⁷⁵ The need for pain control after surgery or in the management of a variety of musculoskeletal-related injuries is an issue regularly faced by the athlete. Postoperative exposure to opioids correlates with long-term use and use disorders, and therefore, the need to use other strategies of pain management is of critical importance.¹⁷⁶

Management

A multidisciplinary approach involving both team physicians and MH providers using a variety of psychotherapeutic, psychosocial, and, if indicated, pharmacological interventions should be used. Management should span the spectrum from acute withdrawal syndromes to long-term addiction treatment services.

No clinical trials currently exist for the specific treatment of substance use disorders in athletes. Therefore, proven strategies from the general population should be used that include addressing the underlying issues leading to substance use and the health problems caused by pain during and after an athletic career.^{159,160} Health care providers should recognize that some medications used in cocaine and opioid dependence treatment, such as modafinil and methadone, are banned substances in sport.¹⁵⁹

Health care providers should also have a general understanding of the differences that often exist between athletes

and nonathletes because this awareness improves not only relatability but effectiveness of interventions. A social norms-based intervention is likely more effective when other athletes rather than students in the general population are used as the reference group, especially as research suggests that perceived alcohol consumption among one’s close friends who are athletes is a very strong predictor of personal consumption.¹⁵⁷ Targeted interventions that incorporate health and athletic performance considerations tend to resonate and be more successful for the athletic population as well.¹⁵⁷

Prevention

Like treatment, prevention must be comprehensive and systemic. Use of relaxation training, adequate social support, psychological counseling, and other nonmedication-based pain control/stress management interventions should be promoted as healthier options for coping. Education is a key component of prevention, and the documented risks and alleged benefits of substance use should be openly and honestly discussed, including as a routine topic at health maintenance and preparticipation physicals. There is also a clear need for depression and anxiety screening, especially after an athlete has experienced a challenging illness or injury, to detect conditions that may lead to self-medication. Using the idea of “pain” as the “fifth vital sign” may have contributed to opioid overprescribing, and therefore, its use is no longer advocated by major US medical organizations.¹⁷⁷ Yet it remains important to understand and manage an athlete’s pain with open lines of communication and standardized, validated reporting measures, to prevent self-medication. Direct questioning should be revisited if any evidence of substance use disorders, legal issues, or conduct issues arise. In addition, 60% of college student-athletes felt that drug testing was an effective deterrent for them, and their peers and should be continued.¹⁵⁸

Administrators and coaches should promote clear and consistent team-based policies on substance use, and the inclusion of team leaders in the development of policy and the promotion of expectations to other team members has yielded a reduction in negative consequences from substance use as well as a positive impact on team performance.¹⁵⁷ Opioid overprescription should be avoided, and alternative pain management solutions should be considered. In addition, based on diversion data, athletes should be educated on the proper use and disposal of opioid medications.¹⁶⁹

SELECT MENTAL HEALTH AND PSYCHOLOGICAL ISSUES

Eating Disorders/Disordered Eating

Detection in Athletes

The diagnosis of EDs follows the DSM-5 guidelines.¹⁷⁸ Athletes demonstrate increased prevalence of ED and DE behaviors compared with nonathletes across all age groups and genders.¹⁷⁹ Because of the harmful health and performance effects, it is important to identify those athletes at increased risk of DE as well as those with clinical diagnoses of ED.

Two conditions that exist along the spectrum of EDs and DE include the Female Athlete Triad and Relative Energy Deficiency in Sport. The Female Athlete Triad has been

described as the relationship between energy availability, menstrual function, and bone health.¹⁸⁰ Relative Energy Deficiency in Sport relates the impact of energy deficiency on physiological function including, but not limited to, metabolic rate, menstrual function, bone health, immunity, protein synthesis, and cardiovascular health.¹⁸¹

Athletes at increased risk of ED include those who participate in weight-restricted and aesthetic sports, as well as sports where having a low body weight is seen as an advantage. In addition, female athletes have been shown to have different factors contributing to onset of ED compared with nonathletes.¹⁸² Although both groups had increased risk with low self-worth, peer issues, and comorbid psychiatric disorders, the athletes had sport-specific factors such as performance pressure, team weigh-ins, and injuries.¹⁸² Some characteristics such as perfectionism are seen in the general population as being an increased risk for DE; in athletes, perfectionism may be problematic but can also be beneficial in sport, thereby complicating management of that risk factor.

Several effective screening tools have been developed to identify EDs specific to athlete populations: Athletic Milieu Direct Questionnaire, Brief Eating Disorders in Athletes Questionnaire, Compulsive Exercise Test—Athlete Version, Female Athlete Screening Tool, and Psychologic Screening Test.^{183,184} These screening tools were all developed using DSM-IV criteria and were validated in female athlete populations; however, they may not accurately reflect today’s diagnostic standards and may not translate well into screening male athletes. There is significantly less literature surrounding male athlete EDs, and there may be different characteristics or issues that a female-based screening tool would miss. Although often used in real-world clinical practice as ED screening tools, the SCOFF¹⁸⁵ and the ED Examination interview (EDE-16)¹⁸⁶ are not athlete-specific.

Current preparticipation screening recommendations vary. The Preparticipation Physical Evaluation Monograph, fourth edition, asks 4 gender-neutral questions regarding DE, 3 regarding menstrual history, and 1 regarding history of stress injuries, while the 2014 female athlete triad coalition consensus statement adds 3 additional questions (Table 4).^{180,187}

TABLE 4. Screening Questions for Eating Disorders/Disordered Eating

Questions common to both PPE-4 and the Female Athlete Triad Coalition Consensus
Do you worry about your weight?
Are you trying to or has anyone recommended that you gain or lose weight?
Are you on a special diet or do you avoid certain types of foods?
Have you ever had an eating disorder?
Have you ever had a menstrual period?
How old were you when you had your first menstrual period?
How many periods have you had in the past 12 months?
Have you ever had a stress fracture?
Female athlete triad screening questions
When was your most recent menstrual period?
Are you presently taking any female hormones (estrogen, progesterone, and birth control pills)?
Have you ever been told you have a low bone density (osteopenia or osteoporosis)?

To date, there have been no studies that identify the ideal time to screen. Considerations would include how young to start screening, how frequently to screen, and at what time point(s) relative to a sport season (ie, preseason, during season including at pre-established time points or during routine visits for injuries/illness, or postseason) an athlete should be screened.

Management

The goal of ED management in athletes is to empower the athlete to be able to safely perform as much of their sport or physical activity as possible while taking the steps to undergo treatment. Eating disorder and DE behaviors often have deep roots with multiple comorbidities. The care of an athlete with DE is a long-term process that requires careful monitoring.

Sport participation criteria are outlined in the 2014 Triad Coalition Consensus Statement.¹⁸⁰ Risk factors such as energy availability, body mass index (BMI), menstrual history, bone mineral density, and bony stress injuries are evaluated to provide a cumulative risk stratification to help determine clearance and return-to-play (RTP) guidelines. Relative Energy Deficiency in Sport also presents a similar model for risk assessment (Table 5).¹⁸¹ The Coalition statement includes a decision-based RTP model (Figure 1)¹⁸⁰ that considers medical factors, sport risk modifiers, and other decision modifiers when determining RTP.

Risk stratification, physical examination, laboratory workup, and electrocardiogram (ECG) can help to determine whether the athlete can continue with any level of sport participation, as well as identify those who should be managed at a higher level of care due to increased risk. Not all athletes are safe to manage as outpatients; some require increased levels of care, including intensive outpatient, partial hospital, residential, and inpatient care. Ultimately, these decisions are

based on the athlete’s medical and psychiatric stability. Importantly, an endurance athlete’s low resting heart rate may not be the same indicator of depressed vital signs because it would be in a nonathlete. Unstable vital signs or inappropriate response to exercise and recovery may be more useful to measure in an athlete. Athletes who are suicidal and unable to contract for safety should be managed in an inpatient setting.

The treatment of ED in athletes should involve a multidisciplinary team including a team physician, nutritionist, and MH care provider, preferably all with experience in taking care of athletes. If an athlete is part of a team or setting that includes a certified AT, that AT is a key member of the athletic care network who often has closer, regular interaction with the athlete, and a better understanding of team and coach dynamics. An athlete’s support team may also include parents, teammates, coaches, academic advisors, and others.

The initial goal of treatment, particularly for those who are underweight, is weight restoration aiming for at least 90% of ideal body weight. Psychotherapy has been found to be beneficial in treatment of EDs, with cognitive behavioral therapy (CBT), the primary modality for treatment. However, in a study of adolescents still at home, family-based therapy (Maudsley family therapy) was just as effective as individual therapy by the end of treatment and demonstrated a greater persistence effect over time.¹⁸⁸ Newer studies are looking at electronic CBT programs, which would help eliminate potential time and privacy barriers associated with attending counseling in person. However, the evidence on their effectiveness is currently equivocal.¹⁸⁹

The Food and Drug Administration (FDA) has approved fluoxetine for bulimia nervosa and lisdexamfetamine for binge ED because there is some evidence to support their use for those conditions.¹⁹⁰ Importantly, stimulants such as lisdexamfetamine are banned in higher levels of sport without

High risk: no start red light	Moderate risk: caution yellow light	Low risk: green light
<ul style="list-style-type: none"> ▶ Anorexia nervosa and other serious eating disorders ▶ Other serious medical (psychological and physiological) conditions related to low energy availability ▶ Extreme weight loss techniques leading to dehydration induced haemodynamic instability and other life-threatening conditions 	<ul style="list-style-type: none"> ▶ Prolonged abnormally low % body fat measured by DXA or anthropometry using The International Society for the Advancement of Kinanthropometry ISAK¹⁴¹ or non-ISAK approaches¹⁴² ▶ Substantial weight loss (5–10% body mass in 1 month) ▶ Attenuation of expected growth and development in adolescent athlete ▶ Abnormal menstrual cycle: FHA amenorrhoea >6 months ▶ Menarche >16 years ▶ Abnormal hormonal profile in men ▶ Reduced BMD (either from last measurement or Z-score < -1 SD). ▶ History of 1 or more stress fractures associated with hormonal/menstrual dysfunction and/or low EA ▶ Athletes with physical/psychological complications related to low EA/ disordered eating - ECG abnormalities- Laboratory abnormalities ▶ Prolonged relative energy deficiency ▶ Disordered eating behaviour negatively affecting other team members ▶ Lack of progress in treatment and/or non-compliance 	<ul style="list-style-type: none"> ▶ Healthy eating habits with appropriate energy availability ▶ Normal hormonal and metabolic function ▶ Healthy BMD as expected for sport, age and ethnicity ▶ Healthy musculoskeletal system
<p><small>BMD, bone mineral density; DXA, dual-energy X-ray absorptiometry; EA, energy availability; FHA, functional hypothalamic amenorrhoea; ISAK, International Society for the Advancement of Kinanthropometry</small></p>		
<p><small>Mountjoy M, et al. Br J Sports Med 2014; 48:491–497. Used with permission.</small></p>		




Figure 1. Decision-based RTP model for the female athlete triad. *Return-to-play decision is determined by the primary care or team physician and is based on a complex and comprehensive synthesis of health status, cumulative risk assessment, participation risk, sport and decision modifiers. 25(OH) Vit D, 25-hydroxyvitamin D; BP, blood pressure; CBC, complete blood count; DXA, dual-energy X-ray absorptiometry; TFTs, thyroid function tests; TSH, thyroid stimulating hormone [De Souza MJ, et al. 2014 female Athlete Triad Coalition Consensus Statement on Treatment and return to play of the female Athlete Triad: 1st International Conference held in San Francisco, California, May 2012, and 2nd International Conference held in Indianapolis, Indiana, May 2013. *Br J Sports Med* 2014;48:289. Used with permission].

appropriate documentation and/or therapeutic use exemptions (TUEs). In situations with comorbid MH diagnoses, psychotropic medications may have benefit in treatment of the comorbid conditions.

Exercise is beneficial in many MH disorders, and a review on exercise as treatment in ED demonstrated benefit in some individuals.¹⁹¹ If continued exercise is felt safe for a given athlete, then clear guidelines, close monitoring, and possibly written contracts are suggested. This may be a way to empower a patient with DE to commit to and participate in their treatment program.

Effect on Performance

Some athletes and coaches feel that calorie restriction and weight loss can improve sport performance. This is particularly prevalent in endurance sports such as cross-country and cycling. There are 2 theories that support an initial short-term improvement in sport performance with weight loss, including upregulation of the hypothalamic–pituitary–adrenal axis and potential increased maximum oxygen uptake.¹⁹² However, long-term continuation of calorie restriction leads to progressive deterioration of sport performance.¹⁹³

The complications of DE can range from low energy to dehydration to potential electrolyte abnormalities. All of these can have negative impacts on sport performance.¹⁹⁴ Fogelholm outlined 4 pathways of reduced performance¹⁹³: (1) Glycogen depletion can cause reduced physical and psychological capacity; (2) increased circulating lactate can produce increased muscular pain with exercise; (3) dehydration can limit performance through early fatigue, as well as through increased risk of muscle cramps and heatstroke; and (4) loss of lean mass can lead to decreased muscle strength and aerobic performance.

Athletes who engage in excessive compulsive exercising may exercise more than they can recover and are at risk of overtraining syndrome (OTS). As athletic performance declines, this can exacerbate the psychological stressors leading to the excessive training in the first place. In a study of high school athletes with EDs, those with EDs were twice as likely to sustain a musculoskeletal injury. In another study that classified athletes according to risk category using the 2014 Consensus Statement, female athletes with moderate risk were twice as likely to sustain a bone stress injury compared with those classified as low risk, and high-risk athletes were 4 times as likely to sustain a stress injury compared with low-risk athletes.¹⁹⁵ The direct impact on sport performance and the indirect impact caused by injuries leading to time-loss from sport are important consequences of EDs in athletes.

Prevention

Several studies have evaluated the impact of ED prevention interventions, with most showing a reduction in risk factors and decreased behaviors associated with DE. However, only one study evaluated the outcome of new ED diagnoses following a prevention program.¹⁹⁶ The 1-year intervention was based on a social-cognitive framework, with a focus on enhancing self-esteem by strengthening self-efficacy. The 9-month follow-up showed no new cases of EDs in the intervention group, while 13% of female athletes in the control group developed EDs diagnosed by DSM-IV. Although the authors evaluated female and male athletes, the data were significant only in female athletes.

Other programs that have shown reduced behaviors of DE include peer-led educational programs, education targeted at coaches, and electronic programs.^{189,196,197} Future research on prevention programs should focus on diagnostic outcomes

and evaluate optimal timing to initiate interventions and retention of behavior improvements.

Depression and Suicide

Depression

Detection in Athletes

There are minimal data specific to athletes in terms of the prevalence of depression and suicide. Depression rates in college athletes are likely similar to their nonathlete peers in college populations. A total of 23.7% of college athletes reported clinically significant depressive symptoms over a 5-year period using a validated depression screening tool (the Center for Epidemiological Studies Depression scale) in the largest published study to date in the sports medicine literature.¹⁹⁸

Female athletes are at higher risk for reporting depression symptoms than male athletes in the general population and in the limited studies in athletes. In the available studies of depression in college athletes by gender, female athletes are twice as likely to report clinically significant depression symptoms compared with male athletes. Freshman college athletes seem to be more likely to report depressive symptoms when compared with upper classmen.¹⁹⁹

Other than age and gender, there are risk factors for depression that are more unique to athletes and may increase the likelihood of depression. Athletes can face expectations, both external and internal, that their nonathlete counterparts do not. These unique risk factors can be grouped into 4 categories: (1) an injury causing time away from sport (both competition and practice), (2) the time demands on student-athletes (up to 40 hours per week can be spent in athletic commitments for a college athlete), (3) performance expectations (self-imposed or imposed by others such as coaches, peers, and family), and (4) an identity intrinsically linked to athletic participation.¹³⁸

The type of sport may also be a factor in likelihood of reporting depression symptoms. In one large study, it was found that track and field college athletes reported a statistically significant higher rate of clinical depression symptoms than those in team-based sports. The authors noted that in an individual-based sport such as track, there is only one winner in any given event, whereas in a team-based sport, half the participants are winners.¹⁹⁸

The signs and symptoms of depression may be atypical in athletes. For example, a report of “lack of focus” in the athlete may not be a “lack of interest” in the sport but rather a sign of cognitive functional impairment resulting from depression. Anger, insomnia or hypersomnia, fatigue, or a change from previous level of function or behavior at home or school are other examples of signs and symptoms that may present in athletes. Health care providers, coaching staff, and families should watch for types of coping mechanisms used by athletes, as some, such as substance use, are less healthy than others.

A personal history and/or a family history of depression and other mood disorders are important in the evaluation of an athlete with possible depression, in addition to evaluation of the risk factors noted previously. A good history of previous injuries, especially injuries that caused loss of time from sport, can be helpful; an assessment of how the injuries impacted the individual as an athlete is also important. A history of previous

concussions may be a possible risk for depression later in life as noted in a study of retired National Football League (NFL) athletes, but this has not been reproduced yet in any other population as a unique risk factor.¹²⁹

Screening tools for depression—such as the Centers for Epidemiological Studies Depression, the Beck Depression Inventory, and the Patient Health Questionnaire 2 and 9 (PHQ2 and PHQ9)—have all been validated in a nonathlete population.²⁰⁰ These tools do not make the diagnosis of depression but instead are useful tools to identify those at risk for depression based on the clinically relevant depressive symptoms that are reported. None of these screening tools have been validated in an athletic population.

Management

One of the first steps in treatment is early recognition of depression in athletes. Health care providers should recognize that a concern about “performance,” whether reported by an athlete or a coach, might in fact be a symptom of an underlying clinical condition such as depression or anxiety. Unrecognized or undertreated depression can also interfere with optimal recovery from an injury or other illness and can delay return to play.¹¹⁹ For the athlete with depression, there should be a multidisciplinary team approach; the team may include an MH care provider, ideally one familiar with athletes, a team physician, an AT, and with permission of the athlete, the family, and coaching and other professional staff.

When considering prescription medication in the management of depression in an athlete, there are several factors to consider that are more unique to an athletic population. The prescribing clinician and the athletic health care team should be aware if a medication might have a potential negative impact on performance. The health care provider and team should also be familiar of any side effects that may place an athlete at increased risk for injury, illness, or adverse outcome. Weight gain, sedation, and adverse cardiac effects are potential consequences of antidepressant medications. In addition, the health care team should also be aware of any performance-enhancing effects and the classes of medications banned by governing athletic bodies, and the need for specific documentation regarding such medications. As a general principle, a conservative approach to prescribing medication for depression is preferable in the athlete with depression.

In an athlete with depression without comorbid anxiety, if the decision to prescribe a medication has been made, bupropion can be considered. An alternative for a medication to assist in depression management in an athlete is fluoxetine; this choice may be helpful if there is a comorbid ED. There is some research that supports that fluoxetine does not have a negative impact on performance²⁰¹; other selective serotonin reuptake inhibitors (SSRIs) may be reasonable options as well. If there is an anxiety comorbidity in an athlete, the health care provider could also consider SSRIs, such as escitalopram, sertraline, or fluoxetine.²⁰¹

Nonpharmacological treatment can play a critical role in the management of depression. Cognitive behavioral therapy, acceptance-based therapy, and mindfulness approaches have all been used to treat athletes. None have been shown to be more effective than the others in athletes.¹³⁸ The non-pharmacological approach, such as the pharmacological approach, should be tailored to the individual athlete and their particular clinical situation.

Effect on Performance

Significant functional impairment can result from depression in athletes. Although not well studied in the sports medicine literature to date, if motor or cognitive dysfunction is present, this could result in decreased reaction times or coordination being affected. If the athlete is injured, their depression can lead to poor adherence to the rehabilitation program and thus a slower return to actual competition. A depressed athlete may be less engaged in their sport due to a loss of pleasure in activities they previously enjoyed. Suicidality may be increased in the depressed athlete. Depression may lead to a disruption in important relationships such as those with family, teammates, and coaches. In addition, depression can cause other impairments such as sleep disturbance, which can negatively impact performance.

Prevention

Early recognition may provide an opportunity for prevention of more serious effects of depression. Health care providers should consider baseline evaluation of the athlete population using a validated screening tool. Consider early detection efforts in subpopulations such as those with known risk factors, for example, through follow-up depression screening after an injury that causes time lost from sport, and in first year athletes. Minimizing or eliminating depression risk factors is an important part of a prevention-based approach. These can include injury prevention programs, early identification and treatment for sports-related concussions, peer support groups, and education of peers, coaches, and staff to recognize signs and symptoms of depression in athletes.

Suicide

Detection in Athletes

In a study of NCAA college athletes and suicide, over a 9-year period, there were 35 deaths from suicide identified from a total of 477 student-athletes' deaths and 3.7 million participant seasons.²⁰² Suicide represented 7.3% of all-cause mortality for the NCAA student-athletes, and the overall suicide rate for these athletes was 0.93/100 000 per year. This is lower than the suicide rate of 7.5/100 000 among all college students.²⁰² In the same study, the suicide incidence was higher in male athletes than female athletes (1.35/100 000 vs 0.37/100 000). Of note, football players had the highest rate by sport, 2.25/100 000, and a relative risk of suicide of 2.2, compared with other male nonfootball athletes. A study of retired NFL players, spanning a 50-year period, also found a lower than expected rate of suicide compared with a matched population. There were 12 suicide deaths in this retired NFL population compared with an expected number of 25 over the same period.²⁰³

The interplay between participating in athletics, depression, and suicide is complex and not well understood. For example, although exercise is an accepted form of treatment for depression, college athletes are likely to report depressive symptoms at similar rates as their nonathlete peers.¹⁹⁸ This may be because athletes have unique risk factors for depression as a result of their athletic pursuits. Depression increases the risk for suicide, but it is unclear whether there is a cause and effect relationship rather than an associative one. Furthermore, participating in athletics in college may provide

peer and other support for the individual—a good thing—but at the same time, an athlete may be more willing to engage in high-risk behavior with peers, which could contribute to depression and suicide risk.¹³⁸

Management/Prevention

Treatment is focused on prevention and early identification and intervention, as with depression. Awareness of suicide in athletes and educational programs aimed at athletes, health care providers, coaches, staff, and families are critical for identification of the athlete at risk and appropriate intervention. Many universities are proactively addressing suicide risk in their communities. A public health approach to these preventable deaths can be appropriate in the athletic population. Organizations such as the NCAA are developing educational tools and resources to help their member institutions with preventive approaches to suicide in student-athletes. As with depression, a multidisciplinary care team approach is key to the management of suicidality in athletes, early identification of those at risk, and implementation of proactive prevention programs.

Anxiety and Stress

Detection in Athletes

Anxiety has the potential to enhance or hinder athletic performance. Because some level of anxiety may be normative to sports and competitive environments, it may be difficult to detect the behavioral, physical, and cognitive symptoms of pathologic anxiety in athletes. There are no known validated athlete-specific anxiety screening tools. The GAD-7 is a screening tool that has been used and validated in the general population.²⁰⁴ Some studies using general population screening tools to investigate prevalence of different anxiety disorders in athletes have suggested a slightly higher prevalence of many forms of anxiety in high-level athletes. Among the most relevant for athletes are panic disorder, GAD, social anxiety disorder, and specific phobias as well as anxiety-related disorders of post-traumatic stress disorder (PTSD) and obsessive-compulsive disorder (OCD).

Generalized anxiety disorder has dominant cognitive symptoms that may result in the inability of the athlete to be cognitively present or mindful of the moment at hand. Generalized anxiety disorder occurs in approximately 3.1% of the general population in a given year.²⁰⁵ Although there are minimal prevalence data in athletes, 6 percent of elite French athletes had clinically significant GAD symptoms,²⁰⁶ and 7.1% of Australian athletes had clinically elevated GAD symptoms.¹⁴⁹

Panic disorder may result in athletes being hypersensitive to their own physiological sensations, such as increases in heart rate, and may lead to avoidance of situations that may cause panic attacks, such as a performance or competition. The prevalence of panic disorder is 2.7% in the adult populations²⁰⁵ and is likely to be similar in athletes, with lifetime prevalence of panic disorder of 2.8% as reported in the elite French athlete population.²⁰⁶

In athletes, social anxiety may involve the avoidance of a situation in which one may have difficulty being able to escape, for example, practice, competition, social interactions, team meetings and meals, and services in the athletic training room. Trait levels of social anxiety were related to social

evaluative fears in sport, and social anxiety was positively correlated with avoidance of individual sports but not team sports.²⁰⁷ Lifetime prevalence rate for social anxiety disorder in the United States is estimated at 13%.²⁰⁸ There are no reliable estimates of social anxiety disorder in athletes; however, clinically significant social anxiety symptom rates as high as 22.2% in a male collegiate athlete population and 37.3% in a female collegiate athlete population have been reported.²⁰⁹

In athletes, OCD may involve intrusive thoughts and rigid use of routines that cause impairment. Prevalence of OCD is estimated at 1.6%²⁰⁵ in the general population and 5.2% of a US college-athlete population met full OCD criteria.²¹⁰ Post-traumatic stress disorder in athletes involves physiological symptoms and avoidance of athletic situations or triggers that are associated with the trauma experience. There is a PTSD prevalence of 6.8%²⁰⁵ in the general population. Prevalence of PTSD in athletes is unknown; however, case studies of PTSD in athletes after injury^{211,212} suggest that postinjury PTSD does occur in athletes.

The athletic environment contains multiple risks for anxiety dysfunction. The most common and basic anxiety trigger is athletic competition and fear of failure, or perceived negative consequences of an athletic performance. Social judgment is also a risk factor for anxiety in athletes. Crowd effects, coach or teammate judgment, ambiguous scoring procedures, and any situation in which others judge an athlete have the potential for a heightened anxiety reaction. In addition, higher numbers of past severe injuries, past surgeries, and recent life events, higher levels of career dissatisfaction, and lower levels of social support were related to the occurrence of anxiety symptoms among both current and former elite athletes.²¹³

Management

Some psychotropic medications may be used in athletes to treat anxiety; however, as with medications for depression, caution is noted to understand any potential negative impact on athletic performance, potential athletic performance-enhancing effects, and potential safety risks.²¹⁴ The use of SSRIs, such as fluoxetine, is recommended for athletes, based on preliminary evidence that buspirone may interfere with performance, and lack of such evidence for SSRIs.²¹⁵ In general, as-needed anxiolytics are not recommended for athletic performance anxiety.²¹⁶ Beta-blockers can be either performance enhancing or performance inhibiting and have been banned in various Olympic sports because of their ability to improve fine motor control.²¹⁷ In an International Society for Sports Psychiatry membership survey, the most commonly prescribed medication for anxiety by sport psychiatrists was the SSRI escitalopram.²⁰¹ International Society for Sports Psychiatry members previously reported that they avoided benzodiazepines because of sedation, dependence, impaired reflexes and balance, and cognitive impairment.²¹⁸

Cognitive behavioral therapy for the treatment of anxiety is the primary nonpharmacological intervention. Although there are no randomized controlled trials (RCTs) of CBT interventions specifically within athletes, CBT is well established as an effective treatment method within many other clinical populations.²¹⁹ These interventions generally include a combination of cognitive interventions and exposure to anxiety stimuli and should be delivered by a qualified CBT clinician who has experience providing services to athlete

populations. Additional adjunctive treatments such as exercise, sleep management, mindfulness training, and diet change may also be considered in the context of pharmacological and cognitive behavioral interventions.

Effect on Performance

Anxiety has the potential to both enhance and hinder athletic performance based on the nature of the athletic task, appraisal of the anxiety by the athlete, and the physiological learning history of the athlete. The Yerkes–Dodson law states that an increasing level of anxious arousal has the potential to enhance performance up to a point, beyond which performance is decreased.²²⁰ The behavioral, physical, and cognitive symptoms of anxiety may present particular symptom profiles that may hinder performance in unique ways. As the athlete attempts to control relevant thoughts, feelings, and body sensations (eg, increased heart rate or respiration rate associated with athletic performance), the task of controlling anxiety may interfere with sport engagement. The result can be a cycle in which performance declines and distress increases. Specific implications of these problems often vary according to the forms of anxiety (eg, panic, worry, etc.) that the athlete experiences. For example, athletes with GAD are likely to have performance difficulties or fail to perform to their potential due to continued worry and increased non-task-related cognitive activity; this may then hinder a state of flow or the ability to engage effectively over the course of a performance task. Athletes with panic disorder may miscalculate physiological responses that are a natural byproduct of their athletic task, and thus avoid performance situations.

Athletes with social anxiety disorder experience a shift of cognitive focus to self as opposed to task, as well as seek avoidance of certain social situations. This may lead to performance deterioration and increased risk of withdrawal from sport. Athletes with PTSD may have physiological symptoms resulting in increased muscle tension, hypervigilance, increased startle or fear response, and negative mood symptoms that may result in performance decline and potentially increased risk for injury. In addition, athletes with PTSD symptoms may avoid triggers related to the trauma, which may include actions that are necessary for their sport performance (eg, a specific route in football, set piece in soccer, or arm motion in baseball). Athletes with OCD may have impaired performance if intrusive thoughts interfere with present-moment attention or if they have extreme difficulty in stopping or completing the obsessive-compulsive routine to engage in the actual performance.

Prevention

Many athletes have normal “state anxiety,” meaning they become appropriately anxious before competition, but it does not permeate their entire life.²²¹ Consequently, primary prevention of anxiety may not be the most useful focus for elite athletes. Interventions exist to assist athletes in regulating anxiety before athletic performance, and preliminary studies have been performed on secondary management of anxiety and rehabilitation after injury using mindfulness and CBT.^{222,223}

Overtraining

Detection in Athletes

Overtraining syndrome in athletes occurs when excessive training loads with inadequate rest or recovery periods lead to persistent sport-specific performance deficits combined with mood disturbances.^{224,225} Overtraining syndrome has been reported in 10% to 64% of athletes, seems to have increased risk at higher levels of sport, and has an increased risk of recurrence in an athlete who has been previously diagnosed.²²⁴ Several confounding factors of the athlete lifestyle can contribute to increased risk for OTS including poor nutrition, illness, psychosocial stressors (work, school, coach, and team), and sleep disorders.^{224,225}

Unfortunately, there are no current standardized diagnostic criteria for OTS. Performance testing using time-to-fatigue measures may be more sensitive for evaluating overtraining than graded exercise stress testing or anaerobic testing.²²⁶ Baseline hormonal biomarkers have not been proven to help diagnose OTS. However, a systematic review showed that athletes with OTS may have a blunted hormonal response to maximal exertion stress in prolactin, growth hormone, and adrenocorticotropic hormone measurements.²²⁷ Although some athletes also had blunted catecholamine response to acute stress and others had exacerbated catecholamine response, this may be explained by the difference in endurance versus strength athletes.²²⁷ A study in endurance athletes showed that changes in heart rate and lactate levels were useful to determine over-reached athletes.²²⁸

The psychological response to overtraining has been successfully evaluated using the Profile of Mood States (POMS) rating scale,²²⁴ a seven-question derivative called the Training Distress Scale, which may be more accurate,²²⁹ and a POMS energy index focused on the fatigue and vigor subscores.²³⁰ Another test is the Recovery-Stress Questionnaire for Athletes (RESTQ-Sport) (see below).²³¹ Perfectionistic concerns have been shown to predict training distress, which is a component of OTS.²³² Reaction time, attention measures, heart rate variability, and immune system function are all tests that have been evaluated for detection of overtraining without conclusive evidence at this time.

Management

There is also no single universal ideal biological marker to guide the management of OTS. Blood lactate level may be useful because it has an inverse relationship with muscle and liver glycogen levels. Although lactate levels should not be used alone in monitoring OTS, one finding is that athletes with OTS have diminished maximal lactate concentration levels, while submaximal levels are within normal limits.²²⁴ Along with blood lactate, monitoring heart rate is important, and can identify almost 90% of athletes who may be developing OTS.²²⁴ Measuring creatine kinase and heart rate variability are 2 other biologic markers that have been studied to manage OTS with mixed results.²³¹

Mood disturbances may be a part of the clinical diagnosis of OTS and should be monitored in the athlete with suspected OTS. The Recovery-Stress Questionnaire for Athletes (RESTQ-Sport) is one such tool. It is a 77-point form that inquires about stress (general, emotional, and social), conflicts and pressure, fatigue, lack of energy, and somatic symptoms as well as recovery scales and sport-specific stress scales.²³¹

A management approach to the athlete with OTS should be individually developed. It should include relative or absolute rest depending on the clinical situation at the time. The use of medications (eg, antidepressants) to assist in treatment is still controversial at this time, and they should be used with caution and a full awareness of the potential side effects. Good sleep hygiene and proper nutrition may be useful for the athlete in recovery. Of note, a high carbohydrate diet has been studied in a small cohort (n = 56), with some promising results for the management of OTS.²²⁴

The physical demands and the psychological factors that contribute to the development of OTS should be addressed as part of the individualized treatment plan of the athlete with OTS. Recovery from OTS, once properly recognized and managed, can be measured in weeks to months to years, adding to the challenge of this condition in athletes.

Effect on Performance

By definition, performance is negatively affected in the athlete with OTS. “Underperformance syndrome” has been used in the literature to more accurately describe the overtrained athlete. If the goal of training is to provide loads that result in performance improvement, then OTS results in performance decrement. In OTS, there is an accumulation of stress that causes longer-term decrease in performance and that may include physiological and psychological changes as well as the consequences of dysfunction in these areas.²³³ The difference between “overreaching” and “overtraining” described in the literature involves a 2-week cutoff—that is, a 2-week period of adequate recovery that results in the resumption of previous performance levels—is typically called overreaching in an athlete, and recovery requiring more than 2 weeks may raise the possibility of OTS.²³³

A core component of OTS is the lack of ability to sustain intense exercise resulting in a decrement in performance capacity. Thus, an athlete with OTS may be able to start a training run at their normal pace but be unable to complete the normal training load. Performance testing can be used to measure the extent of this negative impact on performance in an athlete, and this may include time-to-fatigue tests, time trials, or sports-specific performance tests.

Prevention

Monitoring training loads, getting adequate rest periods, and maintaining optimal nutrition and hydration status are all important in preventing the development of OTS.²²⁴ Unfortunately, specific protocols are not known at this time, and this will likely require individual athlete program tailoring. Documenting training loads using rate of perceived exertion and duration of training is important when athletes have shown different perceived training loads than their coaches may intend.²³⁴ The use of psychological mood monitoring to guide training loads has been shown to decrease risk of staleness, a key risk component in OTS.²³⁵

Sleep Disorders

Detection in Athletes

There is a paucity of data associating sleep disorders in athletes with physical and MH disorders. In college students,

there is a high prevalence of common mental disorders comorbid with sleep disorders; students who experienced poor sleep quality had a 2.4 times higher odds of depression, anxiety, and somatoform disorder than those students with good sleep quality.²³⁶ The prevalence of poor sleep quality with high levels of daytime sleepiness in athletes is as high as 50% to 83%,^{237,238} and prevalent poor sleep quality has also been shown in elite athletes with disabilities.²³⁹

Symptoms associated with insomnia can include lack of concentration, irritability, and depression. During yearly psychological evaluations of elite athletes assessed for insomnia symptoms, ongoing sleep problems were reported by just over 20%, with difficulty falling asleep and nocturnal waking reported more often in women.²⁰⁶

Obstructive sleep apnea (OSA) has been linked to cognitive impairment and mood disorders,^{240,241} but few studies look at the prevalence in athletes. Physical characteristics of athletes in certain sports such as American football and rugby may predispose them to OSA.²³⁸ American football players exhibit several risk factors for OSA, including large neck circumference and high BMI. The prevalence of sleep-disordered breathing among collegiate football players is estimated to be 8%,²⁴² and the rate of OSA in professional football players exceeds the population with a 5 to 11 times greater risk.²⁴³

The Athlete Sleep Screening Questionnaire is a specific screening tool for athletes with sleep disorders that have been clinically validated.²⁴⁴ The Insomnia Severity Index²⁴⁵ and the STOP-Bang Questionnaire²⁴⁶ are 2 reliable and valid questionnaires designed for the general public that may also be used for athletes.

Management

High-quality research, while not specific for athletes, has established that insomnia-specific CBT (CBTi) is first-line treatment for sustained improvements in sleep in those with insomnia alone or insomnia comorbid with other conditions.^{247,248} One large study found that 60% of team-sport athletes had no strategy to overcome poor sleep compared with 33% of individual athletes, who used simple relaxation and reading techniques.²⁴⁹

Extending sleep to a minimum of 10 hours a night has been shown to improve mood in elite athletes.^{250,251} Sleep education and optimization programs have led to significant improvements in self-reported total sleep time, sleep efficiency, fatigue, and vigor in athletes.²⁵²

Athletes may prefer a “natural” solution to assist with their sleep problems. Research studying the influence of nutritional interventions is minimal and somewhat inconclusive. A review of multiple studies, none specific to athletes, concluded that diets high in carbohydrate may result in shorter sleep latencies, those high in protein may result in improved sleep quality, and those high in fat may negatively influence total sleep time.²⁵³

High-quality RCTs exploring sleep medications’ influence on reliable measures of athletes’ mental and physical performance is lacking. With benzodiazepine-sedative hypnotics, individuals can have a marked “hangover” effect, and reaction time can be negatively impacted.^{254,255} There is mixed evidence as to whether nonbenzodiazepine benzodiazepine receptor agonist hypnotics such as zolpidem show impairment on measures of psychomotor and physical performance.^{256–258}

Although melatonin is the first choice of psychiatrists for the treatment of insomnia in athletes,²⁰¹ research investigating its use for primary insomnia is inconclusive. There is no evidence that melatonin is effective in the management of most primary sleep disorders with short-term use, although it may reduce sleep-onset latency (the length of time it takes a person to go from complete wakefulness to the first stage of sleep) for persons with delayed sleep-phase syndrome, where the sleep/wake cycle is delayed with respect to the external day/night cycle. Evidence does suggest that short-term use of melatonin is safe.²⁵⁹ Although no decrements in performance were observed the next morning after athletes ingested melatonin, it was not shown to improve their sleep quality.²⁶⁰ Exogenous melatonin is not regulated by the FDA, and therefore, elite athletes enrolled in a drug testing program must use this over-the-counter (OTC) supplement with caution as purity cannot be guaranteed. The prescription medication ramelteon, a melatonin receptor agonist, carries a longer half-life than melatonin and therefore has a greater likelihood of causing hangover effect; it has also not been studied in athletes.²⁶¹

Although trazodone is approved in the United States for the treatment of depression, it is more commonly prescribed off-label for insomnia. No studies have been performed in athletes, and only one RCT comparing treatment with trazodone, zolpidem, and placebo found an absence of significant efficacy for trazodone, but significantly more side effects including headache and somnolence.²⁶²

Antihistamines and other first-generation antihistamines are often included in OTC sleep medications; however, they can have adverse anticholinergic properties (such as dry mouth and urinary retention) and a quicker onset of tolerance and longer duration of action than many prescription sleep aids. There is weak evidence demonstrating an absence of efficacy in the treatment of sleep onset insomnia, with minimal evidence of adverse events in excess of placebo.²⁶³

The primary treatment for OSA is continuous positive airway pressure and weight loss as indicated, although the latter is impractical for athletes in certain sports. No athlete-specific studies have been performed.

Effect on Performance

Sleep plays a major role in recovery and performance capacity in elite athletes. Studies in athletes and nonathletes have linked decreases in both the quality and duration of sleep with detriments in overall health including impaired cognitive functioning and judgment, mood problems, and somatic symptoms,²⁶⁴ as well as an increase in perceived physical exertion and decrease in pain tolerance.²⁶⁵ As sleep deprivation may cause or modulate acute and chronic pain, and pain may disturb sleep by inducing arousals during sleep, these 2 issues can augment each other, and a continuous pain cycle can develop.²⁶⁶

Athletes diagnosed with OTS showed decreased sleep quality, leading to the conclusion that worsened sleep was likely a trigger.²⁶⁷ Seventy percent of athletes who had disrupted sleep before a competition had negative moods of fatigue and tension.²⁶⁸

Sleep loss is associated with an increase in both sympathetic activity and catecholamine levels, which, over time, may lead to altered stress system responsiveness, similar to that seen in mood disorders.²⁶⁹ Sleep deprivation is also a risk factor for illicit substance and alcohol use, violence-related behaviors,

and motor vehicle accidents.^{270–272} Although this can certainly affect performance, whether or not this translates to athletes is unknown.

Prevention

The prevention of sleep disorders and poor sleep habits starts with the identification and elimination of risk factors and behaviors that would reduce the quantity and quality of sleep, establishing and maintaining a regular sleep routine, and ensuring adequate sleep duration depending on age and hours of training each day. Health care providers should discourage the use of computers and other tech devices, and the watching of screens before bedtime because these may perpetuate sleep deficiency and disrupt circadian rhythms, which can have adverse impacts on athletic performance.²⁷³ Recognize that excessive worry and anxiety related to training, competition, academics, or personal relationships may cause significant emotional reactions that decrease sleep quality.

Although athletes are often advised to not exercise before bedtime due to potential overstimulation, evening exercise (2–4 hours before bed) has not been associated with worse sleep.^{274,275} High exercise levels are also related to improved sleep and psychological functioning. Adolescent athletes who trained almost 18 hours a week (vs 4.5 hours a week for the control group) reported better sleep patterns including higher sleep quality, shortened sleep-onset latency, and fewer awakenings after sleep onset. They also reported less tiredness and increased concentration during the day, and significantly fewer anxiety and depressive symptoms.²⁷⁶

Attention-Deficit/Hyperactivity-Deficit

Detection in Athletes

Attention-deficit/hyperactivity deficit (ADHD) is an important issue for the team physician taking care of athletes. The health care provider should be familiar with making the diagnosis of ADHD, the basic management of ADHD, and how medications used to treat ADHD impact exercise and performance.^{277–280} The diagnosis of ADHD should be made when the individual meets the DSM-5 criteria and after the careful evaluation and consideration of competing comorbid diagnoses.

Management

The optimal management and treatment approach for ADHD is an individualized one that includes behavioral therapies and consideration of medications.^{280–286} Assessment of the athlete's history and background, support structures, psychiatric and medical comorbid diagnoses, and previous responses to interventions including compliance and medication side effects is recommended.

The decision to use medications during sport is one that should be made on an individual basis, and team physicians should be aware of the effects of medication treatment options on certain sport and athlete-specific situations. The fear of potential misuse of stimulants or other recreational drugs is not justification for withholding pharmacologic treatment of ADHD,^{287–289} unless there are other risk factors for substance misuse in an individual athlete. There are also some data to suggest that exercise in and of itself can improve cognitive performance in children with ADHD.²⁹⁰

Although groups offer different guidelines, there is insufficient evidence to recommend routine electrocardiogram or additional cardiac testing in athletes without cardiac disease who are on medications for ADHD, unless they are simultaneously on other medications that may impact cardiac health, such as tricyclic antidepressants. The American Heart Association recommends ECG with nearly all treatments for ADHD.²⁹¹ The American Academy of Pediatrics states that there is no need for ECG with stimulant treatment unless there is concern for cardiac disease.^{292–294} If an athlete is at risk by history, family history, and/or physical examination, then ECG and additional workup is indicated.²⁷⁷

There is also no evidence to support withholding stimulant treatment for ADHD due to a fear of sudden cardiac death; however, athletes should be monitored for signs and symptoms suggestive of cardiac disease.^{295–297} A very large retrospective cohort study of over 1.2 million children and young adults and 2.5 million person years follow-up, including almost 375 000 person years of current use of ADHD drugs, did not find any increased risk of cardiovascular events for users of ADHD drugs.²⁹⁸

Team physicians should be aware of and educate athletes on regulations and requirements with pharmacologic treatments of ADHD. Team physicians and athletes should use resources provided by the IOC,²⁹⁹ NCAA,³⁰⁰ and World Anti-doping Association (WADA),³⁰¹ depending on their level of competition, to determine what medications and/or methods are permissible and/or require a TUE to participate. The IOC and WADA allow for the use of stimulant medication, with a TUE form completed. At the NCAA level, stimulants are only allowable for ADHD if institutions are able to submit the “NCAA Medical Exception Documentation Reporting Form to Support the Diagnosis of Attention Deficit Hyperactivity Disorder (ADHD) and Treatment with Banned Stimulant Medication” and supporting documentation to the NCAA in the event the athlete tests positive for stimulants. This documentation must include comprehensive clinical evaluation using DSM-5 criteria; blood pressure and heart rate readings and comments; diagnosis, medication(s), and dosage; and confirmation that nonstimulant medications have been considered.

Effect on Performance

The athletic care network must be aware of the athlete's medications and how the athlete is taking them, as use may vary with individual athlete preference. Some athletes will only take medications episodically for school testing or for studying purposes, whereas others feel that their sport performance is improved on stimulants. Some athletes temporarily stop taking them, so that their sports play appears more random or unpredictable to opponents, which they feel improves their performance.³⁰² When an athlete first starts taking medication, it is important to consider that the medication may affect exercise and performance; thus, medication is best initiated in a low stress (eg, captain's practice) versus high stress (eg, game) activity.²⁷⁷

There may be an increased risk of heat injury in athletes taking medications for ADHD, specifically stimulants and bupropion (the latter has been used off-label to treat ADHD).²⁰¹ Some evidence suggests that athletes taking stimulant medications have elevated core temperatures while exercising, although an increased incidence of exertional heat

injury or heatstroke in these groups has not been reported.^{303–306} Although evidence for the ergogenic effect of medications to treat ADHD is unclear, these medications are often used and misused because of this perception.^{201,307–309}

SUMMARY

The primary goal of this AMSSM position statement is to assist the team physician and other members of the athletic care network with the detection, treatment, and prevention of a select range of psychological issues and MH disorders in athletes. An important component of management is an understanding of pharmacological treatment options including those that may be the most effective with the fewest side effects. This document also addresses topics not fully explored in previous publications about MH in athletes. Critical insight is needed into key personality issues (eg, “athlete identity”), demographic and cultural variables (eg, sexual orientation and gender identification), and environmental conditions (eg, hazing, bullying, and sexual abuse) that can impact athletes, and how interactions among these variables may contribute to MH issues. It is important for the athletic care network to be attuned to risk factors for MH disorders, and to monitor athletic environments that may trigger or exacerbate psychological issues in athletes under their care.

We acknowledge the limited evidence-based data specifically addressing the athlete, and AMSSM supports continued research in this area to validate the optimal strategies for the detection, management, and prevention of MH disorders in athletes. Although not specifically addressed due to the already large scope of this position statement, AMSSM also supports research looking at the potentially increased risk of depression and other MH disorders after sports-related concussions in athletes.

The following summary conclusions and recommendations include the following:

Personality Issues

1. High athletic identity is associated with both positive and negative health and performance outcomes (SORT B).⁸
2. Personality traits and disorders deemed problematic for athletes may be best addressed through psychotherapy (SORT C).⁹

Sexuality and Gender Issues

1. The creation of a strong supportive environment that is welcoming to sexual minorities is key to the MH of the athlete and the sports team (SORT A).^{10,11}
2. Reducing the risk of negative health consequences for the sexual minority athlete starts with education of all stakeholders associated with athletic participation (SORT C).^{12–14}

Hazing

1. Hazing leads to both short- and long-term health ramifications that can affect an individual’s athletic success and ability to participate in sport (SORT C).¹⁵
2. The prevention and management of hazing requires a global investment from athletes, coaches,

administrators, and health care providers centered on a zero-tolerance policy for any form of maltreatment and a focus on positive team building activities that promote dignity and teamwork as opposed to victimization (SORT C).¹⁶

Bullying

1. Bullying in athletics can take on many different forms and be the actions of teammates or coaches. Signs and symptoms of being bullied may vary greatly (SORT C).¹⁷
2. Preventing bullying is the responsibility of all the stakeholders in athletics. Educational programs can be found on the NCAA website (SORT C).¹⁷

Sexual Misconduct

1. Authority figures are more often perpetrators of sexual abuse, but peer athletes are far more likely than coaches to be perpetrators of sexual harassment (SORT C).^{18,19}
2. Populations at higher risk for sexual abuse in youth sports reflect trends in the general population. Those participating at higher levels of competition are also at an increased risk. Sport type, amount of touching, or degree of clothing cover during participation do not seem to correlate with higher rates of abuse (SORT C).^{20–23}

Transitioning from Sport

1. Athletic departments, national governing bodies, and professional leagues should assist athletes who are retiring from their sport with development of a comprehensive preretirement plan addressing issues surrounding their transition out of athletic participation (SORT A).²⁴
2. Long-term psychological effects of career-ending injuries are common for many athletes (SORT C).²⁵

Psychological Response to Injury and Illness

1. Psychological and sociocultural factors have been raised as potential risk factors for injury. Stress consistently demonstrates a relationship with injury risk as well as the ability to rehabilitate from injury and return to sport (SORT B).^{25,26}
2. Cognitive, emotional, and behavioral responses to injury are important in determining outcome (SORT C).²⁷

Self-Medication in Response to Injury/Illness

1. Limited data exist on the use of self-medication by athletes as a coping mechanism. However, certain demographics of athletes are emerging as higher risk groups for medication misuse and for negative MH and other consequences of their use (SORT C).^{28,29}
2. Targeted interventions that incorporate health and athletic performance considerations tend to be more successful for the athletic population, and this includes addressing the underlying issues leading to substance use/self-medication (SORT C).^{30–32}

Eating Disorder/Disordered Eating

1. Annual preparticipation screening for EDs in athletes should be routine (SORT C).^{33–39}
2. Eating disorder prevention programs have benefit in reducing risk for EDs (SORT B).^{33–36,39–42}
3. Cognitive behavioral therapy and family therapy are recommended as treatments for EDs in athletes (SORT B).^{33–36,40,43}

Depression and Suicide

1. Athletes have unique risk factors for depression compared with nonathletes. Early recognition and appropriate management of depression in athletes lead to improved clinical and performance outcomes (SORT C).⁴⁴
2. College student-athletes report depression symptoms at a higher prevalence than previously reported; these rates are comparable with nonathlete college students (SORT B).⁴⁵
3. Suicide incidence in college student-athletes is lower than in college student nonathletes. Football has the highest suicide rate by sport in college athletes (SORT B).⁴⁶

Anxiety/Stress

1. Cognitive behavioral therapy for the treatment of anxiety is the optimal nonpharmacological intervention. Cognitive behavioral therapy is an established and effective treatment method for many clinical populations with different types of anxiety disorders, but there are no RCTs of CBT interventions specifically within athletes (SORT B).⁴⁷
2. Although SSRIs may be considered, as-needed anxiolytics are not recommended for athletic performance anxiety (SORT B).^{48,49}

Overtraining

1. A management approach to the athlete with OTS should be individually developed and should include evaluation for MH stressors and relative or absolute rest depending on the clinical situation at the time (SORT C).⁵⁰
2. Monitoring training loads, getting adequate rest periods, and maintaining optimal nutrition and hydration status are all important in preventing the development of OTS (SORT C).⁵¹

Sleep

1. Although not specific for athletes, insomnia-specific CBT is first-line treatment for sustained improvements in sleep in those with insomnia alone or insomnia comorbid with other MH disorders (SORT A).^{52,53}
2. Benzodiazepine-sedative hypnotics are not recommended for athletes because of their marked “hangover” effect, which includes a negative impact on reaction time (SORT A).^{54,55}
3. Although melatonin has not been shown to improve sleep quality in athletes, short-term use is safe with no decrements in performance (SORT A).^{56,57} Because melatonin is not regulated by the FDA, caution for the presence of impurity is necessary, and it should be purchased as a single-ingredient product from a reputable company.

Attention-Deficit Hyperactivity Disorder

1. The optimal management approach for ADHD is individualized and may include behavior therapies, academic accommodations, pharmacotherapy (eg atomoxetine, amphetamine salts, or methylphenidate formulations), and psychological interventions to manage associated features and comorbid diagnoses (SORT C).^{58–61}
2. The risk of heat illness may be increased in athletes taking ADHD medications. Those taking stimulant medications have elevated core temperatures while exercising, although an increased incidence of exertional heat injury or heatstroke in these groups has not been reported (SORT C).^{62–65}
3. Team physicians should be aware of and educate the athlete on regulations and requirements regarding medication treatment of ADHD (SORT A).^{66–68}

ACKNOWLEDGMENTS

The authors sincerely thank Libby Barton, MD, for her meticulous review of the references and Stephen Paul, MD, for his encouragement and review of the manuscript.

References

1. Herring SA, Kibler W, Putukian M, et al. Sideline preparedness for the team physician: a consensus statement-2012 update. *Med Sci Sports Exerc.* 2012;44:2442–2445.
2. Herring SA, Kibler WB, Putukian M, et al. Team physician consensus statement: 2013 update. *Med Sci Sports Exerc.* 2013;45:1618–1622.
3. National Collegiate Athletic Association Sports Science Institute. *Inter-association Consensus Statement. Best Practices for Understanding and Supporting Student-Athlete Mental Wellness.* Indianapolis, IN: National Collegiate Athletic Association Sport Science Institute. Available at: http://www.ncaa.org/sites/default/files/HS_Mental-Health-Best-Practices_20160317.pdf. Accessed February 2, 2018.
4. Herring SA, Kibler WB, Putukian M, et al. Psychological issues related to illness and injury in athletes and the team physician: a consensus statement—2016 update. *Med Sci Sports Exerc.* 2017;49:1043–1054.
5. Brown GT, Hainline B, Kroshus E, et al. *Mind, Body and Sport: Understanding and Supporting Student-Athlete Mental Wellness.* NCAA Publications; 2014. Available at: <http://www.ncaapublications.com/productdownloads/MindBodySport.pdf>. Accessed February 18, 2018.
6. Haase AM, Prapavessis H, Owens RG. Perfectionism, social physique anxiety and disordered eating: a comparison of male and female elite athletes. *Psych Sport Exerc.* 2002;3:209–222.
7. Smith RE, Smoll FL, Cumming SP. Effects of a motivational climate intervention for coaches on young athletes' sport performance anxiety. *J Sport Exerc Psychol.* 2007;29:39–59.
8. Hendawy HMF, Awad EAA. Personality and personality disorders in athletes. In: Baron DA, Reardon CL, Baron SH, eds. *Clinical Sports Psychiatry: An International Perspective.* Oxford, United Kingdom: Wiley-Blackwell; 2013:53–64.
9. Samadzadeh M, Abbasi M, Shahbazzadegan B. Comparison of sensation seeking and self-esteem with mental health in professional and amateur athletes, and non-athletes. *Proced Soc Behav Sci.* 2011;15:1942–1950.
10. Shariati M, Bakhtiari S. Comparison of personality characteristics athlete and non-athlete student, Islamic Azad University of Ahvaz. *Proced Soc Behav Sci.* 2011;30:2312–2315.
11. McKelvie SJ, Lemieux P, Stout D. Extraversion and neuroticism in contact athletes, no contact athletes and non-athletes: a research note. *Athletic Insight J Sport Psychol.* 2003;5:19–27.
12. Schroth ML. A comparison of sensation seeking among different groups of athletes and nonathletes. *Pers Individ Dif.* 1995;18:219–222.
13. Nia ME, Besharat MA. Comparison of athletes' personality characteristics in individual and team sports. *Proced Soc Behav Sci.* 2010;5:808–812.
14. Hill AP, Hall HK, Appleton PR, et al. Perfectionism and burnout in junior elite soccer players: the mediating influence of unconditional self-acceptance. *Psychol Sport Exerc.* 2009;9:630–644.
15. Frost RO, Marten P, Laharat C, et al. The dimensions of perfectionism. *Cognit Ther Res.* 1990;14:449–468.

16. Soenens B, Elliot J, Goossens L, et al. The intergenerational transmission of perfectionism: parents' psychological control as an intervening variable. *J Fam Psychol.* 2005;19:358–366.
17. Brewer BW, Van Raalte JL, Linder DE. Athletic identity—Hercules muscles or Achilles heel? *Int J Sport Psychol.* 1993;24:237–254.
18. Linville PW. Self-complexity as a cognitive buffer against stress-related illness and depression. *J Pers Soc Psychol.* 1987;52:663–676.
19. Watkins CE, Campbell VL, Nieberding R, et al. Contemporary practice of psychological assessment by clinical psychologists. *Prof Psychol Res Pr.* 1995;26:54–60.
20. Westen D. Divergences between clinical and research methods for assessing personality disorders: implications for research and the evolution of Axis II. *Am J Psychiatry.* 1997;154:895–903.
21. Widiger TA, Boyd SE. Personality disorders assessment instruments. In: Butcher JN, ed. *Oxford Handbook of Personality Assessment.* Oxford, United Kingdom: Oxford University Press; 2012:336–363.
22. Butcher JN. Minnesota multiphasic personality inventory. In: Weiner IB, Craighaid WE, eds. *The Corsini Encyclopedia of Psychology.* Vol 3. Hoboken, NJ: John Wiley & Sons, Inc.; 2010.
23. Stillman MA, Ritvo EC, Glick ID. Psychotherapeutic treatment of athletes and their significant others. In: Baron DA, Reardon CL, Baron SH, eds. *Clinical Sports Psychiatry: An International Perspective.* Oxford, United Kingdom: Wiley-Blackwell; 2013:117–123.
24. Stirling AE, Kerr GA. Perfectionism and mood states among recreational and elite athletes. *Athletic Insight J Sport Psychol.* 2006;8:13–27.
25. Oswald SB, Vargas TM. How safe is the playing field? collegiate coaches' attitudes toward gay, lesbian, and bisexual individuals. *Sport Soc.* 2012; 16:120–132.
26. Greenspan SB, Griffith C, Murtagh EF. LGBTQ youths' school athletic experience: a 40 year content analysis in nine flagship journals. *J LGBT Issues Couns.* 2017;11:190–200.
27. Mountjoy M, Brackenridge C, Arrington M, et al. International Olympic Committee consensus statement: harassment and abuse (non-accidental violence) in sport. *Br J Sports Med.* 2016;50:1019–1029.
28. Rotella RJ, Murray MM. Homophobia, the world of sport, and sport psychology consulting. *Sport Psychol.* 1991;5:355–364.
29. Reardon CL, Baron DA, Baron SH, et al. The role of culture in sports. In: Baron DA, Reardon CL, Baron SH, eds. *Clinical Sports Psychiatry: An International Perspective.* Oxford, United Kingdom: Wiley-Blackwell; 2013:179–187.
30. Kroshus E, Davoren AK. Mental health and substance use of sexual minority college athletes. *J Am Coll Health.* 2016;64:371–379.
31. Barber H, Krane V. Creating a positive climate for lesbian, gay, bisexual, and transgender youths. *J Phys Educ Recreat Dance.* 2007;78:6–8.
32. Mereish EH, Poteat VP. A relational model of sexual minority mental and physical health: the negative effects of shame on relationships, loneliness, and health. *J Couns Psychol.* 2015;62:425–437.
33. Rosario M, Reisner SL, Corliss HL, et al. Sexual-orientation disparities in substance use in emerging adults: a function of stress and attachment paradigms. *Psychol Addict Behav.* 2014;28:790–804.
34. Bauman NJ. The stigma of mental health in athletes: are mental toughness and mental health seen as contradictory in elite sport? *Br J Sports Med.* 2016;50:135–136.
35. Morris JF, Van Raalte JL. Transgender and gender nonconforming athletes: creating safe spaces for all. *J Sport Psychol Action.* 2016;7: 121–132.
36. Lucas-Carr CB, Krance V. What is the T in the LGBT? supporting transgender athletes through sport psychology. *Sport Psychol.* 2011;25: 532–548.
37. Women's Sports Foundation. *Playing in the Closet: Homophobia in Sports.* New York, NY: Women's Sports Foundation; 2011. Available at: <https://www.womenssportsfoundation.org/research/article-and-report/lgbt-issues/playing-in-the-closet/>. Accessed June 22, 2018.
38. National Collegiate Athletic Association. *LGBTQ Resources: NCAA Inclusion Initiative Framework [Internet].* Available at: www.ncaa.org/about/resources/inclusion/lgbtq-resources. Accessed February 3, 2018.
39. National Collegiate Athletic Association. *Champions of Respect: Inclusion of LGBTQ Student-Athletes and Staff in NCAA Programs [Internet].* Available at: www.ncaapublications.com/p-4305-champions-of-respect-inclusion-of-lgbtq-student-athletes-and-staff-in-ncaa-programs.aspx. Accessed February 2, 2018.
40. *Championing LGBTQ Issues in K-12 Education since 1990 [Internet].* 1. Available at: www.glsen.org/?gclid=Cj0KCQiAnuDTBRDUARsAL41eDoyihJzbflrBbKOTm036Omp2BP4KcQjOwCu6HkvNpcjwV3MT27AwaAviMEALw_wcB. Accessed February 5, 2018.
41. *Changing the Game: The GLSEN Sports Project [Internet].* 2. Available at: www.glsen.org/sports. Accessed February 1, 2018.
42. *Go! Athletes 2018 [Internet].* Available at: goathletes.org/. Accessed February 1, 2018.
43. *Athlete Ally: Victory through Unity [Internet].* Available at: <https://www.athletheally.org/about/>. Accessed February 1, 2018.
44. Allan EJ, Orono ME. *Hazing vs. Bullying.* StopHazing.Org. Available at: <https://web.archive.org/web/20150906002200/http://www.stophazing.org/hazing-vs-bullying/>. Accessed March 27, 2018.
45. Gladden R, Vovolo-Kantor A, Hamburger M, et al. *Bullying Surveillance Among Youths: Uniform Definitions for Public Health and Recommended Data Elements, Version 1.0.* Atlanta, GA: National Center for Injury Prevention and Control, Centers for Disease Control and Prevention and U.S. Department of Education; 2014.
46. Allan EJ, Madden M. *Hazing in View: College Students at Risk.* 2008. Available at: http://www.stophazing.org/wp-content/uploads/2014/06/hazing_in_view_web1.pdf. Accessed January 21, 2019.
47. Waldron JJ, Kowalski CL. Crossing the line: rites of passage, team aspects, and ambiguity of hazing. *Res Q Exerc Sport.* 2009;80: 291–302.
48. Gershel JC, Katz-Sidlow RJ, Small E, et al. Hazing of suburban middle school and high school athletes. *J Adolesc Health.* 2003;32:333–335.
49. Leahy T, Pretty G, Tenenbaum G. Perpetrator methodology as a predictor of traumatic symptomatology in adult survivors of childhood sexual abuse. *J Interpers Violence.* 2004;19:521–540.
50. Anderson E, McCormack M, Lee H. Male team sport hazing initiations in a culture of decreasing homophobia. *J Adolesc Res.* 2012;27: 427–448.
51. Hoover NC. *National Survey: Initiation Rites and Athletics for NCAA Sports Teams.* Alfred University; 1999. Available at: http://www.alfred.edu/sports_hazing/docs/hazing.pdf. Accessed January 21, 2019.
52. Irick E. *Student-Athlete Participation 1981-82—2015-16. NCAA Sports Sponsorship and Participation Rates Report.* Indianapolis, IN: The National Collegiate Athletic Association; 2016.
53. Hoover NC, Pollard N. *High School Hazing Initiation Rites in American High Schools: A National Survey.* Alfred, NY: Alfred University; 2000.
54. Fields S, Collins C, Comstock R. Violence in youth sports: hazing, brawling and foul play. *Br J Sports Med.* 2010;44:32–37.
55. Hoover J, Milner C. Are hazing and bullying related to love and belongingness? *Reclaim Child Youth.* 1998;7:138–141.
56. Johnson J. Through the liminal: a comparative analysis of communities and rites of passage in sport hazing and initiations. *Can J Sociol.* 2011; 36:199–227.
57. Diamond A, Callahan S, Chain K, et al. Qualitative review of hazing in collegiate and school sports: consequences from a lack of culture, knowledge and responsiveness. *Br J Sports Med.* 2016;50:149–153.
58. Wilfert M. *NCAA Study of Student-Athlete Social Environments.* Research presented at the 2014 NCAA Convention, San Diego, California, January 2014.
59. Neal TL, Diamond AB, Goldman S, et al. Interassociation recommendations for developing a plan to recognize and refer student-athletes with psychological concerns at the secondary school level: a consensus statement. *J Athl Train.* 2015;50:231–249.
60. Hoover NC, Pollard NJ. *Initiation Rites in American High Schools: A National Survey.* Alfred Univ Web site; 2014. Available at: http://www.alfred.edu/hs_hazing/. Accessed November 17, 2018.
61. Van Raalte J, Cornelius AE, Linder DE, et al. The relationship between hazing and team cohesion. *J Sport Behav.* 2007;30:491–507.
62. *Bullying [Internet].* Merriam-Webster.Com: Merriam-Webster; 2018. Available at: www.merriam-webster.com/dictionary/bullying. Accessed February 17, 2018.
63. Schinnerer JL. *The Consequences of a Verbally Abusive Athletic Coach [Internet].* PsychCentral. Available at: psychcentral.com/lib/the-consequences-of-verbally-abusive-athletic-coaches/. Accessed February 17, 2018.
64. Evans B, Adler A, MacDonald D, et al. Bullying victimization and perpetration among adolescent sport teammates. *Pediatr Exerc Sci* 2016; 28:296–303.
65. Swigonski NL, Enneking BA, Hendrix KS. Bullying behavior by athletic coaches. *Pediatrics.* 2014;133:273–275.
66. Sicking J. *Bullying Still Occurs in College, Professors Find [Internet].* Indiana State University Newsroom; 2011. Available at: www2.indstate.edu/news/news.php?newsid=2904. Accessed May 5, 2018.
67. National Collegiate Athletic Association. *Cyberbullying. Mind, Body, and Sport: Interpersonal Violence and Student-Athlete Population: An Excerpt from the Sport Science Institute's Guide to Understanding and Supporting Student-Athlete Mental Wellness [Internet].*

- Available at: www.ncaa.org/sport-science-institute/mind-body-and-sport-interpersonal-violence-and-student-athlete-population. Accessed May 5, 2018.
68. National Collegiate Athletics Association. *NCAA Study of Student Athlete Social Environments (2012–2016) January 2017 Preliminary Report [Internet]*. 2017. Available at: www.ncaa.org/sites/default/files/2017RES_NCAA_Convention_Social_Environments_present_20170807.pdf. Accessed August 7, 2017.
 69. National Centre Against Bullying. *Signs of Bullying [Internet]*. Available at: www.ncab.org.au/bullying-advice/bullying-for-parents/signs-of-bullying/. Accessed May 5, 2018).
 70. Saylor CF, Nida SA, Williams KD, et al. Bullying and ostracism screening scales BOSS: development and application. *J Child Health Care*. 2012; 41:322–343.
 71. Felix ED, Sharkey JD, Greif Green JG, et al. Getting precise and pragmatic about the assessment of bullying: the development of the California Bullying Victimization Scale. *Aggress Behav*. 2011;37: 234–247.
 72. University of Arizona C.A.T.S. *Life Skills. Step up [Internet]*. Available at: stepupprogram.org/about/. Accessed February 4, 2018.
 73. NCAA Sport Science Institute. *Bystander Intervention [Internet]*. Available at: www.ncaa.org/sport-science-institute/topics/bystander-intervention. Accessed February 14, 2018.
 74. Marks S, Mountjoy M, Marcus M. Sexual harassment and abuse in sport: the role of the team doctor. *Br J Sports Med*. 2012;46:905–908.
 75. National Council of Youth Sports. *Reports on Trends and Participation in Organized Youth Sports*. 2008. Available at: <http://www.ncys.org/pdfs/2008/2008-ncys-market-research-report.pdf>. Accessed January 1, 2020.
 76. Stirling A, Bridges E, Cruz E, et al. Canadian Academy of Sport and Exercise Medicine position paper: abuse, harassment, and bullying in sport. *Clin J Sport Med*. 2011;21:385–391.
 77. Fasting K, Chroni S, Hervik SE, et al. Sexual harassment in sport toward females in three European countries. *Int Rev Sociol Sport*. 2011;46:76–89.
 78. Leahy T, Pretty G, Tenebaum G. Prevalence of sexual abuse in organised competitive sport in Australia. *J Sex Aggress*. 2002;8:16–36.
 79. Toftegaard NJ. The forbidden zone: intimacy, sexual relations and misconduct in the relationship between coaches and athletes. *Int Rev Sociol Sport*. 2001;36:165–182.
 80. Brackenridge CH, Bishopp D, Moussali S, et al. The characteristics of sexual abuse in sport: a multidimensional scaling analysis of events described in media reports. *Int J Sport Exerc Psychol*. 2008;16:385–406.
 81. Kirby S, Greaves L, Hankivsky O. *The Dome of Silence: Sexual Harassment and Abuse in Sport*. London, United Kingdom: Zed Books; 2000.
 82. Cense M, Brackenridge CH. Temporal and developmental risk factors for sexual harassment and abuse in sport. *Eur Phy Educ Rev*. 2001;7: 61–79.
 83. Allan E. *Hazing in View: College Students at Risk: Initial Findings from the National Study of Student Hazing*. Darby, PA: DIANE Publishing; 2009.
 84. Brackenridge C, Fasting K, Kirby S, et al. *Protecting Children from Violence in Sport: A Review with a Focus on Industrialized Countries*. Florence, Italy: United Nations Children's Fund (UNICEF); 2010.
 85. Vertommen T, Schipper-van Veldhoven N, Hartill MJ. Sexual harassment and abuse in sport: the NOC*NSF helpline. *Int Rev Sociol Sport*. 2015;50:822–839.
 86. Fasting K, Brackenridge CH, Sundgot-Borgen J. Prevalence of sexual harassment among Norwegian female elite athletes in relation to sport type. *Int Rev Sociol Sport*. 2004;39:373–386.
 87. Fasting K, Brackenridge CH, Sundgot-Borgen J. Experiences of sexual harassment and abuse amongst Norwegian elite female athletes and nonathletes. *Res Q Exerc Sport*. 2003;74:84–97.
 88. Waldron J, Lynn Q, Krane V. Duct tape, icy hot & paddles: narratives of initiation onto US male sport teams. *Sport Educ Soc*. 2011;16:111–125.
 89. Brackenridge CH, Kirby S. Playing safe: assessing the risk of sexual abuse to elite child athletes. *Int Rev Sociol Sport*. 1997;32:407–418.
 90. Alexander K, Stafford A, Lewis R. *The Experiences of Children Participating in Organized Sport in the UK*. Edinburgh, Scotland: The University of Edinburgh/NSPCC Child Protection Research Centre; 2011. Available at: <https://www.nspcc.org.uk/globalassets/documents/research-reports/experiences-children-participating-organised-sport-uk-main-report.pdf>. Accessed March 29, 2018.
 91. Malkin K, Johnston L, Brackenridge CH. A critical evaluation of training needs for child protection in UK sport. *Managing Sport Leis*. 2000;5: 151–160.
 92. Brackenridge CH, Fasting K. Sexual harassment and abuse in sport: the research context. *J Sex Aggress*. 2002;8:3–15.
 93. Brackenridge C. *Burden or Benefit? An Evaluation of SportsScotland's Child Protection Programme with Governing Bodies of Sport*. Edinburgh, Scotland: SportsScotland; 2004.
 94. Taylor AN. *Sometimes It's Necessary to Blame the Victim*. Washington, DC: The Chronicle of Higher Education; 2011.
 95. Zierler S, Feingold L, Laufer D, et al. Adult survivors of childhood sexual abuse and subsequent risk of HIV infection. *Am J Public Health*. 1991; 81:572–575.
 96. Lerner BH. Young doctors learn quickly in the hot seat. *The New York Times*. 2006. Available at: <https://www.nytimes.com/2006/03/14/health/young-doctors-learn-quickly-in-the-hot-seat.html>. Accessed January 21, 2019.
 97. Wilfert M; NCAA Education Services. *Building New Traditions: Hazing Prevention in College Athletics*. Indianapolis, IN: The National Collegiate Athletic Association; 2007.
 98. Gurnham D. Victim-blame as a symptom of rape myth acceptance? Another look at how young people in England understand sexual consent. *Leg Stud*. 2016;36:258–278.
 99. Freetly AJH, Kane EW. Men's and women's perceptions of non-consensual sexual intercourse. *Sex Roles*. 1995;33:785–802.
 100. Fasting K, Brackenridge CH, Walseth K. Consequences of sexual harassment in sport. *J Sex Aggress*. 2002;8:37–48.
 101. IOC Medical Commission Expert Panel. Consensus statement on sexual harassment and abuse in sport. 2007. Available at: https://stilledm.olympic.org/media/Document%20Library/OlympicOrg/News/20070802-IOC-adopts-Consensus-Statement-on-sexual-harassment-and-abuse-in-sport/EN-Sexual-Harassment-Abuse-In-Sport-report-1125.pdf#_ga=2.41144171.850791067.1522330555-100913010.1519307960. Accessed March 29, 2018.
 102. Fisher LA, Wrisberg CA. How to handle athlete's transition out of sport. *Athl Ther Today*. 2007;12:49–50.
 103. Miller L, Buttal FP. Are NCAA Division I athletes prepared for end of the athletic career transition? A literature review. *J Evid Inf Soc Work*. 2018; 15:52–70.
 104. Lavalley D. The effect of a life development intervention on sports career transition adjustment. *Sport Psychol*. 2005;19:193–202.
 105. National Collegiate Athletic Association. *Moving on!: A Physical Activity Transition Program for Student-Athletes [Internet]*. Available at: <http://www.ncaa.org/about/resources/research/movingphysicalactivitytransition>. Accessed January 23, 2018).
 106. National Collegiate Athletic Association. *An Introduction to Mind, Body and Sport: The NCAA's Chief Medical Officer Weighs in on the Sport Science Institute's New Guide to Student-Athlete Mental Health [Internet]*. Available at: www.ncaa.org/sport-science-institute/introduction-mind-body-and-sport. Accessed January 23, 2018.
 107. Stoltenburg AL, Kamphoff CS, Lindstrom Bremer KL. Transitioning out of sport: the psychosocial effects of collegiate athletes' career-ending injuries. *Athletic Insight J*. 2011;3:115–133.
 108. Kleiber DA, Brock SC. The effect of career-ending injuries on the subsequent well-being of elite college athletes. *Sociol Sport J*. 1992;9:70–75.
 109. Cadigan JM, Littlefield AK, Martens MP, et al. Transitions into and out of intercollegiate athletic involvement and risky drinking. *J Stud Alcohol Drugs*. 2013;74:21–29.
 110. Lavalley D, Robinson HK. In pursuit of an identity: a qualitative exploration of retirement from women's artistic gymnastics. *Psychol Sport Exerc*. 2007;8:119–141.
 111. Arvinen-Barrow M, Hurley D, Ruiz MC. Transitioning out of professional sport: the psychosocial impact of career-ending injuries among elite Irish Rugby Football Union players. *J Clin Sport Psychol*. 2017;11:67–84.
 112. National Collegiate Athletic Association. *Interassociation Consensus Document: Understanding and Supporting Student-Athlete Mental Wellness. Mental Health Best Practices*. Available at: http://www.ncaa.org/sites/default/files/SSL_MentalHealthBestPractices_Web_20170921.pdf. Accessed February 18, 2018.
 113. Neal TL, Diamond AB, Goldman S, et al. Inter-Association recommendations for developing a plan to recognize and refer student-athletes with psychological concerns at the collegiate level: an executive summary of a consensus statement. *J Athl Train*. 2013;48: 716–720.
 114. Putukian M. The psychological response to injury in student athletes: a narrative review with a focus on mental health. *Br J Sports Med*. 2016; 50:145–148.
 115. Rice SM, Purcell R, DeSilva S, et al. The mental health of elite athletes: a narrative systematic review. *Sports Med*. 2016;46:1333–1353.

116. Ivarsson A, Johnson U, Anderson MB, et al. Psychosocial factors and sport injuries: meta-analyses for prediction and prevention. *Sports Med.* 2017;47:353–365.
117. Ivarsson A, Johnson U, Podlog L. Psychological predictors of injury occurrence: a prospective investigation of professional Swedish soccer players. *J Sport Rehabil.* 2013;22:19–26.
118. Ivarsson A, Johnson U, Lindwall M, et al. Psychosocial stress as a predictor of injury in elite junior soccer: a latent growth curve analysis. *J Sci Med Sport.* 2014;17:366–370.
119. Ardern CL, Taylor NF, Feller JA, et al. A systematic review of the psychological factors associated with returning to sport following injury. *Br J Sport Med.* 2013;47:1120–1126.
120. Wiese-Bjornstal DM. Psychology and socioculture affect injury risk, response, and recovery in high-intensity athletes: a consensus statement. *Scand J Med Sci Sports.* 2010;20(suppl 2):103–111.
121. Wiese-Bjornstal DM, Smith AM, Shaffer SM, et al. An integrated model of response to sport injury: psychological and sociological dynamics. *J Appl Sport Psychol.* 1998;10:46–69.
122. Nippert AH, Smith AM. Psychological stress related to injury and impact on sport performance. *Phys Med Rehabil Clin N Am.* 2008;19:399–418.
123. Glazer JL. Eating disorders among male athletes. *Curr Sports Med Rep.* 2008;7:332–337.
124. Nattiv A, Loucks AB, Manore MM, et al. American College of Sports Medicine position stand. The female athlete triad. *Med Sci Sports Exerc.* 2007;39:1867–1882.
125. Nattiv A, Puffer JC, Green GA. Lifestyles and health risks of collegiate athletes: a multi-center study. *Clin J Sports Med.* 1997;7:262–272.
126. Armstrong S, Omen-Early J: social connectedness, self-esteem, and depression symptomatology among collegiate athletes versus non-athletes. *J Am Coll Health.* 2009;57:521–526.
127. Miller BE, Miller MN, Verheghe R, et al. Alcohol misuse among college athletes: self medication for psychiatric symptoms? *J Drug Educ.* 2002;32:41–52.
128. Guskiewicz KM, Marshall SW, Bailes J, et al. Recurrent concussion and risk of depression in retired professional football players. *Med Sci Sports Exerc.* 2007;39:903–909.
129. Kerr ZY, Marshall SW, Harding HP, et al. Nine-year risk of depression diagnosis increases with increasing self-reported concussions in retired professional football players. *Am J Sports Med.* 2012;40:2206–2212.
130. Smith AM, Milliner EK. Injured athletes and the risk of suicide. *J Athl Train.* 1994;29:337–341.
131. Kellman M. Preventing overtraining in athletes in high-intensity sports and stress/recovery monitoring. *Scand J Med Sci Sports.* 2010;20(suppl 2):95–102.
132. Staufenbiel SM, Penninx BW, Spijker AT, et al. Hair cortisol, stress exposure, and mental health in humans: a systematic review. *Psychoneuroendocrinology.* 2013;38:1220–1235.
133. Beals KA, Manore MM. Disorders of the female athlete triad among collegiate athletes. *Int J Sport Nutr Exerc Metab.* 2002;12:281–293.
134. Weigand S, Cohen J, Merenstein D. Susceptibility for depression in current and retired student athletes. *Sports Health.* 2013;5:263–266.
135. Lindqvist AS, Moberg T, Ehmborg C, et al. Increased mortality rate and suicide in Swedish former elite male athletes in power sports. *Scand J Med Sci Sports.* 2014;24:1000–1005.
136. Huang JH, Jacobs DF, Derevensky JL, et al. Gambling and health risk behaviors among US college student-athletes: findings from a national study. *J Adolesc Health.* 2007;40:390–397.
137. Weinstock J, Whelan JP, Meyers AW, et al. Gambling behavior of student-athletes and a student cohort: what are the odds? *J Gambl Stud.* 2007;23:13–24.
138. Wolanin A, Gross M, Hong E. Depression in athletes: prevalence and risk factors. *Curr Sports Med Rep.* 2015;14:56–60.
139. Forsdyke D, Smith A, Jones M, et al. Psychosocial factors associated with outcomes of sports injury rehabilitation in competitive athletes: a mixed studies systematic review. *Br J Sports Med.* 2016;50:537–544.
140. Edvardsson A, Ivarsson A, Johnson U. Is a cognitive behavioural feedback intervention useful to reduce injury risk in junior football players? *J Sports Sci Med.* 2012;11:331–338.
141. Pensgaard AM, Ivarsson A, Nilstad A, et al. Psychosocial stress factors, including the relationship with the coach, and their influence on acute and overuse injury risk in elite female football players. *BMJ Open Sport Exerc Med.* 2018;4:e000317.
142. van der Does HTD, Brink MS, Otter RTA, et al. Injury risk is increased by changes in perceived recovery of team sport players. *Clin J Sport Med.* 2017;27:46–51.
143. Nicholls AR, Levy AR, Grice A, et al. Stress appraisals, coping, and coping effectiveness among international cross-country runners during training and competition. *Eur J Sport Sci.* 2009;9:285–293.
144. Devantier C. Psychological predictors of injury among professional soccer players. *Sport Sci Rev.* 2011;20:5–36.
145. Ardern CL. Anterior cruciate ligament reconstruction-not exactly a one-way ticket back to the preinjury level: a review of contextual factors affecting return to sport after surgery. *Sports Health.* 2015;7:224–230.
146. Czuppon S, Racette BA, Klein SE, et al. Variables associated with return to sport following anterior cruciate ligament reconstruction: a systematic review. *Br J Sports Med.* 2014;48:356–364.
147. Glazer DD. Development and preliminary validation of the injury-psychological readiness to return to sport (I-PRRS) scale. *J Athl Train.* 2009;44:185–189.
148. Tripp DA, Stanish W, Ebel-Lam A, et al. Fear of reinjury, negative affect, and catastrophizing predicting return to sport in recreational athletes with anterior cruciate ligament injuries at 1 year postsurgery. *Rehabil Psychol.* 2007;52:74–81.
149. Gulliver A, Griffiths KM, Mackinnon A, et al. The mental health of Australian elite athletes. *J Sci Med Sport.* 2015;18:255–261.
150. Proctor SL, Boan-Lenzo C. Prevalence of depressive symptoms in male intercollegiate student-athletes and nonathletes. *J Clin Sport Psych.* 2010;4:204–220.
151. Hammond T, Gialloreti C, Kubas H, et al. The prevalence of failure-based depression among elite athletes. *Clin J Sport Med.* 2013;23:273–277.
152. Gignac M, Cao X, Ramanathan S, et al. Perceived personal importance of exercise and fears of re-injury: a longitudinal study of psychological factors related to activity after anterior cruciate ligament reconstruction. *BMC Sports Sci Med Rehabil.* 2015;7:4.
153. Flanagan DC, Everhart JS, Glassman AH. Psychological factors affecting rehabilitation and outcomes following elective orthopaedic surgery. *J Am Acad Orthop Surg.* 2015;23:563–570.
154. Podlog L, Dimmock J, Miller J. A review of return to sport concerns following injury rehabilitation: Practitioner strategies for enhancing recovery outcomes. *Phys Ther Sport.* 2011;12:36–42.
155. Podlog LW, Banham SM, Wadey R, et al. Psychological readiness to return to competitive sport following injury: a qualitative study. *Sport Psychol.* 2015;29:1–14.
156. Ardern CL, Taylor NF, Feller JA, et al. Sports participation 2 years after anterior cruciate ligament reconstruction in athletes who had not returned to sport at 1 year: a prospective follow-up of physical function and psychological factors in 122 athletes. *Am J Sports Med.* 2015;43:848–856.
157. Martens MP, Dam-O'Connor K, Beck NC. A systematic review of college student-athlete drinking: prevalence rates, sport-related factors, and interventions. *J Subst Abuse Treat.* 2006;31:305–316.
158. NCAA Student-Athlete Substance Use Study: Executive Summary August 2014. NCAA.org. Available at: <http://www.ncaa.org/about/resources/research/ncaa-student-athlete-substance-use-study-executive-summary-august-2014>. Accessed June 21, 2018.
159. Gil F, Guerra de Andrade A, Castaldelli-Maia JM. Discussing prevalence, impacts and treatment of substance use disorders in athletes. *Int Rev Psychiatry.* 2016;28:572–578.
160. Brisola-Santos MB, Mello e Gallinaro JG, Sampaio-Junior B, et al. Prevalence and correlates of cannabis use among athletes: a review. *Am J Addict.* 2016;25:518–528.
161. Crisp A, Sedgwick P, Halek C, et al. Why may teenage girls persist in smoking? *J Adolesc.* 1999;22:657–672.
162. Stice E, Hayward C, Cameron RP, et al. Body-image and eating disturbances predict onset of depression among female adolescents: a longitudinal study. *J Abnorm Psychol.* 2000;109:43844.
163. Crow S, Eisenberg ME, Story M, et al. Psychosocial and behavioral correlates of dieting among overweight and non-overweight adolescents. *J Adolesc Health.* 2006;38:569–574.
164. Green GA, Uryasz FD, Petr TA, et al. NCAA study of substance use and abuse habits of college student-athletes. *Clin J Sport Med.* 2001;11:51–56.
165. Yusko DA, Buckman JF, White HR, et al. Risk for excessive alcohol use and drinking-related problems in college student athletes. *Addict Behav.* 2008;33:1546–1556.
166. US Department of Health and Human Services. HHS Acting Secretary Declares Public Health Emergency to Address National Opioid Crisis. 2017. Available at: <https://www.hhs.gov/about/news/2017/10/26/hhs-acting-secretary-declares-public-health-emergency-address-national-opioid-crisis.html>. Accessed September 16, 2018.

167. Veliz P, Epstein-Ngo QM, Meier E, et al. Painfully obvious: a longitudinal examination of medical use and misuse of opioid medication among adolescent sports participants. *J Adolesc Health*. 2014;54:333–340.
168. Veliz P, Epstein-Ngo QM, Austic E, et al. Opioid use among interscholastic sports participants: an exploratory study from a sample of college students. *Res Q Exerc Sport*. 2015;86:205–211.
169. Veliz P, Boyd CJ, McCabe SE. Nonmedical prescription opioid and heroin use among adolescents who engage in sports and exercise. *Pediatrics*. 2016;138:e20160677.
170. McCabe SE, West BT, Teter JT, et al. Characteristics associated with the diversion of controlled medications among adolescents. *Drug and Alcohol Depend*. 2011;118:452–458.
171. Cottler LB, Abdallah AB, Cummings SM, et al. Injury, pain, and prescription opioid use among former National Football League (NFL) players. *Drug Alcohol Depend*. 2011;116:188–194.
172. Veliz P, Boyd CJ, McCabe SE. Nonmedical use of prescription opioids and heroin use among adolescents involved in competitive sports. *J Adolesc Health*. 2017;60:346–349.
173. Veliz PT, Boyd C, McCabe SE. Playing through pain: sports participation and non-medical use of opioid medications among adolescents. *Am J Public Health*. 2013;103:e28–e30.
174. Tricker R. Painkilling drugs in collegiate athletics: knowledge, attitudes and use of student athletes. *J Drug Educ*. 2000;30:313–324.
175. Compton WM, Jones CM, Baldwin GT. Relationship between nonmedical prescription-opioid use and heroin use. *N Engl J Med*. 2016;374:154–163.
176. Kelly MA. Addressing the opioid epidemic with multimodal pain management. *Am J Orthop*. 2016;45:S6–S8.
177. Levy N, Sturgess J, Mills P. “Pain as the fifth vital sign” and dependence on the “numerical pain scale” is being abandoned in the US: why? *Br J Anesth*. 2018;120:435–438.
178. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 5th ed. Washington, DC: American Psychiatric Association; 2013.
179. Joy E, Kussman A, Nattiv A. Update on eating disorders in athletes: a comprehensive narrative review with a focus on clinical assessment and management. *Br J Sports Med*. 2016;50:154–162.
180. De Souza MJ, Nattiv A, Joy E, et al. 2014 female athlete triad coalition consensus statement on treatment and return to play of the female athlete triad: 1st International Conference held in San Francisco, CA, May 2012 and 2nd International Conference held in Indianapolis, IN, May 2013. *Clin J Sport Med*. 2014;24:96–119.
181. Mountjoy M, Sundgot-Borgen J, Burke L, et al. The IOC consensus statement: beyond the female athlete triad—relative energy deficiency in sport (RED-S). *Br J Sports Med*. 2014;48:491–497.
182. Arthur-Cameselle J, Sossin K, Quatromoni P. A qualitative analysis of factors related to eating disorder onset in female collegiate athletes and non-athletes. *Eat Disord*. 2017;25:199–215.
183. Knapp J, Aerni G, Anderson J. Eating disorders in female athletes: use of screening tools. *Curr Sports Med Rep*. 2014;13:214–218.
184. Plateau CR, Arcelus J, Meyer C. Detecting eating psychopathology in female athletes by asking about exercise: use of the compulsive exercise test. *Eur Eat Disord Rev*. 2017;25:618–624.
185. Morgan JF, Reid F, Lacey JH. The SCOFF questionnaire: assessment of a new screening tool for eating disorders. *BMJ*. 1999;319:1467–1468.
186. Fairburn CG, Cooper Z, O’Connor ME. Eating disorder examination (edition 16.0D). In: Fairburn CG, ed. *Cognitive Behavior Therapy and Eating Disorders*. New York, NY: The Guilford Press; 2008:265–308.
187. American Academy of Family Physicians, American Academy of Pediatrics, American College of Sports Medicine, American Medical Society for Sports Medicine, American Orthopaedic Society for Sports Medicine, American Osteopathic Academy of Sports Medicine. *PPE Preparation Physical Evaluation*. 4th ed. Minneapolis, MN: McGraw-Hill; 2010.
188. Couturier J, Kimber M, Szatmari P. Efficacy of family-based treatment for adolescents with eating disorders: a systematic review and meta-analysis. *Int J Eat Disord*. 2013;46:3–11.
189. Loucas CE, Fairburn CG, Whittington C, et al. E-therapy in the treatment and prevention of eating disorders: a systematic review and meta-analysis. *Behav Res Ther*. 2014;63:122–131.
190. David H, Attia E. Pharmacotherapy of eating disorders. *Curr Opin Psych*. 2017;30:452–457.
191. Cook BJ, Wonderlich SA, Mitchell JE, et al. Exercise in eating disorders treatment: systematic review and proposal of guidelines. *Med Sci Sports Exerc*. 2016;48:1408–1414.
192. Johnson MD. Disordered eating in active and athletic women. *Clin Sports Med*. 1994;13:355–369.
193. Fogelholm M. Effects of bodyweight reduction on sports performance. *Sports Med*. 1994;18:249–267.
194. El Ghoch M, Soave F, Calugi S, et al. Eating disorders, physical fitness and sport performance: a systematic review. *Nutrients*. 2013;5:5140–5160.
195. Tenforde AS, Carlson JL, Chang A, et al. Association of the female athlete triad risk assessment stratification to the development of bone stress injuries in collegiate athletes. *Am J Sports Med*. 2017;45:302–310.
196. Martinson M, Bahr R, Børrensen R, et al. Preventing eating disorders among young elite athletes: a randomized controlled trial. *Med Sci Sports Exerc*. 2014;46:435–447.
197. Bar RJ, Cassin SE, Dionne MM. Eating disorder prevention initiatives for athletes: a review. *Eur J Sport Sci*. 2016;16:325–335.
198. Wolanin A, Hong E, Marks D, et al. Prevalence of clinically elevated depressive symptoms in college athletes and differences by gender and sport. *Br J Sports Med*. 2016;50:167–171.
199. Yang J, Peek-Asa C, Corlette JD. Prevalence of and risk factors associated with symptoms of depression in competitive collegiate student athletes. *Clin J Sports Med*. 2007;17:481–487.
200. Smarr KL, Keefer AL. Measures of depression and depressive symptoms. *Arthritis Care*. 2011;63:S454–S466.
201. Reardon CL, Creado S. Psychiatric medication preferences of sports psychiatrists. *Phys Sportsmed*. 2016;44:397–402.
202. Rao AL, Asif IM, Drezner JA, et al. Suicide in national collegiate athletic association athletes: a 9-year analysis of the NCAA resolutions database. *Sports Health*. 2015;7:452–457.
203. Lehman EJ, Hein MJ, Gersic CM. Suicide mortality among retired national football league players who played 5 or more seasons. *Am J Sports Med*. 2016;44:2486–2491.
204. Spitzer R, Kroenke K, Williams JBW, et al. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med*. 2006;166:1092–1097.
205. Kessler RC, Berglund P, Demler O, et al. Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psych*. 2005;62:593–602.
206. Schaal K, Tafflet M, Nassif H, et al. Psychological balance in high level athletes: gender-based differences and sport-specific patterns. *PLoS One*. 2011;6:e19007.
207. Norton PJ, Burn JA, Hope DA, et al. Generalization of social anxiety to sporting and athletic situations: gender, sports involvement, and parental pressure. *Depress Anxiety*. 2000;12:193–202.
208. Kessler RC, Petukhova M, Sampson NA, et al. Twelve-month and lifetime prevalence and lifetime morbid risk of anxiety and mood disorders in the United States. *Int J Methods Psych Res*. 2012;21:169–184.
209. Storch EA, Storch JB, Killiany EM, et al. Self-reported psychopathology in athletes: a comparison of intercollegiate student-athletes and non-athletes. *J Sport Behav*. 2005;28:86–98.
210. Cromer L, Kaier E, Davis J, et al. OCD in college athletes. *Am J Psychiatry*. 2017;174:595–597.
211. Appaneal RN, Perna FM, Larkin KT. Psychophysiological response to severe sport injury among competitive male athletes: a preliminary investigation. *J Clin Sport Psychol*. 2007;1:68–88.
212. McArdle S. Psychological rehabilitation from anterior cruciate ligament–medial collateral ligament reconstructive surgery: a case study. *Sports Health*. 2010;2:73–77.
213. Gouttebarga V, Frings-Dresen MHW, Sluiter JK. Mental and psychosocial health among current and former professional footballers. *Occup Med*. 2015;65:190–196.
214. Reardon CL. The sports psychiatrist and psychiatric medication. *Intl Rev Psychiatry*. 2016;28:606–613.
215. Baron DA, Reardon CL, Baron SH, eds. *Clinical Sports Psychiatry: An International Perspective*. Sussex, United Kingdom: John Wiley & Sons; 2013.
216. Patel DR, Omar H, Terry M. Sport-related performance anxiety in young female athletes. *J Pediatr Adolesc Gynecol*. 2010;23:325–335.
217. Reardon CL, Factor RM. Sport psychiatry. *Sports Med*. 2010;40:961–980.
218. Baum AL. Psychopharmacology in athletes. In Begel D and Burton R, eds. *Sport Psychiatry: Theory and Practice*. New York, NY: W.W. Norton Publishing; 2000:249–259.

219. Otte C. Cognitive behavioral therapy in anxiety disorders: current state of the evidence. *Dialogues Clin Neurosci*. 2011;13:413–421.
220. Yerkes RM, Dodson JD. The relation of strength of stimulus to rapidity of habit formation. *J Comp Neurol Psychol*. 1908;18:459–482.
221. Kamm RL. Interviewing principles for the psychiatrically aware sports medicine physician. *Clin Sports Med*. 2005;24:745–769.
222. Mahoney J, Hanrahan SJ. A brief educational intervention using acceptance and commitment therapy: four injured athletes' experiences. *J Clin Sport Psychol*. 2011;5:252–273.
223. Shortway KM, Wolanin A, Block-Lerner J, et al. Acceptance and commitment therapy for injured athletes: development and preliminary Feasibility of the return to ACTion protocol. *J Clin Sport Psychol*. 2018;12:4–26.
224. Meeusen R, Duclos M, Foster C, et al. Prevention, diagnosis, and treatment of the overtraining syndrome: joint consensus statement of the European College of Sport Science and the American College of Sports Medicine. *Med Sci Sports Exerc*. 2013;45:186–205.
225. Schweltnus M, Soligard T, Alonso JM, et al. How much is too much? (part 2) International Olympic Committee consensus statement on load in sport and risk of illness. *Br J Sports Med*. 2016;50:1043–1052.
226. Urhausen A, Gabriel HH, Weiler B, et al. Ergometric and psychological findings during overtraining: a long-term follow-up study in endurance athletes. *Int J Sports Med*. 1998;19:114–120.
227. Cadejani FA, Kater CE. Hormonal aspects of overtraining syndrome: a systematic review. *BMC Sports Sci Med Rehabil*. 2017;9:14.
228. Le Meur Y, Hausswirth C, Natta F, et al. A multidisciplinary approach to overreaching detection in endurance trained athletes. *J Appl Physiol*. 2013;114:411–420.
229. Raglin JS, Morgan WP. Development of a scale for use in monitoring training-induced distress in athletes. *Int J Sports Med*. 1994;15:84–88.
230. Kenttä G, Hassmén P, Raglin J. Mood state monitoring of training and recovery in elite kayakers. *Eur J Sport Sci*. 2006;4:245–253.
231. Carfagno D, Hendrix J. Overtraining syndrome in the athlete: current clinical practice. *Curr Sports Med Rep*. 2014;13:45–51.
232. Madigan DJ, Stoeber J, Passfield L. Perfectionism and training distress in junior athletes: a longitudinal investigation. *J Sports Sci*. 2017;35:470–475.
233. Lewis N, Collins D, Pedlar C, et al. Can clinicians and scientists explain and prevent unexplained underperformance syndrome in elite athletes: an interdisciplinary perspective and 2016 update. *Br Med J Open Sport Exerc Med*. 2015;1:1–10.
234. Brink MS, Frencken WGP, Jordet G, et al. Coaches' and players' perceptions of training dose: not a perfect match. *Int J Sports Physiol Perform*. 2014;9:497–502.
235. Berglund B, Säfström H. Psychological monitoring and modulation of training load of world-class canoeists. *Med Sci Sports Exerc*. 1994;26:1036–1040.
236. Byrd KL, Gelaye B, Tadesse MG, et al. Sleep disturbances and common mental disorders in college students. *Health Behav Policy Rev*. 2014;1:229–237.
237. Drew M, Vlahovich N, Hughes D, et al. Prevalence of illness, poor mental health and sleep quality and low energy availability prior to the 2016 Summer Olympic Games. *Br J Sports Med*. 2018;52:47–53.
238. Swinbourne R, Gill N, Vaile J, et al. Prevalence of poor sleep quality, sleepiness and obstructive sleep apnoea risk factors in athletes. *Eur J Sport Sci*. 2016;16:850–858.
239. Silva A, Queiroz SS, Winckler C, et al. Sleep quality evaluation, chronotype, sleepiness and anxiety of Paralympic Brazilian athletes: Beijing 2008 Paralympic Games. *Br J Sports Med*. 2012;46:150–154.
240. Stranks EK, Crowe SF. The cognitive effects of obstructive sleep apnea: an updated meta-analysis. *Arch Clin Neuropsychol*. 2016;31:186–193.
241. Haddock N, Wells ME. The association between treated and untreated obstructive sleep apnea and depression. *Neurodiagn J*. 2018;58:30–39.
242. Dobrosielski DA, Nichols D, Ford J, et al. Estimating the prevalence of sleep-disordered breathing among collegiate football players. *Respir Care*. 2016;61:1144–1150.
243. George CF, Kab V. Sleep-disordered breathing in the National Football League is not a trivial matter. *Sleep*. 2011;34:245.
244. Bender AM, Lawson D, Werthner P, et al. The clinical validation of the athlete sleep screening questionnaire: an instrument to identify athletes that need further sleep assessment. *Sports Med Open*. 2018;4:23.
245. Gagnon C, Bélanger L, Ivers H, et al. Validation of the insomnia severity index in primary care. *Am Board Fam Med*. 2013;26:701–710.
246. Chiu HY, Chen PY, Chuang LP, et al. Diagnostic accuracy of the Berlin questionnaire, STOP-BANG, STOP, and Epworth sleepiness scale in detecting obstructive sleep apnea: a bivariate meta-analysis. *Sleep Med Rev*. 2017;36:57–70.
247. Taylor DJ, Zimmerman MR, Gardner CE, et al. A pilot randomized controlled trial of the effects of cognitive-behavioral therapy for insomnia on sleep and daytime functioning in college students. *Behav Ther*. 2014;45:376–389.
248. Carney CE, Edinger JD, Kuchibhatla M, et al. Cognitive behavioral insomnia therapy for those with insomnia and depression: a randomized controlled clinical trial. *Sleep*. 2017;40.
249. Juliff LE, Halson SL, Peiffer JJ. Understanding sleep disturbance in athletes prior to important competitions. *J Sci Med Sport*. 2015;18:13–18.
250. Mah CD. Extended sleep and the effects on mood and athletic performance in collegiate swimmers. *Sleep*. 2008;31:A128.
251. Mah CD, Mah KE, Kezirian EJ, et al. The effects of sleep extension on the athletic performance of collegiate basketball players. *Sleep*. 2011;34:943–950.
252. Van Ryswyk E, Weeks R, Bandick L, et al. A novel sleep optimisation programme to improve athletes' well-being and performance. *Eur J Sport Sci*. 2017;17:144–151.
253. Halson SL. Sleep in elite athletes and nutritional interventions to enhance sleep. *Sports Med*. 2014;44:13–23.
254. Charles RB, Kirkham AJ, Guyatt AR, et al. Psychomotor, pulmonary and exercise responses to sleep medication. *Br J Clin Pharmacol*. 1987;24:191–197.
255. Grobler LA, Schweltnus MP, Trichard C, et al. Comparative effects of zopiclone and loperazolam on psychomotor and physical performance in active individuals. *Clin J Sport Med*. 2000;10:123–128.
256. Blin O, Micallef J, Audebert C, et al. A double-blind, placebo-and flurazepam-controlled investigation of the residual psychomotor and cognitive effects of modified release zolpidem in young healthy volunteers. *J Clin Psychopharmacol*. 2006;26:284–289.
257. Ito SU, Kanbayashi T, Takemura T, et al. Acute effects of zolpidem on daytime alertness, psychomotor and physical performance. *Neurosci Res*. 2007;59:309–313.
258. Paul MA, Gray G, Kenny G, et al. Impact of melatonin, zaleplon, zopiclone, and temazepam on psychomotor performance. *Aviat Space Environ Med*. 2003;74:1263–1270.
259. Buscemi N, Vandermeer B, Hooton N, et al. The efficacy and safety of exogenous melatonin for primary sleep disorders. *J Gen Intern Med*. 2005;20:1151–1158.
260. Atkinson G, Buckley P, Edwards B, et al. Are there hangover-effects on physical performance when melatonin is ingested by athletes before nocturnal sleep? *Int J Sports Med*. 2001;22:232–234.
261. Atkin T, Comai S, Gobbi G. Drugs for insomnia beyond benzodiazepines: pharmacology, clinical applications, and discovery. *Pharmacol Rev*. 2018;70:197–245.
262. Walsh JK, Erman M, Erwin CW, et al. Subjective hypnotic efficacy of trazodone and zolpidem in DSM-III-R primary insomnia. *Hum Psychopharmacol*. 1998;13:191–198.
263. Sateia MJ, Buysse DJ, Krystal AD, et al. Clinical practice guideline for the pharmacologic treatment of chronic insomnia in adults: an American Academy of Sleep Medicine clinical practice guideline. *J Clin Sleep Med*. 2017;13:307–349.
264. Fullagar HH, Skorski S, Duffield R, et al. Sleep and athletic performance: the effects of sleep loss on exercise performance, and physiological and cognitive responses to exercise. *Sports Med*. 2015;45:161–186.
265. Haack M, Mullington JM. Sustained sleep restriction reduces emotional and physical well-being. *Pain*. 2005;119:56–64.
266. Lautenbacher S, Kundermann B, Krieg JC. Sleep deprivation and pain perception. *Sleep Med Rev*. 2006;10:357–369.
267. Cadejani FA, Kater CE. Body composition, metabolism, sleep, psychological and eating patterns of overtraining syndrome: results of the EROS study (EROS-PROFILE). *J Sports Sci*. 2018;11:1–9.
268. Lastella M, Lovell GP, Sargent C. Athletes' precompetitive sleep behaviour and its relationship with subsequent precompetitive mood and performance. *Eur J Sport Sci*. 2014;14:S123–S130.
269. Meerlo P, Sgoifo A, Suchecki D. Restricted and disrupted sleep: effects on autonomic function, neuroendocrine stress systems and stress responsivity. *Sleep Med Rev*. 2008;12:197–210.
270. Paiva T, Gaspar T, Matos MG. Mutual relations between sleep deprivation, sleep stealers and risk behaviours in adolescents. *Sleep Sci*. 2016;9:7–13.
271. Hildenbrand AK, Daly BP, Nicholls E, et al. Increased risk for school violence-related behaviors among adolescents with insufficient sleep. *J Sch Health*. 2013;83:408–414.
272. Taylor DJ, Bramoweth AD. Patterns and consequences of inadequate sleep in college students: substance use and motor vehicle accidents. *J Adolesc Health*. 2010;46:610–612.

273. Chang AM, Aeschbach D, Duffy JF, et al. Evening use of light-emitting eReaders negatively affects sleep, circadian timing, and next-morning alertness. *Proc Natl Acad Sci*. 2015;112:1232–1237.
274. Myllymäki T, Kyröläinen H, Savolainen K, et al. Effects of vigorous late-night exercise on sleep quality and cardiac autonomic activity. *J Sleep Res*. 2011;20:146–153.
275. Buman MP, Phillips BA, Youngstedt SD, et al. Does nighttime exercise really disturb sleep? Results from the 2013 national sleep Foundation sleep in America Poll. *Sleep Med*. 2014;15:755–761.
276. Brand S, Gerber M, Beck J, et al. High exercise levels are related to favorable sleep patterns and psychological functioning in adolescents: a comparison of athletes and controls. *J Adolesc Health*. 2010;46:133–141.
277. Putukian M, Kreher JB, Coppel DB, et al. Attention deficit hyperactivity disorder and the athlete: an American Medical Society for Sports Medicine position statement. *Clin J Sport Med*. 2011;21:392–401.
278. Stewman CG, Liebman C, Fink L, et al. Attention deficit Hyperactivity disorder: unique considerations in athletes. *Sports Health*. 2018;10:40–46.
279. Pujalte GGA, Maynard JR, Thurston MJ, et al. Considerations in the care of athletes with attention deficit hyperactivity disorder. *Clin J Sports Med*. 2019;29:245–256.
280. Reardon CL, Factor RM. Considerations in the use of stimulants in sport. *Sports Med*. 2016;46:611–617.
281. Jadad AR, Boyle M, Cunningham C, et al. Treatment of attention-deficit/hyperactivity disorder. *Evid Rep Technol Assess*. 1999;11:i–viii, 1–341.
282. MTA Cooperative Group. 14-month randomized clinical trial of treatment strategies for attention deficit hyperactivity disorder. *Arch Gen Psychiatry*. 1999;56:1073–1086.
283. Jensen PS, Hinshaw SP, Swanson JM, et al. Findings from the NIMH multimodal treatment study of ADHD (MTA): implications and applications for primary care providers. *J Dev Behav Pediatr*. 2001;22:60–73.
284. MTA Cooperative Group. National Institute of Mental Health Multimodal Treatment Study of ADHD follow-up: 24-month outcomes of treatment strategies for attention-deficit/hyperactivity disorder. *Pediatrics*. 2004;113:754–761.
285. MTA Cooperative Group. National Institute of Mental Health Multimodal Treatment Study of ADHD follow-up: changes in effectiveness and growth after the end of treatment. *Pediatrics*. 2004;113:762–769.
286. Abikoff H, Hechtman L, Klein RG, et al. Social functioning in children with ADHD treated with long-term methylphenidate and multimodal psychosocial treatment. *J Am Acad Child Adolesc Psychiatry*. 2004;43:820–829.
287. Faraone SV, Biederman J, Jetton JG, et al. Attention deficit disorder and conduct disorder: longitudinal evidence for a familial subtype. *Psychol Med*. 1997;27:291–300.
288. Biederman J, Wilens T, Mick S, et al. Pharmacotherapy of attention-deficit/hyperactivity disorder reduces risk for substance use disorder. *Pediatrics*. 1999;104:e20.
289. Wilens TE, Faraone SV, Biederman J, et al. Does stimulant therapy of attention-deficit/hyperactivity disorder beget later substance abuse? A meta-analytic review of the literature. *Pediatrics*. 2003;111:179–185.
290. Ziareis S, Jansen P. Effects of physical activity on executive function and motor performance in children with ADHD. *Res Dev Disabil*. 2015;38:181–191.
291. Vetter VL, Elia J, Erickson C, et al. Cardiovascular monitoring of children and adolescents with heart disease receiving medications for attention deficit/hyperactivity disorder: a scientific statement from the American Heart Association Council on Cardiovascular Disease in the Young Congenital Cardiac Defects Committee and the Council on Cardiovascular Nursing. *Circulation*. 2008;117:2407–2423.
292. Perrin JM, Friedman RA, Knilans TK, et al. Cardiovascular monitoring and stimulant drugs for attention-deficit/hyperactivity disorder. *Pediatrics*. 2008;122:451–453.
293. American Academy of Pediatrics, Committee on Quality Improvement, Subcommittee on Attention-Deficit/Hyperactivity Disorder. Clinical practice guideline: diagnosis and evaluation of the child with attention-deficit/hyperactivity disorder. *Pediatrics*. 2000;105:1158–1170.
294. American Academy of Pediatrics, Subcommittee on Attention-Deficit/Hyperactivity Disorder and Committee on Quality Improvement. Clinical practice guideline: treatment of school-aged child with attention-deficit/hyperactivity disorder. *Pediatrics*. 2001;108:1033–1044.
295. Drezner JA. Sudden cardiac death in young athletes: causes, athlete's heart, and screening guidelines. *Postgrad Med*. 2000;108:37–44, 47–50.
296. Wilens TE, Hammerness PG, Biederman J, et al. Blood pressure changes associated with medication treatment of adults with attention-deficit/hyperactivity disorder. *J Clin Psychiatry*. 2005;66:253–259.
297. Knight M. Stimulant-drug therapy for attention-deficit disorder (with or without hyperactivity) and sudden cardiac death. *Pediatrics*. 2007;119:154–155.
298. Cooper WO, Habel LA, Sox CM, et al. ADHD drugs and serious cardiovascular events in children and adults. *N Engl J Med*. 2011;365:1896–1904.
299. International Olympic Committee (IOC). *Anti Doping Rules*. Available at: <https://www.olympic.org/~media/Document%20Library/OlympicOrg/IOC/What-We-Do/Protecting-Clean-Athletes/Fight-against-doping/EN-Anti-Doping-Rules-PyeongChang2018.pdf?la=en>. Accessed February 1, 2019.
300. NCAA Drug Testing Program. Available at: <http://www.ncaa.org/sport-science-institute/ncaa-drug-testing-program>. Accessed August 8, 2018.
301. *World Anti-doping Agency Drug Testing*. Available at: <https://www.wada-ama.org/>. Accessed August 8, 2018.
302. Pelham WE Jr, McBurnett K, Harper GW, et al. Methylphenidate and baseball playing in ADHD children: who's on first? *J Consult Clin Psychol*. 1990;58:130–133.
303. Jacobs I, Bell DG. Effects of acute modafinil ingestion on exercise time to exhaustion. *Med Sci Sports Exerc*. 2004;36:1078–1082.
304. Roelands B, Hasegawa H, Watson P, et al. The effects of acute dopamine reuptake inhibition on performance. *Med Sci Sports Exerc*. 2008;40:879–885.
305. Watson P, Hasegawa H, Roelands B, et al. Acute dopamine/noradrenaline reuptake inhibition enhances human exercise performance in warm, but not temperate conditions. *J Physiol*. 2005;565:873–883.
306. McLellan TM, Ducharme MB, Canini F, et al. Effect of modafinil on core temperature during sustained wakefulness and exercise in a warm environment. *Aviat Space Environ Med*. 2002;73:1079–1088.
307. DeSantis AD, Webb EM, Noar SM, et al. Illicit use of prescription ADHD medications on a college campus: a multimethodological approach. *J Am Coll Health*. 2008;57:315–323.
308. DuPont RL, Coleman JJ, Bucher RH, et al. Characteristics and motives of college students who engage in nonmedical use of methylphenidate. *Am J Addict*. 2008;17:167–171.
309. American College Association. American College Health Association National College Assessment (ACHA-NCHA): Spring 2005 Reference Group Data Report (Abridged). *J Amer Coll Health*. 2006;54:5–16.