

Supplemental Material

Supplemental table 1 – Inflammatory markers in COVID-19 patients, on admission, and at time of AKI (acute kidney injury) onset.

Cas e #	D-Dimer (<229ng/ml DDU)	D-Dimer (<229ng/ml DDU)	Ferritin (15-150ng/ml)	Ferritin (15-150ng/ml)	C-Reactive Protein (0.00-0.4mg/dl)	C-Reactive Protein (0.00-0.4mg/dl)	Procalcitonin (0.02-0.10ng/ml)	Procalcitonin (0.02-0.10ng/ml)
	On Admission	At the time of AKI	On Admission	At the time of AKI	On Admission	At the time of AKI	On Admission	At the time of AKI
1	805		1105		84.8		2.02	
2	165	417	275	377	4.23	7.65	0.13	0.99
3	308	1168	1693	4839	12.07	52.77	0.17	5.82
4	622	13212	1728	2239	19.62	14.97	1.16	0.57
5	1133	2576	NA	430.2	205.4	116.7	NA	0.4
6	NA		NA		NA		NA	
7	411	5927	189	2517	10.35	8.02	0.11	2.1
8	1353		1985		14.53		1.6	
9	1147		269.4		12.3		NA	
10	2158		1912		173.2		NA	

Supplementary table 1: Inflammatory markers in COVID-19 patients, on admission, and at time of AKI (acute kidney injury) onset. Patients 1,6,8,9,10 presented with AKI at the time of admission.