

## Comparison between placenta accreta scoring system, ultrasound staging and clinical classification

Xinrui Yang<sup>1</sup> (Email: [yangxinrui9695@sina.com](mailto:yangxinrui9695@sina.com) Phone: +86 13521095118)

Weiran Zheng<sup>1</sup> Jie Yan<sup>1</sup> Huixia Yang<sup>1\*</sup>

<sup>1</sup> Department of Obstetrics and Gynecology, Peking University First Hospital  
Beijing 100034, China

\*Corresponding Author: Huixia Yang, Email: [yanghuixia@bjmu.edu.cn](mailto:yanghuixia@bjmu.edu.cn)

**Table S1.** Placenta accreta scoring system\*

	Items	0 point	1 point	2 points	3 points
Risk factors	Previous abortion or miscarriage	No	Yes	-	-
	Previous cesarean sections	0	1	2	≥3
	Placenta previa	No	Marginal	Partial	Complete
	Multiple vascular lacunae	No	Yes	Swiss-cheese appearance	-
Ultrasound	Disruption of the myometrium	Continuous	Disrupted	Disappeared	-
	Boundaries between the placenta and the myometrium	Clear	Obscure	Disappeared	-
Features	Uterine serosa-posterior bladder wall interface	Clear	Obscure	Disappeared	-
	Sub-placental hypervascularity	No	Turbulent blood flow	Bridging vessels	-
	Extending into the bladder	No	Yes	-	-
	Extending into the cervix	No	Yes	-	-

\* adapted from Weiran Zheng et al.