SDC 3. Pearson correlation matrix of the associations among changes in food reward, eating behavior traits and body composition in the whole group

		ΔLiking	ΔWanting	ΔRestraint	ΔDisinhibition	ΔSuscep. hunger	ΔBinge eating	ΔBMI	ΔBody weight	ΔBody fat percentage	ΔFat mass	ΔFat-free mass
ΔLiking	*r*	1	0.458	-0.128	0.022	0.027	0.099	0.004	0.017	0.006	0.054	-0.085
	p-value		<0.001	0.344	0.872	0.845	0.468	0.973	0.900	0.962	0.682	0.523
ΔWanting	*r*		1	0.015	0.117	0.206	0.254	0.082	0.098	0.046	0.076	0.043
	p-value			0.915	0.386	0.124	0.059	0.535	0.458	0.728	0.569	0.746
ΔRestraint	*r*			1	0.156	-0.021	-0.199	-0.250	-0.266	-0.218	-0.271	0.028
	p-value				0.243	0.875	0.138	0.059	0.044	0.099	0.040	0.835
ΔDisinhibition	*r*				1	0.290	0.412	0.288	0.297	0.237	0.315	-0.055
	p-value					0.028	0.001	0.028	0.023	0.074	0.016	0.680
ΔSuscep. hunger	*r*					1	0.330	0.368	0.356	0.334	0.347	0.013
	p-value						0.012	0.005	0.006	0.010	0.008	0.925
ΔBinge eating	*r*						1	0.315	0.327	0.197	0.283	0.094
	p-value							0.017	0.013	0.142	0.033	0.486
ΔBMI	*r*							1	0.994	0.799	0.914	0.143
	p-value								<0.001	<0.001	<0.001	0.276
ΔBody weight	*r*								1	0.802	0.921	0.140
	p-value									<0.001	<0.001	0.286
ΔBody fat percentage	*r*									1	0.956	-0.438
	p-value										<0.001	<0.001
ΔFat mass	*r*										1	-0.257
	p-value											0.047
ΔFat-free mass	*r*											1
	p-value											