SDC 5. Pearson correlation matrix of the associations among changes in food reward, eating behavior traits and body composition in Controls

		ΔLiking	ΔWanting	ΔRestraint	ΔDisinhibition	ΔSuscep. hunger	ΔBinge eating	ΔBMI	ΔBody weight	ΔBody fat percentage	ΔFat mass	ΔFat-free mass
ΔLiking	*r*	1	0.316	-0.160	-0.084	0.179	-0.203	0.061	0.056	0.081	0.141	-0.167
	p-value		0.270	0.619	0.796	0.578	0.526	0.835	0.850	0.783	0.632	0.569
ΔWanting	*r*		1	-0.111	-0.028	0.436	0.565	-0.094	-0.113	-0.055	-0.060	-0.062
	p-value			0.730	0.930	0.156	0.055	0.749	0.701	0.851	0.839	0.834
ΔRestraint	*r*			1	0.288	0.016	-0.191	0.209	0.178	-0.057	-0.031	0.369
	p-value				0.364	0.961	0.551	0.515	0.579	0.860	0.925	0.238
ΔDisinhibition	*r*				1	-0.193	-0.049	0.498	0.495	0.623	0.563	-0.363
	p-value					0.549	0.879	0.099	0.101	0.031	0.057	0.247
ΔSuscep. hunger	*r*					1	0.239	0.163	0.134	-0.193	-0.104	0.450
	p-value						0.455	0.613	0.679	0.549	0.748	0.142
ΔBinge eating	*r*						1	-0.160	-0.168	-0.111	-0.147	0.004
	p-value							0.619	0.601	0.732	0.649	0.991
ΔBMI	*r*							1	0.997	0.764	0.874	-0.160
	p-value								<0.001	0.001	<0.001	0.584
ΔBody weight	*r*								1	0.775	0.889	-0.189
	p-value									0.001	<0.001	0.517
ΔBody fat percentage	*r*									1	0.968	-0.741
	p-value										<0.001	0.002
ΔFat mass	*r*										1	-0.617
	p-value											0.019
ΔFat-free mass	*r*											1
	p-value											