

eTable 1: Observed vs expected deaths and Intensive Care Unit admissions in the lower and higher-risk groups according to disease modifying therapy.

	Lower-risk					Higher-risk				
		Death		ICU			Death		ICU	
		Observed	Expected	Observed	Expected		Observed	Expected	Observed	Expected
Other DMT	n=579	2	1,21	4	2,35	n=270	4	2,62	5	2,87
	RR (95%CI)	1.65 (0.20-5.94)		1.70 (0.46-4.33)		RR (95%CI)	1.53 (0.42-3.86)		1.74 (0.57-4.02)	
Interferon	n=89	0	0,17	0	0,35	n=43	0	0,22	0	0,34
	RR (95%CI)	0		0		RR (95%CI)	0		0	
Anti-CD20	n=66	0	0,10	1	0,23	n=110	3	0,56	2	0,79
	RR (95%CI)	0		4.39 (0.11-23.4)		RR (95%CI)	5.40 (1.11-15.25)		2.53 (0.31-8.93)	
No therapy	n=75	0	0,19	0	0,35	n=130	13	3,99	10	2,74
	RR (95%CI)	0		0		RR (95%CI)	3.26 (1.77-5.37)		3.65 (1.78-6.50)	

RR=Relative risk, CI=Confidence Interval, DMT= disease modifying therapy, ICU= Intensive Care Unit