

Appendix Table A1. Percentage of corrected refractive errors (n=3,468).

Group	Definition	Total (%)	Male	Female
Myopia	SE \leq -0.50D in at least one of the eyes	12.85 (95% CI: 11.06—14.82)	4.08 (95% CI: 3.69—6.13)	8.04 (95% CI: 6.60—9.67)
Bilateral myopia	SE \leq -0.50D in both eyes	9.54 (95% CI: 7.98—11.29)	3.70 (95% CI: 2.73—4.89)	5.83 (95% CI: 4.60—7.27)
Clinically significant myopia	SE \leq -0.75D in at least one eye	11.51 (95% CI: 9.80—13.40)	4.25 (95% CI: 3.21—5.52)	7.25 (95% CI: 5.88—8.82)
High myopia	SE \leq -5.00D in at least one eye	0.8 (95% CI: 0.43—1.54)	0.47 (95% CI: 0.17—1.02)	0.39 (95% CI: 0.12—0.91)
Hyperopia	SE \geq +2.00D in at least one eye	0.47% (95% CI: 0.17— 1.02)	0.47% (95% CI: 0.17— 1.02)	none
Astigmatism	cylinder \geq 0.75D in at least one eye	9.93% (95% CI: 8.34—11.71)	4.10% (95% CI: 3.07—5.34)	5.83 % (95% CI: 4.60—7.27)
Emmetropia	-0.50 < SE < +0.50D in at least one eye	4.81% (95% CI: 3.69—6.13)	1.81% (95% CI: 1.11—2.70)	2.99 % (95% CI: 2.12—4.09)

D = diopter; CI = Confident Interval; SE = Spherical Equivalent