

Table, Supplemental Digital Content 5: Secondary Outcomes – Void While Standing

	Event Rate (%)	95% CI	<i>p</i>-value	Q-value	<i>I</i>²	N, Studies	N, Patients
Void While Standing	79.0	62.5-89.4	0.001	25.09	4.33	25	729
All Procedures							
Transmen	73.0	45.9-89.6	0.092	20.64	46.71	12	661
Cismen	82.8	70.7-90.6	<0.001	8.14	0	13	68
PU	86.8	61.8-96.4	0.009	8.62	0	15	508
SU	63.2	37.9-82.8	0.307	4.75	0	6	62
RFFF	83.6	62.0-94.1	0.005	16.11	0.68	17	618
Transmen	76.7	39.8-94.2	0.145	13.32	39.96	9	495
Cismen	90.5	79.0-96.0	<0.001	1.03	0	8	51
PU	88.8	58.1-97.8	0.020	6.69	0	12	508
SU	60.4	26.0-86.9	0.573	2.12	5.82	3	50
Other Procedure	67.1	39.5-86.4	0.220	4.39	0	8	111
Transmen	71.1	15.4-97.1	0.498	1.46	0	3	94
Cismen	67.0	41.7-85.2	0.183	2.39	0	5	17
PU	77.0	44.9-93.2	0.094	0.09	36.45	3	26
SU	70.3	28.1-93.5	0.348	1.83	0	3	12

Footnotes:

PU = Primary Urethroplasty, SU = Staged Urethroplasty