

Variable	Number (%)
Patients	40
Mean age $\pm$ SD, yr	55.3 $\pm$ 6.9
Mean BMI $\pm$ SD, kg/m <sup>2</sup>	27.6 $\pm$ 3.7
Stage	
0	9 (22.5%)
1	5 (12.5%)
2	6 (15%)
3	19 (47.5%)
4	1 (2.5%)
Flap utilized	
AICAP	5 (12.5%)
MICAP	5 (12.5%)
LICAP/LTAP	18 (45%)
MSLD	12 (30%)
Nipple pedicle	
Inferior	23 (57.5%)
Medial	10 (25%)
Superior	5 (12.5%)
Lateral	2 (5%)
Breast Ptosis	
Grade 1	11 (27.5%)
Grade 2	26 (65%)
Grade 3	3 (7.5%)
Operative times $\pm$ SD, minutes	115 + 21.7
AICAP, MICAP, LICAP/LTAP	183 $\pm$ 27.5
Immediate MSLD	170 + 21.2
Delayed MSLD	

Tumor size, cm	4.0 ± 2.0
All	2.4 ± 0.59
AICAP	2.1 ± 0.48
MICAP	4.1 ± 1.1
LICAP/LTAP	5.3 ± 1.73
MSLD	
Specimen weight, grams	
All	152 ± 102
AICAP	49 ± 11
MICAP	47 ± 15
LICAP/LTAP	173 ± 60
MSLD	260 ± 80
Follow-up, months	13.6 ± 6.5

**SDC 1. Demographics, operative details, and oncological characteristics of patients undergoing combined Wise-pattern mastopexy and autologous flap reconstruction.**