


SUPPLEMENTAL DIGITAL CONTENT 6

HIV-1 RNA <40 copies/mL at Week 96 (Snapshot analysis, N=272) by subgroup.



#FAA, number of fully-active antiretrovirals according to study entry criteria; GSS, genotypic susceptibility score; PSS, phenotypic susceptibility score; OBT, optimized background therapy; OSS, overall susceptibility score; S-GSS, Stanford GSS. Only subgroups with ≥ 20 participants are shown. ^aResponse rates in other subgroups were American Indian/Alaskan Native: 57% (4/7), Native Hawaiian/Other Pacific Islander: 100% (1/1), Asian: 100% (2/2), and other: 71% (12/17). ^bResponse rate for rest of world was 88% (7/8). ^cResponse rate for #FAA = 0 was 31% (5/16). ^dResponse rates for S-GSS = 0 and missing were 100% (1/1) and 13% (1/8), respectively. ^eResponse rates for GSS = 0 and missing were 0% (0/1) and 11% (1/9), respectively. ^fResponse rate for PSS missing was 20% (2/10). ^gResponse rate for OSS missing was 47% (7/15). ^hResponse rate for OSS-new missing was 50% (7/14). ⁱIncluded for comparison with other OBT susceptibility scores.