

Supplemental Digital Content 1, Table S1. ACTG Staging of KS

	Good Risk (All of the following)	Poor Risk (Any of the following)
T	T_0	T_1
Tumor Extent	No edema, ulceration, extensive oral disease, or visceral disease	Edema or ulceration, extensive oral mucosa KS, or visceral KS
I	I_0	I_1
Immune System	CD4 > 150 cells/mL	CD4 < 150 cells/mL
S	S_0	S_1
Systemic Illness	No opportunistic infections or thrush No "B" symptoms Karnofsky Score \geq 70 %	History of opportunistic infections or thrush "B" symptoms ^a present Karnofsky Score < 70 % other HIV related disease

a. "B" symptoms – unexplained fever; night sweats; >10 % involuntary weight loss or diarrhea persisting > 2 weeks

From: Krown SE, et. al. Kaposi's sarcoma in the acquired immune deficiency syndrome: a proposal for uniform evaluation, response, and staging criteria. AIDS Clinical Trials Group Oncology Committee. J Clin Oncol 1989.¹³

Krown SE, Testa MA, Huang J, et al. AIDS-related Kaposi's sarcoma: prospective validation of the AIDS Clinical Trials Group staging classification. J Clin Oncol 1997; 15:3085-3092.¹⁴