

SUPPLEMENTARY TABLE 1. Patient characteristics

Patient ^a	Sex	Age	Lymphocytes×10 ⁹ /L				Monocytes×10 ⁹ /L				CRP mg/mL			
			Pre-Tx	3 m	6 m	12 m	pre-Tx	3 m	6 m	12 m	Pre-Tx	3 m	6 m	12 m
High IL-10 frequency group														
1	M	54	2.35	1.66	1.64	2.21	0.59	0.50	0.40	0.37	13.5	53.0	24.4	3.5
2	F	24	0.78	1.67	1.74	2.29	0.23	0.46	0.38	0.31	2.1	0.0	0.7	0.0
3	M	47	1.01	1.38	0.82	2.88	0.18	0.22	0.33	0.48	0.0	1.0	3.6	2.0
4	F	35	1.08	0.77	1.27	0.83	0.34	0.14	0.24	0.26	0.0	0.0	0.0	0.0
Low IL-10 frequency group														
5	M	56	1.09	1.20	2.25	1.87	0.25	0.24	0.23	0.18	0.0	0.0	0.0	0.0
6	M	68	0.89	0.35	0.26	0.54	0.17	0.42	0.36	0.36	20.0	12.1	0.5	0.0
7	M	41	0.68	0.81	0.93	0.94	0.26	0.29	0.09	0.22	4.8	9.9	36.3	5.7
8	M	50	1.13	1.47	1.52	ND	0.20	0.28	0.27	ND	0.6	0.0	0.7	ND
9	M	37	1.15	0.91	ND	ND	1.15	0.91	ND	ND	2.9	3.3	ND	ND
10	M	67	1.43	0.99	1.14	1.09	0.35	0.12	0.30	0.28	1.1	5.6	1.9	0.0
11	F	62	1.12	2.07	1.88	ND	0.36	0.54	0.42	ND	2.0	1.1	0.5	ND
12	F	66	1.16	1.32	ND	ND	0.56	0.43	0.55	0.00	16.8	1.5	ND	0.0

^a All patients had a well-functioning graft at 12 months after transplantation (median creatinine level 124 $\mu\text{mol/L}$ [range, 97–180 $\mu\text{mol/L}$]).
ND, not done.