

Supplemental tables

Table S1 - Answers to components of QRDS by hospitalization status

Question	Overall N (111)	Non- Hospitalized N (78)	Hospitalized N (33)	p-value
MEMORY AND RECALL				
Since being diagnosed with COVID-19, have you noticed changes to your memory and recall?				0.12
No obvious memory loss or inconsistent forgetfulness that does not interfere with function in everyday activities	76 (68.5%)	58 (74%)	18 (55%)	
Consistent mild forgetfulness or partial recollection of events that may interfere with performing your daily activities, repeat questions/statements, misplace items, forgets appointments	23 (20.7%)	13 (17%)	10 (30%)	
Mild to moderate memory loss; more noticeable for recent events; interferes with performing everyday activities	9 (8.1%)	6 (8%)	3 (9%)	
Moderate to severe memory loss; only highly learned information remembered; new information rapidly forgotten	3 (2.7%)	1 (1%)	2 (6%)	
Severe memory loss, almost impossible to recall new information, long term memory may be affected	0	0	0	
median (IQR)	0 (0, .5)	0 (0, .5)	0 (0, .5)	0.043
ORIENTATION				
Since being diagnosed with COVID-19, have you noticed changes in your orientation?				0.38
Fully oriented to person, place, and time nearly all the time	82 (73.9%)	59 (76%)	18 (55%)	
Slight difficulty keeping track of time, may forget day or date more frequently than the past	25 (22.5%)	18 (23%)	10 (30%)	
Mild to moderate difficulty in keeping track of time and sequence of events; forgets month or year; oriented to familiar places but gets confused outside familiar areas; gets lost or wanders	2 (1.8%)	1 (1%)	3 (9%)	
Moderate to severe difficulty, usually disoriented to time and place (familiar and unfamiliar); frequently dwells in past	1 (0.9%)	1 (1%)	2 (6%)	
Only oriented to your name, although may recognize other family members	0	0	0	
Did not answer	1 (1%)	0 (0%)	1 (3%)	
median (IQR)	0 (0, .5)	0 (0, 0)	0 (0, .5)	0.59
DECISION MAKING AND PROBLEM-SOLVING ABILITIES				
Since being diagnosed with COVID-19, have you noticed changes in your decision making and problem solving?				0.006
Solves everyday problems without difficulty; handles personal business and financial matters well; decision-making abilities consistent with past performance	73 (65.8%)	58 (74%)	15 (45%)	
Slight impairment or takes longer to solve problems; trouble with abstract concepts, decisions still sound	30 (27.0%)	17 (22%)	13 (39%)	
Moderate difficulty with handling problems and making decisions; defers many decisions to others; social judgment and behavior may be slightly impaired; loss of insight	8 (7.2%)	3 (4%)	5 (15%)	

Severely impaired in handling problems, making only simple personal decisions, social judgment and behavior often impaired, lacks insight	0	0	0	0.002
Unable to make decisions or solve problems, others make nearly all decisions for you	0	0	0	
median (IQR)	0 (0, .5)	0 (0, .5)	.5 (0, .5)	
ACTIVITIES OUTSIDE THE HOME				
Since being diagnosed with COVID-19, have you noticed changes in your ability to do activities outside the home?				0.043
Independent in function at the usual level of performance in profession, shopping, community and religious activities, volunteering, or social groups	83 (74.8%)	63 (81%)	20 (61%)	
Slight impairment in these activities compared to previous performance; slight change in driving skills; still able to handle emergency situations	25 (22.5%)	14 (18%)	11 (33%)	
Unable to function independently but still may attend and be engaged; appears normal to others; notable changes in driving skills; concern about ability to handle emergency situations	2 (1.8%)	1 (1%)	1 (3%)	
No pretense of independent function outside the home; appears well enough to be taken to activities outside the family home but generally needs to be accompanied	1 (0.9%)	0 (0%)	1 (3%)	
No independent function or activities; appear too ill to be taken to activities outside the home	0	0	0	
median (IQR)	0 (0, .5)	0 (0, 0)	0 (0, .5)	0.022
FUNCTION AT HOME AND HOBBY ACTIVITIES				
Since being diagnosed with COVID-19, have you noticed changes in your function at home and hobby activities?				0.027
Chores at home, hobbies and personal interests are well maintained compared with past performance	86 (77.5%)	66 (85%)	20 (61%)	
Slight impairment or less interest in these activities; trouble operating appliances (particularly new purchases)	15 (13.5%)	8 (10%)	7 (21%)	
Mild but definite impairment in home and hobby function; more difficult chores or tasks abandoned; more complicated hobbies and interests given up	7 (6.3%)	3 (4%)	4 (12%)	
Only simple chores preserved, very restricted interest in hobbies which are poorly maintained	3 (2.7%)	1 (1%)	2 (6%)	
No meaningful function in the household chores or with prior hobbies	0	0	0	
median (IQR)	0 (0, 0)	0 (0, 0)	0 (0, .5)	0.005
TOILETING AND PERSONAL HYGIENE				
Since being diagnosed with COVID-19, have you noticed changes in your ability in toileting and personal hygiene?				0.69
Fully capable of self-care (dressing, grooming, washing, bathing, toileting)	103 (92.8%)	73 (94%)	30 (91%)	
Slight changes in abilities and attention to these activities	8 (7.2%)	5 (6%)	3 (9%)	
Needs prompting to complete these activities but may still complete independently	0	0	0	
Requires some assistance in dressing, hygiene, keeping of personal items; occasionally incontinent	0	0	0	
Requires significant help with personal care and hygiene; frequent incontinence	0	0	0	

median (IQR)	0 (0, 0)	0 (0, 0)	0 (0, 0)	0.62
BEHAVIOR AND PERSONALITY CHANGES				
Since being diagnosed with COVID-19, have you noticed changes in your behavior and personality?				0.015
Socially appropriate behavior in public and private; no changes in personality	82 (73.9%)	61 (78%)	21 (64%)	
Questionable or very mild changes in behavior, personality, emotional control, appropriateness of choices	14 (12.6%)	6 (8%)	8 (24%)	
Mild changes in behavior or personality	12 (10.8%)	10 (13%)	2 (6%)	
Moderate behavior or personality changes, affects interactions with others; may be avoided by friends, neighbors, or distant relatives	2 (1.8%)	0 (0%)	2 (6%)	
Severe behavior or personality changes; making interactions with others often unpleasant or avoided	1 (0.9%)	1 (1%)	0 (0%)	
Since being diagnosed with COVID-19, have you noticed changes in your behavior and personality? median (IQR)	0 (0, .5)	0 (0, 0)	0 (0, .5)	0.17
LANGUAGE AND COMMUNICATION ABILITIES				
Since being diagnosed with COVID-19, have you noticed changes to your language and communication abilities?				0.13
No language difficulty or occasional word searching; reads and writes as in the past	72 (64.9%)	54 (69%)	18 (55%)	
Consistent mild word finding difficulties, using descriptive terms or takes longer to get point across, mild problems with comprehension, decreased conversation; may affect reading and writing	32 (28.8%)	21 (27%)	11 (33%)	
Moderate word finding difficulty in speech, cannot name objects, marked reduction in work production; reduced comprehension, conversation, writing, and/or reading	3 (2.7%)	2 (3%)	1 (3%)	
Moderate to severe impairments in speech production or comprehension; has difficulty in communicating thoughts to others; limited ability to read or write	4 (3.6%)	1 (1%)	3 (9%)	
Severe deficits in language and communication; little to no understandable speech is produced	0	0	0	
median (IQR)	0 (0, .5)	0 (0, .5)	0 (0, .5)	0.095
MOOD				
Since being diagnosed with COVID-19, have you noticed changes to your mood?				0.099
No changes in mood, interest, or motivation level	62 (55.9%)	49 (63%)	13 (39%)	
Occasional sadness, depression, anxiety, nervousness, or loss of interest/motivation	33 (29.7%)	20 (26%)	13 (39%)	
Daily mild issues with sadness, depression, anxiety, nervousness, or loss of interest/motivation	10 (9.0%)	6 (8%)	4 (12%)	
Moderate issues with sadness, depression, anxiety, nervousness, or loss of interest/motivation	5 (4.5%)	2 (3%)	3 (9%)	
Severe issues with sadness, depression, anxiety, nervousness, or loss of interest/motivation	0	1 (1%)	0 (0%)	
median (IQR)	0 (0, .5)	0 (0, .5)	.5 (0, .5)	0.023
If you have had changes in mood, were those mood changes present prior to COVID?				0.21
No	50 (45.0%)	36 (46%)	14 (42%)	
Yes	27 (24.3%)	15 (19%)	12 (36%)	
No answer	34 (30.6%)	27 (35%)	7 (21%)	

ATTENTION AND CONCENTRATION

Since being diagnosed with COVID-19, have you noticed changes to your attention and concentration?				0.10
Normal attention, concentration, and interaction with environment and surroundings	66 (59.5%)	50 (64%)	16 (48%)	
Mild problems with attention, concentration, and interaction with environment and surroundings, may appear drowsy during day	35 (31.5%)	23 (29%)	12 (36%)	
Moderate problems with attention and concentration, may have staring spells or spend time with eyes closed, increased daytime sleepiness	8 (7.2%)	3 (4%)	5 (15%)	
Significant portion of the day is spend sleeping, not paying attention to environment, when having a conversation may say things that are illogical or not consistent with topic	2 (1.8%)	2 (3%)	0 (0%)	
Limited to no ability to pay attention to external environment or surroundings	0	0	0	
median (IQR)	0 (0, .5)	0 (0, .5)	.5 (0, .5)	0.098

Table S2- Answers to components of QRDS by time from SARS-CoV2 infection				
Factor	Overall N (111)	<6 months N (76)	>6 months N (35)	p-value
MEMORY AND RECALL				
Since being diagnosed with COVID-19, have you noticed changes to your memory and recall?				0.81
No obvious memory loss or inconsistent forgetfulness that does not interfere with function in everyday activities	76 (68.5%)	53 (70%)	23 (66%)	
Consistent mild forgetfulness or partial recollection of events that may interfere with performing your daily activities, repeat questions/statements, misplace items, forgets appointments	23 (20.7%)	14 (18%)	9 (26%)	
Mild to moderate memory loss; more noticeable for recent events; interferes with performing everyday activities	9 (8.1%)	7 (9%)	2 (6%)	
Moderate to severe memory loss; only highly learned information remembered; new information rapidly forgotten	3 (2.7%)	2 (3%)	1 (3%)	
Severe memory loss, almost impossible to recall new information, long term memory may be affected	0	0	0	
median (IQR)	0 (0, .5)	0 (0, .5)	0 (0, .5)	0.78
ORIENTATION				
Since being diagnosed with COVID-19, have you noticed changes in your ability to				0.41
Fully oriented to person, place, and time nearly all the time	82 (73.9%)	58 (76%)	24 (69%)	
Slight difficulty keeping track of time, may forget day or date more frequently than the past	25 (22.5%)	17 (22%)	8 (23%)	
Mild to moderate difficulty in keeping track of time and sequence of events; forgets month or year; oriented to familiar places but gets confused outside familiar areas; gets lost or wanders	2 (1.8%)	1 (1%)	1 (3%)	
Moderate to severe difficulty, usually disoriented to time and place (familiar and unfamiliar); frequently dwells in past	1 (0.9%)	0 (0%)	1 (3%)	
Only oriented to your name, although may recognize other family members	0	0	0	
Did not answer	1 (0.9%)	0 (0%)	1 (3%)	
median (IQR)	0 (0, .5)	0 (0, 0)	0 (0, .5)	0.46
DECISION MAKING AND PROBLEM-SOLVING ABILITIES				
Since being diagnosed with COVID-19, have you noticed changes in your decision making and problem solving?				0.30
Solves everyday problems without difficulty; handles personal business and financial matters well; decision-making abilities consistent with past performance	73 (65.8%)	53 (70%)	20 (57%)	
Slight impairment or takes longer to solve problems; trouble with abstract concepts, decisions still sound	30 (27.0%)	19 (25%)	11 (31%)	
Moderate difficulty with handling problems and making decisions; defers many decisions to others; social	8 (7.2%)	4 (5%)	4 (11%)	

judgment and behavior may be slightly impaired; loss of insight				0.16
Severely impaired in handling problems, making only simple personal decisions, social judgment and behavior often impaired, lacks insight	0	0	0	
Unable to make decisions or solve problems, others make nearly all decisions for you	0	0	0	
median (IQR)	0 (0, .5)	0 (0, .5)	0 (0, .5)	
ACTIVITIES OUTSIDE THE HOME				
Since being diagnosed with COVID-19, have you noticed changes in your ability to do activities outside the home?				0.099
Independent in function at the usual level of performance in profession, shopping, community and religious activities, volunteering, or social groups	83 (74.8%)	61 (80%)	22 (63%)	
Slight impairment in these activities compared to previous performance; slight change in driving skills; still able to handle emergency situations	25 (22.5%)	14 (18%)	11 (31%)	
Unable to function independently but still may attend and be engaged; appears normal to others; notable changes in driving skills; concern about ability to handle emergency situations	2 (1.8%)	1 (1%)	1 (3%)	
No pretense of independent function outside the home; appears well enough to be taken to activities outside the family home but generally needs to be accompanied	1 (0.9%)	0 (0%)	1 (3%)	
No independent function or activities; appear too ill to be taken to activities outside the home	0	0	0	
median (IQR)	0 (0, .5)	0 (0, 0)	0 (0, .5)	0.043
FUNCTION AT HOME AND HOBBY ACTIVITIES				
Since being diagnosed with COVID-19, have you noticed changes in your ability to				0.11
Chores at home, hobbies and personal interests are well maintained compared with past performance	86 (77.5%)	60 (79%)	26 (74%)	
Slight impairment or less interest in these activities; trouble operating appliances (particularly new purchases)	15 (13.5%)	11 (14%)	4 (11%)	
Mild but definite impairment in home and hobby function; more difficult chores or tasks abandoned; more complicated hobbies and interests given up	7 (6.3%)	2 (3%)	5 (14%)	
Only simple chores preserved, very restricted interest in hobbies which are poorly maintained	3 (2.7%)	3 (4%)	0 (0%)	
No meaningful function in the household chores or with prior hobbies	0	0	0	
median (IQR)	0 (0, 0)	0 (0, 0)	0 (0, .5)	0.53
TOILETING AND PERSONAL HYGIENE				
Since being diagnosed with COVID-19, have you noticed changes in your ability in toileting and personal hygiene?				0.26
Fully capable of self-care (dressing, grooming, washing, bathing, toileting)	103 (92.8%)	72 (95%)	31 (89%)	
Slight changes in abilities and attention to these activities	8 (7.2%)	4 (5%)	4 (11%)	
Needs prompting to complete these activities but may still complete independently	0	0	0	

Requires some assistance in dressing, hygiene, keeping of personal items; occasionally incontinent	0	0	0	0.25
Requires significant help with personal care and hygiene; frequent incontinence	0	0	0	
median (IQR)	0 (0, 0)	0 (0, 0)	0 (0, 0)	
BEHAVIOR AND PERSONALITY CHANGES				
Since being diagnosed with COVID-19, have you noticed changes in your behavior and personality?				0.68
Socially appropriate behavior in public and private; no changes in personality	82 (73.9%)	54 (71%)	28 (80%)	
Questionable or very mild changes in behavior, personality, emotional control, appropriateness of choices	14 (12.6%)	10 (13%)	4 (11%)	
Mild changes in behavior or personality	12 (10.8%)	10 (13%)	2 (6%)	
Moderate behavior or personality changes, affects interactions with others; may be avoided by friends, neighbors, or distant relatives	2 (1.8%)	1 (1%)	1 (3%)	
Severe behavior or personality changes; making interactions with others often unpleasant or avoided	1 (0.9%)	1 (1%)	0 (0%)	
median (IQR)	0 (0, .5)	0 (0, .5)	0 (0, 0)	0.30
LANGUAGE AND COMMUNICATION ABILITIES				
Since being diagnosed with COVID-19, have you noticed changes to your language and communication abilities?				0.97
No language difficulty or occasional word searching; reads and writes as in the past	72 (64.9%)	50 (66%)	22 (63%)	
Consistent mild word finding difficulties, using descriptive terms or takes longer to get point across, mild problems with comprehension, decreased conversation; may affect reading and writing	32 (28.8%)	21 (28%)	11 (31%)	
Moderate word finding difficulty in speech, cannot name objects, marked reduction in work production; reduced comprehension, conversation, writing, and/or reading	3 (2.7%)	2 (3%)	1 (3%)	
Moderate to severe impairments in speech production or comprehension; has difficulty in communicating thoughts to others; limited ability to read or write	4 (3.6%)	3 (4%)	1 (3%)	
Severe deficits in language and communication; little to no understandable speech is produced	0	0	0	
median (IQR)	0 (0, .5)	0 (0, .5)	0 (0, .5)	0.81
MOOD				
Since being diagnosed with COVID-19, have you noticed changes to your mood?				0.13
No changes in mood, interest, or motivation level	62 (55.9%)	48 (63%)	14 (40%)	
Occasional sadness, depression, anxiety, nervousness, or loss of interest/motivation	33 (29.7%)	18 (24%)	15 (43%)	
Daily mild issues with sadness, depression, anxiety, nervousness, or loss of interest/motivation	10 (9.0%)	6 (8%)	4 (11%)	
Moderate issues with sadness, depression, anxiety, nervousness, or loss of interest/motivation	5 (4.5%)	3 (4%)	2 (6%)	
Severe issues with sadness, depression, anxiety, nervousness, or loss of interest/motivation	1 (0.9%)	1 (1%)	0 (0%)	

Since being diagnosed with COVID-19, have you noticed changes to your mood? median (IQR)	0 (0, .5)	0 (0, .5)	0.5 (0, .5)	0.044
If you have had changes in mood, were those mood changes present prior to COVID?				0.22
No	50 (45.0%)	34 (45%)	16 (46%)	
Yes	27 (24.3%)	14 (18%)	13 (37%)	
No answer	34 (30.6%)	28 (37%)	6 (17%)	
ATTENTION AND CONCENTRATION				
Since being diagnosed with COVID-19, have you noticed changes to your attention and concentration?				0.52
Normal attention, concentration, and interaction with environment and surroundings	66 (59.5%)	46 (61%)	20 (57%)	
Mild problems with attention, concentration, and interaction with environment and surroundings, may appear drowsy during day	35 (31.5%)	25 (33%)	10 (29%)	
Moderate problems with attention and concentration, may have staring spells or spend time with eyes closed, increased daytime sleepiness	8 (7.2%)	4 (5%)	4 (11%)	
Significant portion of the day is spend sleeping, not paying attention to environment, when having a conversation may say things that are illogical or not consistent with topic	2 (1.8%)	1 (1%)	1 (3%)	
Limited to no ability to pay attention to external environment or surroundings	0	0	0	
median (IQR)	0 (0, .5)	0 (0, .5)	0 (0, .5)	0.55

Table S3 - Answers to components of PHQ-9 by hospitalization status				
Question	Overall N (111)	Non-Hospitalized N (76)	Hospitalized N (32)	p-value
1. Little interest or pleasure in doing things				0.18
Not at all	68 (61.3%)	48 (62%)	20 (61%)	
Several Days	29 (26.1%)	23 (29%)	6 (18%)	
More than half the days	5 (4.5%)	2 (3%)	3 (9%)	
Nearly every day	6 (5.4%)	3 (4%)	3 (9%)	
No answer	3 (2.7%)	2 (3%)	1 (3%)	
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	0.63
2. Feeling down, depressed, or hopeless				0.14
Not at all	65 (58.6%)	48 (62%)	17 (52%)	
Several Days	35 (31.5%)	26 (33%)	9 (27%)	
More than half the days	6 (5.4%)	2 (3%)	4 (12%)	
Nearly every day	2 (1.8%)	1 (1%)	1 (3%)	
No answer	3 (2.7%)	1 (1%)	2 (6%)	
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	0.27
3. Trouble falling or staying asleep, or sleeping too much				0.18
Not at all	49 (44.1%)	39 (50%)	10 (30%)	
Several Days	26 (23.4%)	17 (22%)	9 (27%)	
More than half the days	20 (18.0%)	13 (17%)	7 (21%)	
Nearly every day	15 (13.5%)	8 (10%)	7 (21%)	
No answer	1 (0.9%)	1 (1%)	0 (0%)	
median (IQR)	1 (0, 2)	0 (0, 2)	1 (0, 2)	0.038
4. Feeling tired or having little energy				0.30
Not at all	40 (36.0%)	30 (38%)	10 (30%)	
Several Days	34 (30.6%)	26 (33%)	8 (24%)	
More than half the days	23 (20.7%)	14 (18%)	9 (27%)	
Nearly every day	13 (11.7%)	7 (9%)	6 (18%)	
No answer	1 (0.9%)	1 (1%)	0 (0%)	
median (IQR)	1 (0, 2)	1 (0, 2)	1 (0, 2)	0.11
5. Poor appetite or over eating				0.067
Not at all	72 (64.9%)	56 (72%)	16 (48%)	
Several Days	19 (17.1%)	12 (15%)	7 (21%)	
More than half the days	15 (13.5%)	7 (9%)	8 (24%)	
Nearly every day	5 (4.5%)	3 (4%)	2 (6%)	
median (IQR)	0 (0, 1)	0 (0, 1)	1 (0, 2)	0.015
6. Feeling bad about yourself-or that you are a failure or have let yourself or your family down				0.055
Not at all	84 (75.7%)	64 (82%)	20 (61%)	
Several Days	22 (19.8%)	12 (15%)	10 (30%)	

More than half the days	3 (2.7%)	1 (1%)	2 (6%)	0.014
Nearly every day	2 (1.8%)	1 (1%)	1 (3%)	
median (IQR)	0 (0, 0)	0 (0, 0)	0 (0, 1)	
7. Trouble concentrating on things, such as reading the newspaper or watching TV				0.54
Not at all	65 (58.6%)	47 (60%)	18 (55%)	
Several Days	35 (31.5%)	22 (28%)	13 (39%)	
More than half the days	8 (7.2%)	7 (9%)	1 (3%)	
Nearly every day	3 (2.7%)	2 (3%)	1 (3%)	
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	0.77
8. Moving or speaking so slowly that other people could have noticed? Or the opposite—being so fidgety or restless that you have been moving around a lot more than usual				0.70
Not at all	97 (87.4%)	69 (88%)	28 (85%)	
Several Days	10 (9.0%)	7 (9%)	3 (9%)	
More than half the days	4 (3.6%)	2 (3%)	2 (6%)	
median (IQR)	0 (0, 0)	0 (0, 0)	0 (0, 0)	0.57
9. Thoughts that you would be better off dead or hurting yourself in some way				0.14
Not at all	103 (92.8%)	75 (96%)	28 (85%)	
Several Days	6 (5.4%)	3 (4%)	3 (9%)	
More than half the days	1 (0.9%)	0 (0%)	1 (3%)	
No answer	1 (0.9%)	0 (0%)	1 (3%)	
median (IQR)	0 (0, 0)	0 (0, 0)	0 (0, 0)	0.088
If you checked off any problems, how difficult have these problems made it for you to do your work, take care of things at home, or get along with other people?				0.36
Not difficult at all	58 (52.3%)	41 (53%)	17 (52%)	
Somewhat difficult	35 (31.5%)	23 (29%)	12 (36%)	
Very difficult	3 (2.7%)	1 (1%)	2 (6%)	
No answer	15 (13.5%)	13 (17%)	2 (6%)	

Table S4- Responses to components of PHQ-9 by time from SARS-CoV2 infection diagnosis

Question	Overall N (111)	<6-months N (76)	>6-months N (35)	p- value
1. Little interest or pleasure in doing things				
Not at all	68 (61.3%)	48 (63%)	20 (57%)	0.90
Several Days	29 (26.1%)	19 (25%)	10 (29%)	
More than half the days	5 (4.5%)	3 (4%)	2 (6%)	
Nearly every day	6 (5.4%)	4 (5%)	2 (6%)	
No answer	3 (2.7%)	2 (3%)	1 (3%)	
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	0.55
2. Feeling down, depressed, or hopeless				
Not at all	65 (58.6%)	47 (62%)	18 (51%)	0.82
Several Days	35 (31.5%)	24 (32%)	11 (31%)	
More than half the days	6 (5.4%)	4 (5%)	2 (6%)	
Nearly every day	2 (1.8%)	1 (1%)	1 (3%)	
No answer	3 (2.7%)	0 (0%)	3 (9%)	
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	0.55
3. Trouble falling or staying asleep, or sleeping too much				
Not at all	49 (44.1%)	35 (46%)	14 (40%)	0.85
Several Days	26 (23.4%)	18 (24%)	8 (23%)	
More than half the days	20 (18.0%)	13 (17%)	7 (20%)	
Nearly every day	15 (13.5%)	9 (12%)	6 (17%)	
No answer	1 (0.9%)	1 (1%)	0 (0%)	
median (IQR)	1 (0, 2)	1 (0, 2)	1 (0, 2)	0.40
4. Feeling tired or having little energy				
Not at all	40 (36.0%)	27 (36%)	13 (37%)	0.84
Several Days	34 (30.6%)	25 (33%)	9 (26%)	
More than half the days	23 (20.7%)	15 (20%)	8 (23%)	
Nearly every day	13 (11.7%)	8 (11%)	5 (14%)	
No answer	1 (0.9%)	1 (1%)	0 (0%)	
median (IQR)	1 (0, 2)	1 (0, 2)	1 (0, 2)	0.74
5. Poor appetite or over eating				
Not at all	72 (64.9%)	54 (71%)	18 (51%)	0.054
Several Days	19 (17.1%)	12 (16%)	7 (20%)	
More than half the days	15 (13.5%)	6 (8%)	9 (26%)	
Nearly every day	5 (4.5%)	4 (5%)	1 (3%)	
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 2)	0.043
6. Feeling bad about yourself-or that you are a failure or have let yourself or your family down				
Not at all	84 (75.7%)	59 (78%)	25 (71%)	0.46
Several Days	22 (19.8%)	15 (20%)	7 (20%)	
More than half the days	3 (2.7%)	1 (1%)	2 (6%)	

Nearly every day	2 (1.8%)	1 (1%)	1 (3%)	
median (IQR)	0 (0, 0)	0 (0, 0)	0 (0, 1)	0.41
7. Trouble concentrating on things, such as reading the newspaper or watching TV				
Not at all	65 (58.6%)	45 (59%)	20 (57%)	0.98
Several Days	35 (31.5%)	23 (30%)	12 (34%)	
More than half the days	8 (7.2%)	6 (8%)	2 (6%)	
Nearly every day	3 (2.7%)	2 (3%)	1 (3%)	
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	0.91
8. Moving or speaking so slowly that other people could have noticed? Or the opposite—being so fidgety or restless that you have been moving around a lot more than usual				
Not at all	97 (87.4%)	68 (89%)	29 (83%)	0.39
Several Days	10 (9.0%)	5 (7%)	5 (14%)	
More than half the days	4 (3.6%)	3 (4%)	1 (3%)	
median (IQR)	0 (0, 0)	0 (0, 0)	0 (0, 0)	0.36
9. Thoughts that you would be better off dead or hurting yourself in some way				
Not at all	103 (92.8%)	71 (93%)	32 (91%)	0.44
Several Days	6 (5.4%)	4 (5%)	2 (6%)	
More than half the days	1 (0.9%)	0 (0%)	1 (3%)	
No answer	1 (0.9%)	1 (1%)	0 (0%)	
median (IQR)	0 (0, 0)	0 (0, 0)	0 (0, 0)	0.50
If you checked off any problems, how difficult have these problems made it for you to do your work, take care of things at home, or get along with other people?				
Not difficult at all	58 (52.3%)	43 (57%)	15 (43%)	0.32
Somewhat difficult	35 (31.5%)	21 (28%)	14 (40%)	
Very difficult	3 (2.7%)	2 (3%)	1 (3%)	
No answer	15 (13.5%)	10 (13%)	5 (14%)	

Table S5- Responses to components of GAD-7 by hospitalization status				
Question	Overall N (111)	Non-Hospitalized N (78)	Hospitalized N (33)	p-value
Over the last 2 weeks, how often have you been bothered by feeling nervous, anxious, or on edge?				
Not at all	68 (61.3%)	50 (64%)	18 (55%)	0.13
Several days	36 (32.4%)	24 (31%)	12 (36%)	
More than half of the days	4 (3.6%)	1 (1%)	3 (9%)	
Nearly everyday	3 (2.7%)	3 (4%)	0 (0%)	
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	0.33
Over the last 2 weeks, how often have you been bothered by not being able to stop, or control, worrying?				
Not at all	77 (69.4%)	56 (72%)	21 (64%)	0.73
Several days	26 (23.4%)	17 (22%)	9 (27%)	
More than half of the days	5 (4.5%)	3 (4%)	2 (6%)	
Nearly everyday	3 (2.7%)	2 (3%)	1 (3%)	
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	0.39
Over the last 2 weeks, how often have you been bothered by worrying too much about different things?				
Not at all	73 (65.8%)	51 (65%)	22 (67%)	0.44
Several days	29 (26.1%)	22 (28%)	7 (21%)	
More than half of the days	6 (5.4%)	3 (4%)	3 (9%)	
Nearly everyday	2 (1.8%)	1 (1%)	1 (3%)	
No answer	1 (0.9%)	1 (1%)	0 (0%)	
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	0.85
Over the last 2 weeks, how often have you had trouble relaxing?				
Not at all	70 (63.1%)	50 (64%)	20 (61%)	0.59
Several days	30 (27.0%)	22 (28%)	8 (24%)	
More than half of the days	8 (7.2%)	4 (5%)	4 (12%)	
Nearly everyday	3 (2.7%)	2 (3%)	1 (3%)	
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	0.58
Over the last 2 weeks, how often have you been bothered by being so restless that it's hard to sit still?				
Not at all	95 (85.6%)	66 (85%)	29 (88%)	0.19
Several days	10 (9.0%)	9 (12%)	1 (3%)	
More than half of the days	3 (2.7%)	1 (1%)	2 (6%)	
Nearly everyday	2 (1.8%)	2 (3%)	0 (0%)	
No answer	1 (0.9%)	0 (0%)	1 (3%)	
median (IQR)	0 (0, 0)	0 (0, 0)	0 (0, 0)	0.44
Over the last 2 weeks, how often have you been bothered by becoming easily annoyed or irritable?				

Not at all	63 (56.8%)	47 (60%)	16 (48%)	0.17
Several days	39 (35.1%)	27 (35%)	12 (36%)	
More than half of the days	6 (5.4%)	2 (3%)	4 (12%)	
Nearly everyday	1 (0.9%)	1 (1%)	0 (0%)	
No answer	2 (1.8%)	1 (1%)	1 (3%)	
median (IQR)	0 (0, 1)	0 (0, 1)	.5 (0, 1)	0.19
Over the last 2 weeks, how often have you been bothered by feeling afraid as if something awful might happen?				
Not at all	82 (73.9%)	61 (78%)	21 (64%)	0.099
Several days	26 (23.4%)	16 (21%)	10 (30%)	
More than half of the days	1 (0.9%)	0 (0%)	1 (3%)	
No answer	2 (1.8%)	1 (1%)	1 (3%)	
median (IQR)	0 (0, 0)	0 (0, 0)	0 (0, 1)	0.12

Table S6- Responses to components of GAD-7 by time from SARS-CoV2 infection diagnosis

Question	Overall N (111)	<6-months N (76)	>6-months N (35)	p- value
Over the last 2 weeks, how often have you been bothered by feeling nervous, anxious, or on edge?				
Not at all	68 (61.3%)	47 (62%)	21 (60%)	0.27
Several days	36 (32.4%)	26 (34%)	10 (29%)	
More than half of the days	4 (3.6%)	1 (1%)	3 (9%)	
Nearly everyday	3 (2.7%)	2 (3%)	1 (3%)	
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	
Over the last 2 weeks, how often have you been bothered by not being able to stop, or control, worrying?				
Not at all	77 (69.4%)	55 (72%)	22 (63%)	0.48
Several days	26 (23.4%)	17 (22%)	9 (26%)	
More than half of the days	5 (4.5%)	2 (3%)	3 (9%)	
Nearly everyday	3 (2.7%)	2 (3%)	1 (3%)	
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	
Over the last 2 weeks, how often have you been bothered by worrying too much about different things?				
Not at all	73 (65.8%)	49 (64%)	24 (69%)	0.46
Several days	29 (26.1%)	22 (29%)	7 (20%)	
More than half of the days	6 (5.4%)	3 (4%)	3 (9%)	
Nearly everyday	2 (1.8%)	1 (1%)	1 (3%)	
No answer	1 (0.9%)	1 (1%)	0 (0%)	
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	0.92
Over the last 2 weeks, how often have you had trouble relaxing?				
Not at all	70 (63.1%)	51 (67%)	19 (54%)	0.42
Several days	30 (27.0%)	19 (25%)	11 (31%)	
More than half of the days	8 (7.2%)	4 (5%)	4 (11%)	
Nearly everyday	3 (2.7%)	2 (3%)	1 (3%)	
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	
Over the last 2 weeks, how often have you been bothered by being so restless that it's hard to sit still?				
Not at all	95 (85.6%)	66 (87%)	29 (83%)	0.50
Several days	10 (9.0%)	7 (9%)	3 (9%)	
More than half of the days	3 (2.7%)	1 (1%)	2 (6%)	
Nearly everyday	2 (1.8%)	2 (3%)	0 (0%)	
No answer	1 (0.9%)	0 (0%)	1 (3%)	
median (IQR)	0 (0, 0)	0 (0, 0)	0 (0, 0)	0.82
Over the last 2 weeks, how often have you been bothered by becoming easily annoyed or irritable?				
Not at all	63 (56.8%)	40 (53%)	23 (66%)	0.63
Several days	39 (35.1%)	29 (38%)	10 (29%)	
More than half of the days	6 (5.4%)	4 (5%)	2 (6%)	
Nearly everyday	1 (0.9%)	1 (1%)	0 (0%)	
No answer	2 (1.8%)	2 (3%)	0 (0%)	

median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	0.27
Over the last 2 weeks, how often have you been bothered by feeling afraid as if something awful might happen?				
Not at all	82 (73.9%)	58 (76%)	24 (69%)	0.25
Several days	26 (23.4%)	16 (21%)	10 (29%)	
More than half of the days	1 (0.9%)	0 (0%)	1 (3%)	
No answer	2 (1.8%)	2 (3%)	0 (0%)	
median (IQR)	0 (0, 0)	0 (0, 0)	0 (0, 1)	0.24

Table S7- Responses to components of IES-6 by hospitalization status				
Question	Overall N (111)	Not- Hospitalized N (78)	Hospitalized N (33)	p-value
After reading each item, indicate how distressing each difficulty has been for you during the <u>past 7 days</u> with respect to <u>your COVID19 diagnosis</u> . How much were you distressed or bothered by these difficulties? In the questions, the word “it” refers to <u>your COVID19 infection</u> .				
1. I thought about it when I did not mean to				
Not at all	65 (58.6%)	48 (62%)	17 (52%)	0.47
A little bit	34 (30.6%)	22 (28%)	12 (36%)	
Moderately	5 (4.5%)	5 (6%)	0 (0%)	
Quite a bit	4 (3.6%)	3 (4%)	1 (3%)	
No answer	3 (2.7%)	0 (0%)	3 (9%)	
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	0.86
2. I felt watchful or on guard.				
Not at all	63 (56.8%)	46 (59%)	17 (52%)	0.55
A little bit	29 (26.1%)	18 (23%)	11 (33%)	
Moderately	9 (8.1%)	8 (10%)	1 (3%)	
Quite a bit	6 (5.4%)	5 (6%)	1 (3%)	
Extremely	1 (0.9%)	1 (1%)	0 (0%)	
No answer	3 (2.7%)	0 (0%)	3 (9%)	0.81
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	
3. Other things kept making me think about it.				
Not at all	64 (57.7%)	46 (59%)	18 (55%)	0.51
A little bit	28 (25.2%)	21 (27%)	7 (21%)	
Moderately	11 (9.9%)	6 (8%)	5 (15%)	
Quite a bit	4 (3.6%)	4 (5%)	0 (0%)	
Extremely	1 (0.9%)	1 (1%)	0 (0%)	
No answer	3 (2.7%)	0 (0%)	3 (9%)	0.93
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	
4. I was aware that I still had a lot of feelings about it, but I didn't deal with them				
Not at all	81 (73.0%)	59 (76%)	22 (67%)	0.37
A little bit	19 (17.1%)	15 (19%)	4 (12%)	
Moderately	4 (3.6%)	2 (3%)	2 (6%)	
Quite a bit	4 (3.6%)	2 (3%)	2 (6%)	
No answer	3 (2.7%)	0 (0%)	3 (9%)	
median (IQR)	0 (0, .5)	0 (0, 0)	0 (0, 1)	0.66
5. I tried not to think about it.				
Not at all	66 (59.5%)	48 (62%)	18 (55%)	0.84
A little bit	29 (26.1%)	21 (27%)	8 (24%)	
Moderately	8 (7.2%)	6 (8%)	2 (6%)	
Quite a bit	2 (1.8%)	1 (1%)	1 (3%)	

Extremely	2 (1.8%)	1 (1%)	1 (3%)	
No answer	4 (3.6%)	1 (1%)	3 (9%)	
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	0.74
6. I had trouble concentrating.				
Not at all	71 (64.0%)	55 (71%)	16 (48%)	
A little bit	23 (20.7%)	14 (18%)	9 (27%)	
Moderately	8 (7.2%)	5 (6%)	3 (9%)	
Quite a bit	3 (2.7%)	2 (3%)	1 (3%)	
Extremely	3 (2.7%)	2 (3%)	1 (3%)	
No answer	3 (2.7%)	0 (0%)	3 (9%)	0.41
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	0.11

Table S8- Responses to components of IES-6 by time from SARS-CoV2 infection diagnosis

Question	Overall N (111)	<6 month N (76)	>6 months N (35)	p-value
After reading each item, indicate how distressing each difficulty has been for you during the past 7 days with respect to <u>your COVID19 diagnosis</u> . How much were you distressed or bothered by these difficulties? In the questions, the word “it” refers to <u>your COVID19 infection</u> .				
1. I thought about it when I did not mean to.				
Not at all	65 (58.6%)	45 (59%)	20 (57%)	1.00
A little bit	34 (30.6%)	23 (30%)	11 (31%)	
Moderately	5 (4.5%)	4 (5%)	1 (3%)	
Quite a bit	4 (3.6%)	3 (4%)	1 (3%)	
No answer	3 (2.7%)	1 (1%)	2 (6%)	
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	0.86
2. I felt watchful or on guard.				
Not at all	63 (56.8%)	45 (59%)	18 (51%)	0.85
A little bit	29 (26.1%)	18 (24%)	11 (31%)	
Moderately	9 (8.1%)	7 (9%)	2 (6%)	
Quite a bit	6 (5.4%)	4 (5%)	2 (6%)	
Extremely	1 (0.9%)	1 (1%)	0 (0%)	
No answer	3 (2.7%)	1 (1%)	2 (6%)	0.77
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	
3. Other things kept making me think about it.				
Not at all	64 (57.7%)	43 (57%)	21 (60%)	0.59
A little bit	28 (25.2%)	22 (29%)	6 (17%)	
Moderately	11 (9.9%)	6 (8%)	5 (14%)	
Quite a bit	4 (3.6%)	3 (4%)	1 (3%)	
Extremely	1 (0.9%)	1 (1%)	0 (0%)	
No answer	3 (2.7%)	1 (1%)	2 (6%)	0.71
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	
4. I was aware that I still had a lot of feelings about it, but I didn’t deal with them				
Not at all	81 (73.0%)	57 (75%)	24 (69%)	0.28
A little bit	19 (17.1%)	14 (18%)	5 (14%)	
Moderately	4 (3.6%)	1 (1%)	3 (9%)	
Quite a bit	4 (3.6%)	3 (4%)	1 (3%)	
No answer	3 (2.7%)	1 (1%)	2 (6%)	
median (IQR)	0 (0, .5)	0 (0, 0)	0 (0, 1)	0.63
5. I tried not to think about it.				
Not at all	66 (59.5%)	46 (61%)	20 (57%)	0.34
A little bit	29 (26.1%)	21 (28%)	8 (23%)	
Moderately	8 (7.2%)	5 (7%)	3 (9%)	
Quite a bit	2 (1.8%)	2 (3%)	0 (0%)	

Extremely	2 (1.8%)	0 (0%)	2 (6%)	
No answer	4 (3.6%)	2 (3%)	2 (6%)	
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	0.71
6. I had trouble concentrating.				
Not at all	71 (64.0%)	51 (67%)	20 (57%)	
A little bit	23 (20.7%)	13 (17%)	10 (29%)	
Moderately	8 (7.2%)	6 (8%)	2 (6%)	
Quite a bit	3 (2.7%)	3 (4%)	0 (0%)	
Extremely	3 (2.7%)	2 (3%)	1 (3%)	
No answer	3 (2.7%)	1 (1%)	2 (6%)	0.51
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	0.66

Table S9- Responses to components of EQ5D-5L by hospitalization status				
Question	All N (111)	Not- Hospitalized N (78)	Hospitalized N (33)	p-value
MOBILITY				
I have no problems in walking about	78 (70.3%)	57 (73%)	21 (64%)	0.47
I have slight problems in walking about	21 (18.9%)	15 (19%)	6 (18%)	
I have moderate problems in walking about	9 (8.1%)	6 (8%)	3 (9%)	
I have severe problems in walking about	1 (0.9%)	0 (0%)	1 (3%)	
No answer	2 (1.8%)	0 (0%)	2 (6%)	
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	0.50
SELF-CARE				
I have no problems washing or dressing myself	101 (91.0%)	72 (92%)	29 (88%)	0.58
I have slight problems washing or dressing myself	6 (5.4%)	5 (6%)	1 (3%)	
I have moderate problems washing or dressing myself	1 (0.9%)	1 (1%)	0 (0%)	
I have severe problems washing or dressing myself	1 (0.9%)	0 (0%)	1 (3%)	
No answer	2 (1.8%)	0 (0%)	2 (6%)	
median (IQR)	0 (0, 0)	0 (0, 0)	0 (0, 0)	0.85
USUAL ACTIVITIES (eg, work, study, housework, family or leisure activities)				
I have no problems doing my usual activities	68 (61.3%)	50 (64%)	18 (55%)	0.26
I have slight problems doing my usual activities	32 (28.8%)	24 (31%)	8 (24%)	
I have moderate problems doing my usual activities	5 (4.5%)	2 (3%)	3 (9%)	
I have severe problems doing my usual activities	4 (3.6%)	2 (3%)	2 (6%)	
No answer	2 (1.8%)	0 (0%)	2 (6%)	
median (IQR)	0 (0, 1)	0 (0, 1)	0 (0, 1)	0.36
PAIN / DISCOMFORT				
I have no pain or discomfort	60 (54.1%)	51 (65%)	9 (27%)	0.003
I have slight pain or discomfort	28 (25.2%)	16 (21%)	12 (36%)	
I have moderate pain or discomfort	18 (16.2%)	10 (13%)	8 (24%)	
I have severe pain or discomfort	3 (2.7%)	1 (1%)	2 (6%)	
No answer	2 (1.8%)	0 (0%)	2 (6%)	
median (IQR)	0 (0, 1)	0 (0, 1)	1 (0, 2)	<0.001
ANXIETY / DEPRESSION:				
I am not anxious or depressed	56 (50.5%)	44 (56%)	12 (36%)	0.15
I am slightly anxious or depressed	39 (35.1%)	27 (35%)	12 (36%)	
I am moderately anxious or depressed	12 (10.8%)	6 (8%)	6 (18%)	
I am severely anxious or depressed	2 (1.8%)	1 (1%)	1 (3%)	
No answer	2 (1.8%)	0 (0%)	2 (6%)	
median (IQR)	0 (0, 1)	0 (0, 1)	1 (0, 1)	0.048

Table S10- Responses to components of EQ5D-5L by time from SARS-CoV2 infection				
Question	All N (111)	<6-months N (76)	>6-months N (35)	p- value
MOBILITY				
I have no problems in walking about	78 (70.3%)	58 (76%)	20 (57%)	0.13
I have slight problems in walking about	21 (18.9%)	12 (16%)	9 (26%)	
I have moderate problems in walking about	9 (8.1%)	5 (7%)	4 (11%)	
I have severe problems in walking about	1 (0.9%)	0 (0%)	1 (3%)	
No answer	2 (1.8%)	1 (1%)	1 (3%)	
median (IQR)	0 (0, 1)	0 (0, 0)	0 (0, 1)	0.042
SELF-CARE				
I have no problems washing or dressing myself	101 (91.0%)	71 (93%)	30 (86%)	0.29
I have slight problems washing or dressing myself	6 (5.4%)	3 (4%)	3 (9%)	
I have moderate problems washing or dressing myself	1 (0.9%)	1 (1%)	0 (0%)	
I have severe problems washing or dressing myself	1 (0.9%)	0 (0%)	1 (3%)	
No answer	2 (1.8%)	1 (1%)	1 (3%)	
median (IQR)	0 (0, 0)	0 (0, 0)	0 (0, 0)	0.23
USUAL ACTIVITIES (eg, work, study, housework, family or leisure activities)				
I have no problems doing my usual activities	68 (61.3%)	51 (67%)	17 (49%)	0.16
I have slight problems doing my usual activities	32 (28.8%)	20 (26%)	12 (34%)	
I have moderate problems doing my usual activities	5 (4.5%)	2 (3%)	3 (9%)	
I have severe problems doing my usual activities	4 (3.6%)	2 (3%)	2 (6%)	
No answer	2 (1.8%)	1 (1%)	1 (3%)	
median (IQR)	0 (0, 1)	0 (0, 1)	.5 (0, 1)	0.052
PAIN / DISCOMFORT				
I have no pain or discomfort	60 (54.1%)	45 (59%)	15 (43%)	0.11
I have slight pain or discomfort	28 (25.2%)	20 (26%)	8 (23%)	
I have moderate pain or discomfort	18 (16.2%)	8 (11%)	10 (29%)	
I have severe pain or discomfort	3 (2.7%)	2 (3%)	1 (3%)	
No answer	2 (1.8%)	1 (1%)	1 (3%)	
median (IQR)	0 (0, 1)	0 (0, 1)	1 (0, 2)	0.053
ANXIETY / DEPRESSION:				
I am not anxious or depressed	56 (50.5%)	41 (54%)	15 (43%)	0.48
I am slightly anxious or depressed	39 (35.1%)	24 (32%)	15 (43%)	
I am moderately anxious or depressed	12 (10.8%)	9 (12%)	3 (9%)	
I am severely anxious or depressed	2 (1.8%)	1 (1%)	1 (3%)	
No answer	2 (1.8%)	1 (1%)	1 (3%)	
median (IQR)	0 (0, 1)	0 (0, 1)	1 (0, 1)	0.42

Table S11- Responses to components of PROMIS Global Physical Health by hospitalization status				
Question	All N (111)	Not- Hospitalized N (78)	Hospitalized N (33)	p-value
How would you rate your pain on average? 0 is no pain and 10 is the worst imaginable pain				
median (IQR)	1 (0, 3)	0 (0, 3)	3 (0, 5) (n=30)	0.003
In general, how would you rate your physical health?				
A) Poor	10 (9.0%)	5 (6%)	5 (15%)	0.28
B) Fair	26 (23.4%)	20 (26%)	6 (18%)	
C) Good	48 (43.2%)	32 (41%)	16 (48%)	
D) Very good	20 (18.0%)	17 (22%)	3 (9%)	
E) Excellent	5 (4.5%)	4 (5%)	1 (3%)	
No answer	2 (1.8%)	0 (0%)	2 (6%)	
median (IQR)	3 (2, 3)	3 (2, 4)	3 (2, 3)	0.21
To what extent are you able to carry out your everyday physical activities such as walking, climbing stairs, carrying groceries, or moving a chair				
A) Not at all	3 (2.7%)	2 (3%)	1 (3%)	0.015
B) A little	9 (8.1%)	4 (5%)	5 (15%)	
C) Moderately	18 (16.2%)	11 (14%)	7 (21%)	
D) Mostly	23 (20.7%)	14 (18%)	9 (27%)	
E) Completely	55 (49.5%)	47 (60%)	8 (24%)	
No answer	3 (2.7%)	0 (0%)	3 (9%)	
median (IQR)	5 (3, 5)	5 (4, 5)	4 (3, 5)	0.002
How would you rate your fatigue on average?				
A) Very severe	3 (2.7%)	0 (0%)	3 (9%)	0.041
B) Severe	8 (7.2%)	4 (5%)	4 (12%)	
C) Moderate	35 (31.5%)	27 (35%)	8 (24%)	
D) Mild	46 (41.4%)	34 (44%)	12 (36%)	
E) None	16 (14.4%)	13 (17%)	3 (9%)	
No answer	3 (2.7%)	0 (0%)	3 (9%)	
median (IQR)	4 (3, 4)	4 (3, 4)	3.5 (3, 4)	0.091

Table S12- Responses to components of Global Physical Health by time from SARS-CoV2 infection				
Question	All N (111)	<6-months N (76)	>6-months N (35)	p-value
How would you rate your pain on average? 0 is no pain and 10 is the worst imaginable pain				
median (IQR)	1 (0, 3)	0 (0, 3)	3 (0, 4)	0.021
In general, how would you rate your physical health?				
A) Poor	10 (9.0%)	5 (7%)	5 (14%)	0.38
B) Fair	26 (23.4%)	19 (25%)	7 (20%)	
C) Good	48 (43.2%)	35 (46%)	13 (37%)	
D) Very good	20 (18.0%)	14 (18%)	6 (17%)	
E) Excellent	5 (4.5%)	2 (3%)	3 (9%)	
No answer	2 (1.8%)	1 (1%)	1 (3%)	
median (IQR)	3 (2, 3)	3 (2, 3)	3 (2, 4)	0.98
To what extent are you able to carry out your everyday physical activities such as walking, climbing stairs, carrying groceries, or moving a chair?				
A) Not at all	3 (2.7%)	2 (3%)	1 (3%)	0.46
B) A little	9 (8.1%)	6 (8%)	3 (9%)	
C) Moderately	18 (16.2%)	10 (13%)	8 (23%)	
D) Mostly	23 (20.7%)	15 (20%)	8 (23%)	
E) Completely	55 (49.5%)	42 (55%)	13 (37%)	
No answer	3 (2.7%)	1 (1%)	2 (6%)	
median (IQR)	5 (3, 5)	5 (4, 5)	4 (3, 5)	0.13
How would you rate your fatigue on average?				
A) Very severe	3 (2.7%)	0 (0%)	3 (9%)	0.14
B) Severe	8 (7.2%)	5 (7%)	3 (9%)	
C) Moderate	35 (31.5%)	26 (34%)	9 (26%)	
D) Mild	46 (41.4%)	33 (43%)	13 (37%)	
E) None	16 (14.4%)	11 (14%)	5 (14%)	
No answer	3 (2.7%)	1 (1%)	2 (6%)	
median (IQR)	4 (3, 4)	4 (3, 4)	4 (3, 4)	0.47

Table S13- Responses to components of BCSS survey by hospitalization status and time from SARS-CoV2 infection							
Question	Overall n (111)	Non- hospitalized n (78)	Hospitalized n (33)	P- value	<6- months n (63)	>6- months N (35)	P- value
How much difficulty did you have breathing today?							
None - unaware of any difficulty	70 (63.1%)	55 (71%)	15 (45%)	0.11	51 (67%)	19 (54%)	0.50
Mild - noticeable when performing strenuous activity (eg, running)	29 (26.1%)	17 (22%)	12 (36%)		18 (24%)	11 (31%)	
Moderate - noticeable even when performing light activity (eg, bed making or carrying groceries)	10 (9.0%)	6 (8%)	4 (12%)		6 (8%)	4 (11%)	
Marked – noticeable when washing or dressing	0	0	0		0	0	
Severe – almost constant, present even when resting	0	0	0		0	0	
No answer	2 (1.8%)	0 (0%)	2 (6%)		1 (1%)	1 (3%)	
median (IQR)	0 (0, 1)	0 (0, 1)	1 (0, 1)	0.035	0 (0, 1)	0 (0, 1)	0.22
How was your cough today?							
No cough - unaware of coughing	75 (67.6%)	59 (76%)	16 (48%)	0.053	55 (72%)	20 (57%)	0.37
Rare - cough now and then	23 (20.7%)	12 (15%)	11 (33%)		14 (18%)	9 (26%)	
Occasional - less than hourly	9 (8.1%)	6 (8%)	3 (9%)		5 (7%)	4 (11%)	
Frequent - one or more times an hour	2 (1.8%)	1 (1%)	1 (3%)		1 (1%)	1 (3%)	
Almost constant – never free of cough or need to cough	0	0	0		0	0	
No answer	2 (1.8%)	0 (0%)	2 (6%)		1 (1%)	1 (3%)	
median (IQR)	0 (0, 1)	0 (0, 0)	0 (0, 1)	0.022	0 (0, 1)	0 (0, 1)	0.12
How much trouble did you have due to sputum today?							
None - unaware of any trouble	93 (83.8%)	68 (87%)	25 (76%)	0.33	67 (88%)	26 (74%)	0.18
Mild - rarely caused trouble	14 (12.6%)	8 (10%)	6 (18%)		7 (9%)	7 (20%)	
Moderate - noticeable trouble	2 (1.8%)	2 (3%)	0 (0%)		1 (1%)	1 (3%)	
Marked – caused a great deal of trouble	0	0	0		0	0	
Severe – almost constant trouble	0	0	0		0	0	

No answer	2 (1.8%)	0 (0%)	2 (6%)		1 (1%)	1 (3%)	
median (IQR)	0 (0, 0)	0 (0, 0)	0 (0, 0)	0.42	0 (0, 0)	0 (0, 0)	0.081

Table S14- Responses to smell, taste, and diarrhea questions by hospitalization status and time from SARS-CoV2 infection

Question	Overall N (111)	Nonhospitalized N (78)	Hospitalized N (33)	P-value	<6- months N (76)	>6- months N (35)	P- value
This question is about your smell sensation today							
I smell things normal	85 (76.6%)	61 (78%)	24 (73%)	0.60	61 (80%)	24 (69%)	0.21
I smell things but my smell sensation is less than before my diagnosis of COVID19	12 (10.8%)	7 (9%)	5 (15%)		5 (7%)	7 (20%)	
I smell things but some things smell different than before my diagnosis of COVID19 or I sometimes smell things that are not there	8 (7.2%)	7 (9%)	1 (3%)		6 (8%)	2 (6%)	
I cannot smell anything	4 (3.6%)	3 (4%)	1 (3%)		3 (4%)	1 (3%)	
No answer	2 (1.8%)	0 (0%)	2 (6%)		1 (1%)	1 (3%)	
This question is about your taste sensation today							
I taste food normal	85 (76.6%)	64 (82%)	21 (64%)	0.23	63 (83%)	22 (63%)	0.049
I taste things but my taste sensation is less than before my diagnosis of COVID19	8 (7.2%)	4 (5%)	4 (12%)		4 (5%)	4 (11%)	
I taste things but some food taste different than before my diagnosis of COVID19 or I sometimes taste things that I did not eat	14 (12.6%)	8 (10%)	6 (18%)		6 (8%)	8 (23%)	
I cannot taste food	2 (1.8%)	2 (3%)	0 (0%)		2 (3%)	0 (0%)	
No answer	2 (1.8%)	0 (0%)	2 (6%)		1 (1%)	1 (3%)	
This question is about your bowel movements over the last 2 weeks							
I have normal bowel movements	72 (64.9%)	52 (67%)	20 (61%)	0.68	49 (64%)	23 (66%)	1.00
I have diarrhea every once a week	10 (9.0%)	8 (10%)	2 (6%)		7 (9%)	3 (9%)	
I have diarrhea every few days	19 (17.1%)	12 (15%)	7 (21%)		13 (17%)	6 (17%)	
I have diarrhea every day	7 (6.3%)	6 (8%)	1 (3%)		5 (7%)	2 (6%)	

No answer	3 (2.7%)	0 (0%)	3 (9%)		2 (3%)	1 (3%)	
-----------	----------	--------	--------	--	--------	--------	--