

Coinvestigators

The following coinvestigators participated in the trial as site investigators:

Shin Hisahara, MD, PhD (Sapporo Medical University Hospital, Hokkaido); Seiji Kikuchi, MD, PhD (National Hospital Organization Hokkaido Medical Center, Hokkaido); Masayuki Baba, MD, PhD (Aomori Prefectural Central Hospital, Aomori); Takashi Abe, MD, PhD (Abe Neurology Clinic, Iwate); Shigeru Sato, MD, PhD (Kohnan Hospital, Miyagi); Etsuro Mori, MD, PhD (Tohoku University Hospital, Miyagi); Ken Nagata, MD, PhD (Research Institute for Brain and Blood Vessels, Akita); Shigeki Arawaka, MD, PhD (Yamagata University Hospital, Yamagata); Akira Tamaoka, MD, PhD (University of Tsukuba Hospital, Ibaraki); Ken-ichi Fujimoto, MD, PhD (Jichi Medical University Hospital, Tochigi); Yasuo Harigaya, MD, PhD (Maebashi Red Cross Hospital, Gunma); Masaki Ikeda, MD, PhD (Gunma University Hospital, Gunma); Hideyoshi Sugimoto, MD, PhD (Saino Clinic, Saitama); Tomoyuki Ohno, MD (Takesato Hospital, Saitama); Koichi Kashiwado, MD, PhD (Kashiwado Hospital, Chiba); Hideji Hashida, MD, PhD (Japanese Red Cross Medical Center, Tokyo); Satoshi Orimo, MD, PhD (Kanto Central Hospital, Tokyo); Taku Hatano, MD, PhD (Juntendo University Hospital, Tokyo); Tadashi Tukamoto, MD, PhD (National Center Hospital, National Center of

Neurology and Psychiatry, Tokyo); Fusako Yokochi, MD, PhD (Tokyo Metropolitan Neurological Hospital, Tokyo); Hideki Takubo, MD, PhD (Ebara Hospital, Tokyo); Chika Nishimura, MD (Kurumi Clinic, Tokyo); Satoe Shimoda, MD, PhD (Ginza Neurological Clinic, Tokyo); Teruo Yokoyama, MD, PhD (National Hospital Organization Sagami National Hospital, Kanagawa); Yuta Manabe, MD, PhD (Yokohama Shintoshin Neurosurgical Hospital, Kanagawa); Hiroaki Hino, MD, PhD (Yokohama Hoya Hospital, Kanagawa); Noriko Kawashima, MD (Kawashima Neurology Clinic, Kanagawa); Takuya Mashima, MD, PhD, and Koki Kikugawa, MD, PhD (Tsubame Rosai Hospital, Niigata); Yasuhiro Kawase, MD (Kawase Neurology Clinic, Niigata); Yoshihisa Takiyama, MD, PhD (University of Yamanashi Hospital, Yamanashi); Yo-ichi Takei, MD, PhD (National Hospital Organization Matsumoto Medical Center, Nagano); Takao Hashimoto, MD, PhD (Aizawa Hospital, Nagano); Yusaku Shimizu, MD, PhD (Ina Central Hospital, Nagano); Yasuyuki Okuma, MD, PhD (Juntendo University Shizuoka Hospital, Shizuoka); Tomokazu Obi, MD, PhD (Shizuoka Institute of Epilepsy and Neurological Disorders, Shizuoka); Tatsuya Hattori, MD, PhD (Honmachi Neurological Clinic, Aichi); Mikiko Kamijo, MD, PhD (Chubu Rosai Hospital, Aichi); Hideyuki Sawada, MD, PhD (National Hospital Organization Utano

Hospital, Kyoto); Ryosuke Takahashi, MD, PhD (Kyoto University Hospital, Kyoto); Masutaro Kanda, MD, PhD (Ijinkai Takeda General Hospital, Kyoto); Yoshihisa Tatsuoka, MD, PhD (Tatsuoka Neurology Clinic, Kyoto); Michiyuki Hayashi, MD, PhD (Hayashi Neurology Clinic, Kyoto); Akihiko Ozaki, MD, PhD (Osaka Saiseikai Nakatsu Hospital, Osaka); Takanori Hazama, MD, PhD (Osaka General Medical Center, Osaka); Kuni Konaka, MD, PhD, and Masaru Yokoe, MD, PhD (Osaka University Hospital, Osaka); Harutoshi Fujimura, MD, PhD (National Hospital Organization Toneyama National Hospital, Osaka); Mitsuo Kodama, MD (Hirakata Kohsai Hospital, Osaka); Tatsushi Toda, MD, PhD (Kobe University Hospital, Hyogo); Kazuo Toda, MD, PhD (Toda Internal Medicine & Rehabilitation Clinic, Hyogo); Chie Yanagihara, MD, PhD, and Yo Nishimura, MD (Nishi-Kobe Medical Center, Hyogo); Kenichi Kashihara, MD, PhD (Okayama Kyokuto Hospital, Okayama); Tomoaki Tsujii, MD, PhD, and Hayato Yabe, MD, PhD (Saiseikai Matsuyama Hospital, Ehime); Masahiro Nagai, MD, PhD (Ehime University Hospital, Ehime); Tomoyo Hashimoto, MD, PhD, and Tomoaki Yuhi, MD, PhD (University of Occupational and Environmental Health Hospital, Fukuoka); Yoshio Tsuboi, MD, PhD (Fukuoka University Hospital, Fukuoka); Mitsuhiro Tsujihata, MD, PhD (Nagasaki Kita Hospital, Nagasaki); Hidenori

Matsuo, MD, PhD (National Hospital Organization Nagasaki Kawatana Medical Center, Nagasaki); Aoi Yoshiiwa, MD, PhD, and Noriyuki Kimura, MD, PhD (Oita University Hospital, Oita); Kazuhito Tsuruta, MD (Junwa Rehabilitation Foundation Junwakai Memorial Hospital, Miyazaki); and Kazunori Okahara, MD (Keimei Memorial Hospital, Miyazaki).