

eTable 1 Descriptive statistics on the continuous outcomes duration (hours) and pain coping (0-10, lower score means less coping) for perimenstrual and non-perimenstrual attacks for women with MRM versus non-MRM

	MRM		Non-MRM	
	Perimenstrual attacks (n=747)	Non-perimenstrual attacks (n=2149)	Perimenstrual attacks (n=183)	Non-perimenstrual attacks (n=1074)
Duration (hours) - incl. 24 h migraine-free, median (IQR)	23.2 (14.0-38.7)	16.9 (10.6-26.6)	13.3 (6.0-31.8)	15.0 (8.6-25.4)
Duration (hours) - incl. 48 h migraine-free, median (IQR)	42.3 (23.0-69.6)	25.2 (14.8-39.5)	20.0 (6.0-41.6)	18.6 (10.2-29.9)
Pain coping score (0-10 scale), mean ± SD	4.9 ± 1.4	5.3 ± 1.2	5.4 ± 1.8	5.5 ± 1.2

Intra-individual means were calculated for perimenstrual attacks and non-perimenstrual attacks prior to group calculations to account for the correlation between migraine attacks within the same participant. Numbers of attacks included in the analyses on duration when migraine-free periods of less than 48 hours were included in the duration were lower than presented in the table (n=661, n=1632, n=170 and n=921 respectively). Note: the adjusted mean difference on pain coping presented in the text is calculated with a linear mixed effects model corrected for potential confounders, and therefore, slightly deviates from the result presented in this table.