

eTable 1. Comparison of the demographic and risk factor profile for patients with EGFRs 1-6 and 7-34 mutations. The p-values were calculated by two-sample t-test. BMI, body mass index; *Significance at p<0.05.

History of Cardiovascular Risk Factor	Whole cohort	Patients with EGFRs 1-6 mutations	Patients with EGFRs 7-34 mutations	p value
n	485	391	94	NA
Age at entry into the study (years (SD))*	50.1 (12.0)	49.1 (11.8)	54.2 (11.8)	0.0002
% male	279 / 485 (57.5%)	225 / 391 (57.5%)	54 / 94 (57.4%)	0.99
Current or ex-smokers	231 / 485 (47.6%)	186 / 391 (47.6%)	45 / 94 (47.9%)	0.96
Alcohol Consumption (>14 units a week)	134 / 364 (36.8%)	113 / 292 (38.7%)	21 / 72 (29.2%)	0.38
BMI (kg/m ² (SD))	28.4 (6.5)	28.0 (6.5)	29.7 (6.5)	0.05
Hypertension	107 / 484 (22.1%)	80 / 390 (20.5%)	27 / 94 (28.7%)	0.11
Hypercholesterolaemia*	224 / 483 (46.4%)	167 / 389 (42.9%)	57 / 94 (60.6%)	0.002
Diabetes mellitus	26 / 485 (5.4%)	19 / 391 (4.9%)	7 / 94 (7.4%)	0.38